



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2014 – 12:41 AM GMT

PDB ID : 2C2C  
Title : REFINEMENT OF THE CRYSTAL STRUCTURE OF OXIDIZED RHO-  
DOSPIRILLUM RUBRUM CYTOCHROME C2  
Authors : Bhatia, G.; Finzel, B.C.; Kraut, J.  
Deposited on : 1983-11-03  
Resolution : 2.00 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	112	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 962 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	112	832	529	137	162	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	ASN	ASP	CONFLICT	UNP P00092
A	73	ASN	ASP	CONFLICT	UNP P00092

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total	O	0	0
			87	87		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CYTOCHROME C2

Chain A: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	32.22Å 37.36Å 84.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.00 42.31 – 1.68	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.00) 99.7 (42.31-1.68)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 1.68Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.172 , (Not available) 0.172 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	12.9	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 49.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 12247 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	962	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	2.35	36/848 (4.2%)	3.52	112/1144 (9.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	3	3

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	LYS	C-O	13.86	1.49	1.23
1	A	50	GLU	CD-OE1	11.93	1.38	1.25
1	A	91	MET	C-O	11.09	1.44	1.23
1	A	1	GLU	CA-CB	9.75	1.75	1.53
1	A	37	GLU	CD-OE1	9.41	1.36	1.25
1	A	1	GLU	C-O	9.10	1.40	1.23
1	A	21	ASP	CG-OD1	8.19	1.44	1.25
1	A	63	THR	CA-CB	7.97	1.74	1.53
1	A	27	LYS	C-O	7.81	1.38	1.23
1	A	9	LYS	CD-CE	6.88	1.68	1.51
1	A	81	LYS	CE-NZ	6.85	1.66	1.49
1	A	8	GLU	CG-CD	6.84	1.62	1.51
1	A	98	ASP	CB-CG	6.74	1.66	1.51
1	A	78	VAL	CB-CG1	-6.56	1.39	1.52
1	A	59	GLY	C-O	6.55	1.34	1.23
1	A	64	GLU	CD-OE1	6.35	1.32	1.25
1	A	29	GLY	CA-C	6.33	1.61	1.51
1	A	90	LYS	CE-NZ	6.30	1.64	1.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	92	THR	N-CA	6.15	1.58	1.46
1	A	9	LYS	CE-NZ	6.12	1.64	1.49
1	A	92	THR	CA-CB	6.10	1.69	1.53
1	A	85	PRO	CA-C	6.08	1.65	1.52
1	A	100	GLU	CD-OE2	6.01	1.32	1.25
1	A	28	VAL	CB-CG2	5.97	1.65	1.52
1	A	99	ASP	CG-OD2	5.96	1.39	1.25
1	A	78	VAL	C-N	-5.82	1.20	1.34
1	A	104	VAL	CB-CG2	-5.73	1.40	1.52
1	A	1	GLU	C-N	-5.71	1.22	1.33
1	A	98	ASP	CA-CB	5.64	1.66	1.53
1	A	63	THR	CB-OG1	5.57	1.54	1.43
1	A	23	GLY	CA-C	-5.45	1.43	1.51
1	A	52	TYR	CE1-CZ	5.33	1.45	1.38
1	A	37	GLU	CD-OE2	5.21	1.31	1.25
1	A	85	PRO	N-CA	5.11	1.55	1.47
1	A	111	LEU	CA-CB	5.01	1.65	1.53
1	A	8	GLU	CD-OE1	-5.01	1.20	1.25

All (112) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	99	ASP	CB-CG-OD2	30.19	145.47	118.30
1	A	98	ASP	CB-CG-OD2	-24.36	96.38	118.30
1	A	98	ASP	CB-CG-OD1	22.29	138.36	118.30
1	A	44	ASP	CB-CG-OD1	19.04	135.43	118.30
1	A	50	GLU	OE1-CD-OE2	18.55	145.56	123.30
1	A	37	GLU	OE1-CD-OE2	17.38	144.15	123.30
1	A	86	LYS	CA-CB-CG	17.25	151.35	113.40
1	A	63	THR	CA-CB-CG2	-16.74	88.97	112.40
1	A	3	ASP	CB-CG-OD2	-14.38	105.36	118.30
1	A	97	LYS	CB-CG-CD	13.69	147.20	111.60
1	A	44	ASP	CB-CG-OD2	-12.90	106.69	118.30
1	A	93	PHE	CB-CG-CD2	-12.75	111.87	120.80
1	A	12	LYS	CB-CG-CD	12.25	143.45	111.60
1	A	78	VAL	CA-CB-CG1	12.04	128.95	110.90
1	A	63	THR	OG1-CB-CG2	-11.55	83.43	110.00
1	A	39	THR	CA-CB-CG2	11.46	128.44	112.40
1	A	85	PRO	O-C-N	11.44	141.00	122.70
1	A	79	LEU	CB-CG-CD2	11.39	130.37	111.00
1	A	1	GLU	O-C-N	11.33	142.46	123.20
1	A	33	PHE	CB-CG-CD1	-11.31	112.88	120.80

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	GLU	C-N-CA	-10.82	99.58	122.30
1	A	112	LYS	CB-CA-C	10.77	131.94	110.40
1	A	70	TYR	CG-CD1-CE1	10.49	129.69	121.30
1	A	99	ASP	OD1-CG-OD2	-10.44	103.47	123.30
1	A	84	ASP	CB-CG-OD2	10.02	127.32	118.30
1	A	54	GLU	OE1-CD-OE2	-9.87	111.46	123.30
1	A	48	TYR	CB-CG-CD2	-9.15	115.51	121.00
1	A	38	ASN	CB-CG-OD1	8.54	138.68	121.60
1	A	9	LYS	CD-CE-NZ	-8.46	92.23	111.70
1	A	99	ASP	CB-CG-OD1	-8.38	110.75	118.30
1	A	57	ALA	O-C-N	-8.38	109.29	122.70
1	A	1	GLU	CA-C-O	-8.34	102.59	120.10
1	A	112	LYS	CA-C-O	8.28	137.49	120.10
1	A	3	ASP	CB-CG-OD1	8.28	125.75	118.30
1	A	54	GLU	CA-C-N	8.18	135.19	117.20
1	A	93	PHE	CB-CG-CD1	7.79	126.26	120.80
1	A	15	LEU	CB-CG-CD1	7.64	123.99	111.00
1	A	79	LEU	CA-CB-CG	7.54	132.65	115.30
1	A	64	GLU	OE1-CD-OE2	7.52	132.33	123.30
1	A	21	ASP	CB-CG-OD2	7.34	124.91	118.30
1	A	40	ALA	N-CA-CB	-7.33	99.84	110.10
1	A	63	THR	CA-CB-OG1	-7.30	93.67	109.00
1	A	54	GLU	CA-C-O	-7.20	104.98	120.10
1	A	80	GLU	CA-CB-CG	7.09	129.00	113.40
1	A	97	LYS	CD-CE-NZ	-7.07	95.44	111.70
1	A	64	GLU	CA-CB-CG	7.06	128.93	113.40
1	A	86	LYS	CB-CA-C	6.86	124.11	110.40
1	A	12	LYS	CA-CB-CG	6.81	128.38	113.40
1	A	92	THR	CA-C-O	-6.66	106.11	120.10
1	A	37	GLU	CG-CD-OE2	-6.62	105.06	118.30
1	A	75	LYS	N-CA-CB	6.61	122.50	110.60
1	A	84	ASP	CB-CG-OD1	-6.61	112.35	118.30
1	A	100	GLU	OE1-CD-OE2	-6.57	115.42	123.30
1	A	92	THR	CA-C-N	6.54	131.59	117.20
1	A	65	ALA	N-CA-CB	-6.52	100.97	110.10
1	A	88	LYS	O-C-N	-6.46	112.36	122.70
1	A	90	LYS	CB-CG-CD	6.33	128.06	111.60
1	A	64	GLU	CG-CD-OE2	-6.32	105.66	118.30
1	A	54	GLU	CG-CD-OE1	6.27	130.85	118.30
1	A	78	VAL	CA-C-O	-6.26	106.96	120.10
1	A	72	LYS	CB-CA-C	-6.25	97.91	110.40
1	A	53	THR	N-CA-CB	-6.25	98.44	110.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	THR	N-CA-CB	6.23	122.14	110.30
1	A	70	TYR	CD1-CG-CD2	-6.20	111.08	117.90
1	A	70	TYR	CB-CG-CD1	6.19	124.71	121.00
1	A	72	LYS	N-CA-CB	6.14	121.66	110.60
1	A	104	VAL	CA-CB-CG1	-6.07	101.80	110.90
1	A	23	GLY	O-C-N	-6.04	112.93	123.20
1	A	62	TRP	O-C-N	-6.01	113.08	122.70
1	A	78	VAL	CG1-CB-CG2	5.99	120.49	110.90
1	A	57	ALA	C-N-CA	5.98	136.65	121.70
1	A	77	PHE	CB-CG-CD1	5.94	124.96	120.80
1	A	85	PRO	CA-N-CD	-5.93	103.20	111.50
1	A	3	ASP	CA-C-N	5.84	130.05	117.20
1	A	95	LEU	CA-CB-CG	5.79	128.61	115.30
1	A	48	TYR	CG-CD1-CE1	-5.77	116.68	121.30
1	A	50	GLU	CG-CD-OE2	-5.76	106.79	118.30
1	A	36	PHE	CZ-CE2-CD2	-5.75	113.20	120.10
1	A	8	GLU	CG-CD-OE2	-5.74	106.81	118.30
1	A	96	THR	CA-C-N	5.74	129.82	117.20
1	A	77	PHE	CG-CD1-CE1	5.72	127.09	120.80
1	A	50	GLU	CB-CG-CD	5.72	129.64	114.20
1	A	52	TYR	CB-CG-CD1	5.68	124.41	121.00
1	A	53	THR	CA-CB-OG1	-5.66	97.11	109.00
1	A	82	SER	CA-C-O	-5.56	108.42	120.10
1	A	51	SER	O-C-N	-5.55	113.81	122.70
1	A	62	TRP	CG-CD1-NE1	-5.53	104.57	110.10
1	A	72	LYS	CD-CE-NZ	5.52	124.40	111.70
1	A	50	GLU	CG-CD-OE1	-5.52	107.26	118.30
1	A	3	ASP	CA-CB-CG	-5.49	101.31	113.40
1	A	84	ASP	C-N-CD	5.49	139.92	128.40
1	A	21	ASP	OD1-CG-OD2	-5.48	112.88	123.30
1	A	33	PHE	CD1-CG-CD2	5.44	125.38	118.30
1	A	91	MET	C-N-CA	-5.44	108.11	121.70
1	A	1	GLU	CB-CA-C	-5.42	99.56	110.40
1	A	100	GLU	CG-CD-OE1	5.41	129.12	118.30
1	A	20	PHE	CD1-CE1-CZ	5.38	126.56	120.10
1	A	76	ALA	CB-CA-C	5.38	118.17	110.10
1	A	36	PHE	CD1-CE1-CZ	-5.32	113.71	120.10
1	A	85	PRO	C-N-CA	-5.30	108.45	121.70
1	A	68	ALA	CB-CA-C	5.28	118.01	110.10
1	A	15	LEU	CA-CB-CG	5.25	127.38	115.30
1	A	62	TRP	CD2-CE3-CZ3	-5.24	111.99	118.80
1	A	36	PHE	CB-CG-CD1	-5.24	117.13	120.80

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	70	TYR	CD1-CE1-CZ	-5.20	115.12	119.80
1	A	36	PHE	CE1-CZ-CE2	5.16	129.28	120.00
1	A	69	ALA	N-CA-CB	-5.11	102.94	110.10
1	A	94	LYS	CB-CA-C	-5.10	100.19	110.40
1	A	8	GLU	OE1-CD-OE2	5.09	129.41	123.30
1	A	85	PRO	CA-C-O	-5.08	108.00	120.20
1	A	43	LYS	O-C-N	-5.08	114.57	122.70
1	A	60	LEU	CB-CG-CD2	-5.03	102.45	111.00

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	39	THR	CB
1	A	110	THR	CB
1	A	112	LYS	CA

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	27	LYS	Mainchain
1	A	68	ALA	Mainchain
1	A	77	PHE	Mainchain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	832	0	795	24	11
2	A	43	0	30	8	0
3	A	87	0	0	5	11
All	All	962	0	825	24	11

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

All (24) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:1:GLU:CA	1:A:1:GLU:CB	1.75	1.60
1:A:17:CYS:SG	2:A:113:HEM:CAC	2.13	1.35
1:A:88:LYS:CB	3:A:185:HOH:O	1.95	1.14
1:A:17:CYS:HG	2:A:113:HEM:CAC	1.64	1.05
1:A:17:CYS:SG	2:A:113:HEM:HAC	2.00	0.97
1:A:100:GLU:O	1:A:104:VAL:HG23	1.71	0.88
1:A:73:ASN:OD1	3:A:201:HOH:O	2.00	0.79
1:A:17:CYS:SG	2:A:113:HEM:CBC	2.73	0.76
1:A:17:CYS:HG	2:A:113:HEM:CBC	2.00	0.74
1:A:1:GLU:C	1:A:1:GLU:CB	2.56	0.73
1:A:17:CYS:SG	2:A:113:HEM:C3C	2.83	0.71
1:A:42:HIS:ND1	2:A:113:HEM:O2A	2.25	0.65
1:A:1:GLU:N	1:A:1:GLU:CB	2.61	0.62
1:A:63:THR:HG22	3:A:141:HOH:O	2.03	0.59
1:A:39:THR:HG21	1:A:56:LYS:HD2	1.85	0.58
1:A:39:THR:HG23	1:A:40:ALA:O	2.04	0.57
1:A:107:TYR:O	1:A:110:THR:HG22	2.05	0.56
1:A:64:GLU:HG3	1:A:105:ILE:HG21	1.88	0.56
1:A:63:THR:CG2	3:A:141:HOH:O	2.59	0.50
1:A:45:ASN:HD22	1:A:45:ASN:H	1.63	0.45
1:A:12:LYS:CD	3:A:205:HOH:O	2.66	0.43
1:A:35:VAL:HG23	1:A:42:HIS:CE1	2.54	0.43
1:A:68:ALA:O	1:A:72:LYS:HE2	2.20	0.40
1:A:91:MET:HB2	2:A:113:HEM:C1D	2.56	0.40

All (11) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:90:LYS:CE	3:A:134:HOH:O[4_556]	1.20	1.00
1:A:63:THR:CB	3:A:203:HOH:O[3_745]	1.29	0.91
1:A:90:LYS:NZ	3:A:134:HOH:O[4_556]	1.32	0.88
1:A:112:LYS:CB	3:A:181:HOH:O[4_456]	1.43	0.77
1:A:64:GLU:OE2	3:A:200:HOH:O[3_745]	1.47	0.73
1:A:112:LYS:CB	3:A:182:HOH:O[4_456]	1.50	0.70
1:A:64:GLU:CB	3:A:200:HOH:O[3_745]	1.67	0.53
1:A:64:GLU:CG	3:A:200:HOH:O[3_745]	1.87	0.33
1:A:64:GLU:CD	3:A:200:HOH:O[3_745]	2.00	0.20
1:A:21:ASP:CB	3:A:183:HOH:O[4_456]	2.05	0.15
1:A:21:ASP:CB	3:A:194:HOH:O[4_456]	2.08	0.12

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	110/112 (98%)	103 (94%)	7 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/89 (90%)	75 (94%)	5 (6%)	25	18

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	45	ASN
1	A	79	LEU
1	A	97	LYS
1	A	104	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	45	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	113	1	49,50,50	4.34	19 (38%)	46,82,82	2.53	16 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	113	1	-	0/14/114/114	0/0/8/8

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	113	HEM	C2B-C1B	-18.06	1.40	1.44
2	A	113	HEM	C2D-C1D	-17.23	1.40	1.44
2	A	113	HEM	CAA-C2A	6.64	1.63	1.52
2	A	113	HEM	C3C-C2C	-5.94	1.33	1.43
2	A	113	HEM	CHB-C1B	5.25	1.43	1.35
2	A	113	HEM	CHA-C4D	4.59	1.42	1.35
2	A	113	HEM	C3D-C2D	-4.11	1.36	1.43
2	A	113	HEM	C3B-C4B	-4.08	1.39	1.44

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	113	HEM	C3B-CAB	3.81	1.52	1.40
2	A	113	HEM	C2C-C1C	3.70	1.54	1.43
2	A	113	HEM	CMA-C3A	3.53	1.59	1.51
2	A	113	HEM	CBB-CAB	3.31	1.48	1.28
2	A	113	HEM	C3C-CAC	3.25	1.50	1.40
2	A	113	HEM	CMD-C2D	3.25	1.57	1.47
2	A	113	HEM	CMB-C2B	3.20	1.57	1.47
2	A	113	HEM	C3D-C4D	-2.58	1.43	1.44
2	A	113	HEM	FE-NA	2.56	2.03	1.92
2	A	113	HEM	C2A-C3A	-2.45	1.30	1.37
2	A	113	HEM	C4A-C3A	2.28	1.43	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	113	HEM	CHC-C1C-NC	6.82	130.66	124.73
2	A	113	HEM	CHD-C1D-ND	-5.59	119.94	124.58
2	A	113	HEM	C4C-NC-C1C	5.38	111.13	105.53
2	A	113	HEM	C4D-ND-C1D	-4.83	100.22	105.16
2	A	113	HEM	C1A-CHA-C4D	-4.79	121.16	127.47
2	A	113	HEM	C1B-NB-C4B	-4.36	100.70	105.16
2	A	113	HEM	C4A-CHB-C1B	-4.03	122.16	127.47
2	A	113	HEM	C3B-C4B-NB	3.80	116.72	114.00
2	A	113	HEM	CBD-CAD-C3D	-2.68	108.53	114.37
2	A	113	HEM	O2A-CGA-O1A	2.58	129.87	123.30
2	A	113	HEM	CHD-C4C-NC	2.58	126.98	124.73
2	A	113	HEM	CAD-CBD-CGD	2.58	121.54	113.48
2	A	113	HEM	O1D-CGD-CBD	-2.52	114.36	123.03
2	A	113	HEM	CMA-C3A-C4A	-2.50	124.77	128.62
2	A	113	HEM	CMC-C2C-C3C	2.39	131.80	126.16
2	A	113	HEM	CAD-C3D-C4D	2.17	128.44	124.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	112/112 (100%)	-0.88	1 (0%) 81 82	7, 14, 25, 51	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	GLU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	HEM	A	113	43/43	0.05	-0.45	3,6,12,17	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.