



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 04:34 PM GMT

PDB ID : 1C3V
Title : DIHYDRODIPICOLINATE REDUCTASE FROM MYCOBACTERIUM TUBERCULOSIS COMPLEXED WITH NADPH AND PDC
Authors : Cirilli, M.; Zheng, R.; Scapin, G.; Blanchard, J.S.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 1999-07-28
Resolution : 2.39 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2207 (2.40-2.40)
Clashscore	79885	2789 (2.40-2.40)
Ramachandran outliers	78287	2736 (2.40-2.40)
Sidechain outliers	78261	2737 (2.40-2.40)
RSRZ outliers	66119	2210 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	245	
1	B	245	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 3778 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIHYDRODIPICOLINATE REDUCTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1808	1141	321	342	4			
1	B	245	Total	C	N	O	S	0	0	0
			1812	1144	322	342	4			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is PYRIDINE-2,6-DICARBOXYLICACID (three-letter code: PDC) (formula: $C_7H_5NO_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			12	7	1	4		
3	B	1	Total	C	N	O	0	0
			12	7	1	4		

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	8	Total 8	O 8	0	0
5	B	17	Total 17	O 17	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DIHYDRODIPICOLINATE REDUCTASE



• Molecule 1: DIHYDRODIPICOLINATE REDUCTASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	118.83Å 118.26Å 79.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.40 – 2.39 30.40 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.40-2.39) 79.4 (30.40-2.50)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.51Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.193 , 0.243 0.198 , 0.252	Depositor DCC
R_{free} test set	649 reflections (4.08%)	DCC
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.146	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.6	EDS
Estimated twinning fraction	0.189 for -k,-h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 15888 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3778	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, PG4, PDC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/1843	0.86	3/2514 (0.1%)
1	B	0.32	0/1847	0.90	3/2518 (0.1%)
All	All	0.32	0/3690	0.88	6/5032 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1083	ARG	NE-CZ-NH2	-22.34	109.13	120.30
1	B	1083	ARG	NE-CZ-NH1	21.12	130.86	120.30
1	A	583	ARG	NE-CZ-NH1	-20.60	110.00	120.30
1	A	583	ARG	NE-CZ-NH2	19.30	129.95	120.30
1	B	1083	ARG	CD-NE-CZ	14.96	144.54	123.60
1	A	583	ARG	CD-NE-CZ	14.37	143.72	123.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1808	0	1815	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1812	0	1826	33	0
2	A	48	0	25	2	0
2	B	48	0	25	2	0
3	A	12	0	5	0	0
3	B	12	0	5	0	0
4	B	13	0	18	1	0
5	A	8	0	0	0	0
5	B	17	0	0	1	0
All	All	3778	0	3719	68	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (68) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:591:LEU:HD22	1:A:733:GLY:HA2	1.68	0.75
1:A:601:ILE:HD12	1:A:734:LEU:HD11	1.71	0.72
1:B:1011:LYS:HE3	2:B:1301:NDP:O1A	1.93	0.68
1:A:542:THR:HG21	1:A:568:ASN:HD22	1.62	0.64
1:B:1042:THR:HG21	1:B:1068:ASN:HD22	1.63	0.64
1:B:1075:GLY:O	2:B:1301:NDP:H2N	2.00	0.62
1:B:1017:VAL:HG13	1:B:1028:LEU:HD11	1.83	0.61
1:A:517:VAL:HG13	1:A:528:LEU:HD11	1.83	0.61
1:B:1054:HIS:HD2	1:B:1056:ASP:H	1.52	0.58
1:B:1208:ARG:HH22	4:B:2000:PG4:H72	1.69	0.58
1:A:554:HIS:HD2	1:A:556:ASP:H	1.51	0.57
1:A:542:THR:CG2	1:A:568:ASN:HD22	2.18	0.57
1:B:1042:THR:CG2	1:B:1068:ASN:HD22	2.18	0.56
1:B:1028:LEU:HD21	1:B:1031:GLU:HG3	1.88	0.56
1:B:1088:GLU:O	1:B:1092:VAL:HG23	2.06	0.55
1:B:1055:PRO:O	1:B:1083:ARG:NH2	2.39	0.55
1:B:1085:GLN:O	1:B:1088:GLU:HB2	2.08	0.54
1:A:528:LEU:HD21	1:A:531:GLU:HG3	1.88	0.53
1:A:510:GLY:HA3	2:A:801:NDP:H52A	1.89	0.53
1:B:1218:VAL:HB	1:B:1219:PRO:HD3	1.91	0.52
1:B:1066:ILE:HG23	1:B:1091:LEU:HD23	1.90	0.52
1:A:718:VAL:HB	1:A:719:PRO:HD3	1.92	0.52
1:A:579:PHE:CE2	1:A:601:ILE:HD13	2.46	0.51
1:A:601:ILE:O	1:A:736:VAL:HA	2.10	0.50
1:B:1079:PHE:CE2	1:B:1101:ILE:HD13	2.46	0.50
1:A:542:THR:HG21	1:A:568:ASN:ND2	2.27	0.49
1:B:1042:THR:HG21	1:B:1068:ASN:ND2	2.28	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:555:PRO:O	1:A:583:ARG:NH2	2.46	0.48
1:A:691:VAL:HB	1:A:711:SER:O	2.14	0.47
1:B:1153:GLU:HG2	1:B:1156:LYS:NZ	2.29	0.47
1:A:575:GLY:O	2:A:801:NDP:H2N	2.13	0.47
1:A:555:PRO:HA	1:A:558:VAL:HG23	1.96	0.47
1:B:1055:PRO:HA	1:B:1058:VAL:HG23	1.96	0.47
1:B:1214:ARG:HG3	1:B:1214:ARG:HH11	1.80	0.47
1:A:714:ARG:HG3	1:A:714:ARG:HH11	1.80	0.47
1:A:653:GLU:HG2	1:A:656:LYS:NZ	2.30	0.46
1:A:554:HIS:CD2	1:A:556:ASP:H	2.33	0.46
1:B:1191:VAL:HB	1:B:1211:SER:O	2.16	0.46
1:A:559:MET:HG2	1:A:583:ARG:HH21	1.80	0.45
1:A:585:GLN:O	1:A:588:GLU:HB2	2.17	0.45
1:A:592:VAL:O	1:A:595:PRO:HD3	2.18	0.44
1:A:559:MET:HE3	1:A:583:ARG:HB3	1.99	0.44
1:A:728:ILE:HG23	1:A:729:ALA:N	2.33	0.44
1:B:1228:ILE:HG23	1:B:1229:ALA:N	2.33	0.43
1:B:1227:ARG:O	1:B:1230:GLU:HB2	2.18	0.43
1:B:1001:MET:N	5:B:2010:HOH:O	2.51	0.43
1:B:1049:VAL:HG11	1:B:1065:LEU:HD22	2.01	0.43
1:B:1239:GLU:HB2	1:B:1240:PRO:HD3	1.99	0.43
1:A:727:ARG:O	1:A:730:GLU:HB2	2.19	0.43
1:A:723:LEU:HG	1:A:741:LEU:HD23	2.00	0.42
1:A:739:GLU:N	1:A:740:PRO:CD	2.82	0.42
1:A:599:VAL:O	1:A:734:LEU:HA	2.20	0.42
1:B:1192:ALA:HB3	1:B:1211:SER:HB2	2.01	0.42
1:B:1239:GLU:H	1:B:1239:GLU:CD	2.22	0.42
1:A:692:ALA:HB3	1:A:711:SER:HB2	2.01	0.42
1:A:549:VAL:HG11	1:A:565:LEU:HD22	2.01	0.41
1:A:627:GLU:HG2	1:A:681:PRO:HD2	2.02	0.41
1:B:1054:HIS:CD2	1:B:1056:ASP:H	2.33	0.41
1:A:549:VAL:HG11	1:A:565:LEU:CD2	2.50	0.41
1:B:1144:ALA:HA	1:B:1194:GLN:NE2	2.35	0.41
1:B:1049:VAL:HG11	1:B:1065:LEU:CD2	2.50	0.41
1:B:1238:LEU:HD12	1:B:1238:LEU:HA	1.96	0.41
1:B:1127:GLU:HG2	1:B:1181:PRO:HD2	2.02	0.41
1:A:644:ALA:HA	1:A:694:GLN:NE2	2.35	0.41
1:A:735:THR:O	1:A:736:VAL:HB	2.21	0.41
1:A:623:PHE:CG	1:A:698:PHE:HB3	2.56	0.41
1:A:607:ILE:HG22	1:B:1203:GLU:OE2	2.20	0.40
1:B:1083:ARG:O	1:B:1087:VAL:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/245 (99%)	223 (92%)	19 (8%)	1 (0%)	43	61
1	B	243/245 (99%)	229 (94%)	13 (5%)	1 (0%)	43	61
All	All	486/490 (99%)	452 (93%)	32 (7%)	2 (0%)	43	61

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	736	VAL
1	B	1095	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/191 (99%)	168 (89%)	21 (11%)	9	12
1	B	190/191 (100%)	171 (90%)	19 (10%)	11	16
All	All	379/382 (99%)	339 (89%)	40 (11%)	10	14

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	545	ASN
1	A	547	GLU
1	A	568	ASN
1	A	582	GLU
1	A	586	GLN
1	A	592	VAL
1	A	596	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	617	LYS
1	A	624	ASP
1	A	630	GLU
1	A	646	ARG
1	A	656	LYS
1	A	668	SER
1	A	687	LEU
1	A	708	ARG
1	A	711	SER
1	A	727	ARG
1	A	738	LEU
1	A	742	LEU
1	A	743	ASP
1	A	745	HIS
1	B	1045	ASN
1	B	1047	GLU
1	B	1068	ASN
1	B	1082	GLU
1	B	1089	SER
1	B	1090	TRP
1	B	1094	LYS
1	B	1096	ASN
1	B	1117	LYS
1	B	1124	ASP
1	B	1130	GLU
1	B	1146	ARG
1	B	1156	LYS
1	B	1168	SER
1	B	1187	LEU
1	B	1208	ARG
1	B	1211	SER
1	B	1227	ARG
1	B	1244	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	554	HIS
1	A	618	GLN
1	A	694	GLN
1	B	1054	HIS
1	B	1086	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1118	GLN
1	B	1194	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	A	801	-	52,52,52	2.09	9 (17%)	80,80,80	2.83	21 (26%)
3	PDC	A	802	-	12,12,12	4.69	9 (75%)	16,16,16	1.88	6 (37%)
2	NDP	B	1301	-	52,52,52	2.09	9 (17%)	80,80,80	2.72	22 (27%)
3	PDC	B	1302	-	12,12,12	4.76	9 (75%)	16,16,16	1.81	6 (37%)
4	PG4	B	2000	-	12,12,12	0.67	0	11,11,11	1.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	801	-	1/1/14/17	0/35/77/77	0/3/5/5
3	PDC	A	802	-	-	0/8/8/8	0/1/1/1
2	NDP	B	1301	-	1/1/14/17	0/35/77/77	0/3/5/5
3	PDC	B	1302	-	-	0/8/8/8	0/1/1/1
4	PG4	B	2000	-	-	0/10/10/10	0/0/0/0

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1302	PDC	C2-C7	-9.32	1.38	1.50
3	A	802	PDC	C2-C7	-9.25	1.38	1.50
2	B	1301	NDP	O7N-C7N	7.50	1.44	1.24
3	B	1302	PDC	C6-C8	-7.36	1.40	1.50
2	A	801	NDP	O7N-C7N	7.34	1.43	1.24
2	A	801	NDP	C4N-C3N	-6.85	1.36	1.50
3	A	802	PDC	C6-C8	-6.71	1.41	1.50
2	B	1301	NDP	C4N-C3N	-6.65	1.37	1.50
3	B	1302	PDC	C2-N1	6.33	1.44	1.34
3	A	802	PDC	C2-N1	6.26	1.44	1.34
3	A	802	PDC	C6-N1	5.67	1.43	1.34
3	B	1302	PDC	C6-N1	5.41	1.42	1.34
3	A	802	PDC	O3-C8	4.68	1.37	1.23
2	B	1301	NDP	C4A-N9A	-4.58	1.31	1.37
3	B	1302	PDC	O3-C8	4.55	1.37	1.23
3	B	1302	PDC	O1-C7	4.32	1.36	1.23
3	A	802	PDC	O1-C7	4.30	1.36	1.23
2	A	801	NDP	C4N-C5N	-4.14	1.40	1.49
2	B	1301	NDP	C4N-C5N	-4.06	1.40	1.49
2	A	801	NDP	C4A-N9A	-4.03	1.31	1.37
2	B	1301	NDP	C2B-C1B	-3.81	1.46	1.52
2	A	801	NDP	C2B-C1B	-3.74	1.46	1.52
2	A	801	NDP	C7N-C3N	-3.70	1.38	1.47
2	A	801	NDP	C6N-C5N	3.67	1.41	1.33
2	B	1301	NDP	C6N-C5N	3.61	1.41	1.33
2	B	1301	NDP	C5A-C4A	-3.48	1.32	1.40
2	A	801	NDP	C5A-C4A	-3.36	1.32	1.40
2	B	1301	NDP	C7N-C3N	-3.36	1.39	1.47
3	B	1302	PDC	C5-C6	2.39	1.43	1.39
3	A	802	PDC	C5-C6	2.39	1.43	1.39
3	B	1302	PDC	O2-C7	2.37	1.37	1.30
3	A	802	PDC	O4-C8	2.36	1.37	1.30
2	B	1301	NDP	C3B-C4B	2.21	1.59	1.53
2	A	801	NDP	C3B-C4B	2.17	1.59	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	802	PDC	O2-C7	2.15	1.36	1.30
3	B	1302	PDC	O4-C8	2.15	1.36	1.30

All (55) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	NDP	O4B-C1B-N9A	14.44	121.87	108.44
2	B	1301	NDP	O4B-C1B-N9A	12.74	120.29	108.44
2	A	801	NDP	C1B-N9A-C4A	9.94	143.81	126.64
2	B	1301	NDP	C1B-N9A-C4A	9.38	142.84	126.64
2	B	1301	NDP	C4N-C3N-C2N	-7.76	112.23	121.68
2	A	801	NDP	C4N-C3N-C2N	-7.61	112.41	121.68
2	B	1301	NDP	C8A-N9A-C1B	-7.05	112.50	126.38
2	A	801	NDP	C8A-N9A-C1B	-7.00	112.59	126.38
2	A	801	NDP	C5N-C4N-C3N	5.01	126.18	112.60
2	B	1301	NDP	C5N-C4N-C3N	4.99	126.11	112.60
2	A	801	NDP	C8A-N9A-C4A	-4.99	103.09	106.90
2	B	1301	NDP	C8A-N9A-C4A	-4.59	103.39	106.90
2	B	1301	NDP	N3A-C2A-N1A	-4.13	125.26	128.71
2	B	1301	NDP	C2B-C1B-N9A	3.91	128.58	113.74
2	B	1301	NDP	O2B-C2B-C1B	3.88	124.71	110.36
2	A	801	NDP	O2B-C2B-C1B	3.73	124.16	110.36
3	B	1302	PDC	C5-C6-N1	-3.71	118.36	122.92
2	A	801	NDP	C2B-C1B-N9A	3.71	127.79	113.74
3	A	802	PDC	C5-C6-N1	-3.67	118.42	122.92
2	A	801	NDP	C4A-C5A-N7A	3.63	112.63	109.52
2	B	1301	NDP	O4B-C1B-C2B	3.54	110.25	106.95
2	A	801	NDP	N3A-C2A-N1A	-3.45	125.82	128.71
2	B	1301	NDP	C4A-C5A-N7A	3.42	112.45	109.52
2	A	801	NDP	O5D-C5D-C4D	3.37	121.30	108.94
2	A	801	NDP	O5B-C5B-C4B	3.28	120.99	108.94
2	B	1301	NDP	O5B-C5B-C4B	3.06	120.17	108.94
2	B	1301	NDP	C4N-C5N-C6N	-3.05	117.02	122.61
2	A	801	NDP	C4N-C5N-C6N	-3.02	117.07	122.61
2	B	1301	NDP	O5D-C5D-C4D	2.91	119.62	108.94
2	A	801	NDP	O3B-C3B-C2B	-2.79	102.98	111.20
2	B	1301	NDP	O3B-C3B-C2B	-2.77	103.05	111.20
2	B	1301	NDP	C5N-C6N-N1N	-2.76	117.39	123.03
2	A	801	NDP	C5N-C6N-N1N	-2.72	117.47	123.03
2	A	801	NDP	C2N-C3N-C7N	2.65	125.27	118.49
3	A	802	PDC	C3-C2-N1	-2.64	119.67	122.92
2	B	1301	NDP	O3B-C3B-C4B	-2.63	103.32	111.08

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	NDP	O3B-C3B-C4B	-2.59	103.44	111.08
3	A	802	PDC	O2-C7-C2	2.46	120.30	114.68
3	B	1302	PDC	O2-C7-C2	2.45	120.28	114.68
2	A	801	NDP	O2B-C2B-C3B	2.44	121.09	111.54
3	A	802	PDC	O4-C8-C6	2.43	120.23	114.68
2	B	1301	NDP	C2N-C3N-C7N	2.41	124.64	118.49
2	B	1301	NDP	O2B-C2B-C3B	2.38	120.87	111.54
3	B	1302	PDC	C3-C2-N1	-2.36	120.02	122.92
2	A	801	NDP	O4B-C1B-C2B	2.29	109.09	106.95
3	B	1302	PDC	C6-N1-C2	2.29	120.74	117.53
2	A	801	NDP	C6A-C5A-C4A	2.27	121.41	117.25
3	A	802	PDC	C6-N1-C2	2.26	120.70	117.53
3	A	802	PDC	O4-C8-O3	-2.19	118.37	123.35
3	B	1302	PDC	O4-C8-C6	2.18	119.67	114.68
2	B	1301	NDP	C6A-C5A-C4A	2.18	121.25	117.25
2	B	1301	NDP	C3N-C2N-N1N	2.12	126.05	123.05
2	B	1301	NDP	C2A-N1A-C6A	-2.09	115.00	118.77
2	A	801	NDP	C3N-C2N-N1N	2.05	125.96	123.05
3	B	1302	PDC	O4-C8-O3	-2.02	118.75	123.35

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	801	NDP	C1B
2	B	1301	NDP	C1B

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/245 (100%)	0.34	14 (5%) 23 21	13, 34, 75, 97	0
1	B	245/245 (100%)	0.26	10 (4%) 35 33	13, 34, 75, 98	0
All	All	490/490 (100%)	0.30	24 (4%) 29 26	13, 34, 76, 98	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	745	HIS	8.8
1	B	1245	HIS	4.9
1	A	543	ASP	4.6
1	A	587	VAL	3.9
1	B	1014	THR	3.2
1	B	1035	GLY	3.0
1	B	1244	LEU	2.9
1	B	1090	TRP	2.8
1	A	586	GLN	2.5
1	B	1243	ASP	2.5
1	A	744	LEU	2.4
1	A	514	THR	2.4
1	A	564	PHE	2.3
1	A	558	VAL	2.3
1	B	1092	VAL	2.3
1	A	580	THR	2.3
1	B	1091	LEU	2.2
1	A	568	ASN	2.2
1	A	582	GLU	2.1
1	B	1032	LEU	2.1
1	A	596	ASN	2.1
1	A	733	GLY	2.0
1	B	1042	THR	2.0
1	A	591	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	PG4	B	2000	13/13	0.15	1.67	24,28,29,32	0
3	PDC	A	802	12/12	0.13	0.55	15,18,24,26	0
3	PDC	B	1302	12/12	0.12	0.01	15,19,20,23	0
2	NDP	A	801	48/48	0.13	-0.45	25,45,63,63	0
2	NDP	B	1301	48/48	0.12	-0.64	27,37,59,61	0

6.5 Other polymers ⓘ

There are no such residues in this entry.