



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 12:34 PM GMT

PDB ID : 1C4G
Title : PHOSPHOGLUCOMUTASE VANADATE BASED TRANSITION STATE
ANALOG COMPLEX
Authors : Baranidharan, S.; Ray Jr., W.J.
Deposited on : 1999-08-24
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	561	
1	B	561	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8877 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROTEIN (ALPHA-D-GLUCOSE 1-PHOSPHATE PHOSPHOGLUCOMUTASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	561	Total	C	N	O	S	0	0	0
			4329	2753	743	817	16			
1	B	561	Total	C	N	O	S	0	0	0
			4329	2753	743	817	16			

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Co	0	0
			1	1		

- Molecule 3 is ALPHA-D-GLUCOSE-1-PHOSPHATE-6-VANADATE (three-letter code: VG1) (formula: C₆H₁₃O₁₂PV).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	O	P	V	0	0
			20	6	12	1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	84	Total	O	0	0
			84	84		
4	B	114	Total	O	0	0
			114	114		



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	174.42Å 174.42Å 101.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.70	Depositor
% Data completeness (in resolution range)	85.9 (6.00-2.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.192 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8877	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CO, VG1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.22	4/4416 (0.1%)	1.06	16/5969 (0.3%)
1	B	1.11	3/4416 (0.1%)	1.05	17/5969 (0.3%)
All	All	1.16	7/8832 (0.1%)	1.05	33/11938 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	416	TRP	CG-CD2	-5.72	1.33	1.43
1	A	488	GLY	N-CA	5.65	1.54	1.46
1	A	239	GLU	CD-OE1	-5.64	1.19	1.25
1	B	392	TRP	CG-CD2	-5.24	1.34	1.43
1	A	445	GLU	CG-CD	5.19	1.59	1.51

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	397	TRP	CD1-CG-CD2	13.96	117.47	106.30
1	B	358	TRP	CD1-CG-CD2	13.45	117.06	106.30
1	A	416	TRP	CD1-CG-CD2	13.44	117.06	106.30
1	B	392	TRP	CD1-CG-CD2	13.30	116.94	106.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	397	TRP	CD1-CG-CD2	13.21	116.87	106.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	321	TYR	Sidechain
1	A	427	TYR	Sidechain
1	A	520	TYR	Sidechain
1	B	516	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4329	0	4332	108	0
1	B	4329	0	4333	115	0
2	A	1	0	0	0	0
3	A	20	0	10	3	0
4	A	84	0	0	4	0
4	B	114	0	0	10	0
All	All	8877	0	8675	223	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 13.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:70:ILE:HG21	1:A:166:LEU:HD21	1.34	1.09
1:B:40:GLN:HG3	1:B:156:TYR:HB2	1.47	0.96
1:B:63:ARG:HH22	1:B:115:ALA:HB2	1.35	0.91
1:A:441:MET:SD	1:A:501:PHE:HB3	2.15	0.86
1:B:391:LEU:HD23	4:B:573:HOH:O	1.77	0.85

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/561 (100%)	461 (82%)	73 (13%)	25 (4%)	4	7
1	B	559/561 (100%)	506 (90%)	39 (7%)	14 (2%)	9	21
All	All	1118/1122 (100%)	967 (86%)	112 (10%)	39 (4%)	6	12

5 of 39 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	446	ALA
1	A	470	ALA
1	A	486	ASN
1	A	487	GLN
1	A	531	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	462/462 (100%)	404 (87%)	58 (13%)	7	16
1	B	462/462 (100%)	410 (89%)	52 (11%)	9	19
All	All	924/924 (100%)	814 (88%)	110 (12%)	8	18

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	515	LEU
1	B	40	GLN
1	B	461	ASN
1	A	523	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	539	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	417	HIS
1	A	532	GLN
1	B	487	GLN
1	A	487	GLN
1	B	14	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	VG1	A	563	1,2	20,20,20	2.18	5 (25%)	24,31,31	1.41	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VG1	A	563	1,2	-	0/7/31/31	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	563	VG1	OV3-V	6.06	1.74	1.63
3	A	563	VG1	OV2-V	4.69	1.71	1.63
3	A	563	VG1	P-O1	-3.16	1.50	1.59
3	A	563	VG1	O5-C5	2.39	1.50	1.44
3	A	563	VG1	O6-C6	-2.20	1.39	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	563	VG1	O5-C5-C6	3.45	113.54	106.61
3	A	563	VG1	O5-C5-C4	-2.03	106.00	109.76
3	A	563	VG1	O2-C2-C1	-2.02	105.64	110.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.