



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 11:13 PM GMT

PDB ID : 1C4W
Title : 1.9 Å STRUCTURE OF A-THIOPHOSPHONATE MODIFIED CHEY D57C
Authors : Halkides, C.J.; McEvoy, M.M.; Matsumura, P.; Volz, K.; Dahlquist, F.W.
Deposited on : 1999-09-28
Resolution : 1.84 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2149 (1.86-1.82)
Ramachandran outliers	78287	2124 (1.86-1.82)
Sidechain outliers	78261	2125 (1.86-1.82)
RSRZ outliers	66119	1857 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	128	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1131 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHEMOTAXIS PROTEIN CHEY.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	128	991	625	163	193	2	8	0	2	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	57	CYQ	ASP	ENGINEERED	UNP P06143

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	140	Total	O	0	0
			140	140		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CHEMOTAXIS PROTEIN CHEY

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.62Å 47.74Å 53.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.84 23.21 – 1.84	Depositor EDS
% Data completeness (in resolution range)	88.3 (10.00-1.84) 96.7 (23.21-1.84)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.17 (at 1.84Å)	Xtriage
Refinement program	PROTIN, PROFFT	Depositor
R, R_{free}	0.208 , (Not available) 0.191 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.483	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 65.9	EDS
Estimated twinning fraction	0.028 for k,h,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 10492 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1131	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CYQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	1/990 (0.1%)	2.12	33/1329 (2.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	56	SER	CA-CB	6.15	1.62	1.52

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ARG	NE-CZ-NH1	23.24	131.92	120.30
1	A	73	ARG	NE-CZ-NH1	13.09	126.84	120.30
1	A	22	ARG	NE-CZ-NH1	12.48	126.54	120.30
1	A	12	ASP	CB-CG-OD1	11.96	129.07	118.30
1	A	64	ASP	CB-CG-OD1	9.59	126.93	118.30
1	A	22	ARG	NE-CZ-NH2	-9.26	115.67	120.30
1	A	18	ARG	NH1-CZ-NH2	-8.43	110.13	119.40
1	A	18	ARG	CD-NE-CZ	8.25	135.15	123.60
1	A	125	GLU	CA-CB-CG	8.11	131.24	113.40
1	A	19	ARG	NE-CZ-NH1	7.99	124.30	120.30
1	A	64	ASP	CB-CG-OD2	-7.92	111.17	118.30
1	A	22	ARG	CD-NE-CZ	7.83	134.57	123.60
1	A	66	LEU	CB-CG-CD2	-7.20	98.77	111.00
1	A	73	ARG	NE-CZ-NH2	-7.14	116.73	120.30
1	A	13	ASP	CB-CG-OD2	-6.56	112.39	118.30
1	A	78	MET	CB-CA-C	-6.51	97.37	110.40
1	A	20	ILE	CA-C-O	6.32	133.36	120.10
1	A	66	LEU	O-C-N	6.03	132.34	122.70
1	A	27	GLU	OE1-CD-OE2	5.99	130.49	123.30
1	A	78	MET	CG-SD-CE	-5.86	90.82	100.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	VAL	O-C-N	5.76	131.91	122.70
1	A	125	GLU	OE1-CD-OE2	5.75	130.21	123.30
1	A	126	LYS	CA-CB-CG	5.65	125.84	113.40
1	A	31	ASN	OD1-CG-ND2	5.58	134.74	121.90
1	A	12	ASP	OD1-CG-OD2	-5.50	112.84	123.30
1	A	35	GLU	CG-CD-OE1	5.36	129.01	118.30
1	A	63	MET	O-C-N	5.21	131.03	122.70
1	A	37	GLU	OE1-CD-OE2	5.18	129.52	123.30
1	A	74	ALA	N-CA-CB	-5.16	102.88	110.10
1	A	14	PHE	CB-CG-CD1	-5.15	117.19	120.80
1	A	27	GLU	O-C-N	-5.15	114.46	122.70
1	A	20	ILE	O-C-N	-5.14	114.48	122.70
1	A	46	LEU	O-C-N	5.12	130.90	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	991	0	1005	29	0
2	A	140	0	0	8	0
All	All	1131	0	1005	29	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 15.

All (29) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:12:ASP:OD2	1:A:57[B]:CYQ:SG	2.18	1.01
1:A:23[B]:ASN:ND2	2:A:236:HOH:O	2.08	0.86
1:A:87:THR:HG21	1:A:106:TYR:CZ	2.27	0.70
1:A:127:LEU:HB3	1:A:129:MET:HG3	1.79	0.65
1:A:23[B]:ASN:CG	2:A:236:HOH:O	2.39	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:46:LEU:HB3	1:A:78:MET:CE	2.33	0.58
1:A:87:THR:HG21	1:A:106:TYR:OH	2.04	0.58
1:A:87:THR:HG21	1:A:106:TYR:CE2	2.39	0.57
1:A:47:GLN:NE2	2:A:304:HOH:O	2.31	0.57
1:A:82:PRO:HG3	1:A:127:LEU:CD2	2.34	0.56
1:A:19:ARG:HD2	2:A:333:HOH:O	2.05	0.55
1:A:7:LYS:HE3	1:A:51:TYR:CZ	2.42	0.55
1:A:4:LYS:NZ	1:A:28:LEU:O	2.39	0.54
1:A:7:LYS:HE3	1:A:51:TYR:CE1	2.42	0.54
1:A:122:LYS:O	1:A:126:LYS:HB2	2.10	0.51
1:A:58:TRP:HE3	1:A:85:MET:HG2	1.77	0.49
1:A:2:ALA:HB2	1:A:124:PHE:CD2	2.48	0.49
1:A:75:ASP:O	1:A:79:SER:HB3	2.13	0.49
1:A:44:ASN:ND2	2:A:250:HOH:O	2.47	0.48
1:A:90:ALA:HB2	1:A:108:VAL:CG2	2.45	0.46
1:A:19:ARG:CD	2:A:333:HOH:O	2.64	0.46
1:A:45:LYS:NZ	2:A:269:HOH:O	2.51	0.44
1:A:53:PHE:CD2	1:A:82:PRO:HB2	2.53	0.43
1:A:19:ARG:HB2	2:A:315:HOH:O	2.18	0.43
1:A:46:LEU:HB3	1:A:78:MET:HE2	2.01	0.43
1:A:58:TRP:CE3	1:A:85:MET:HG2	2.54	0.43
1:A:127:LEU:O	1:A:128:GLY:C	2.57	0.43
1:A:90:ALA:HB2	1:A:108:VAL:HG21	2.01	0.42
1:A:43:LEU:O	1:A:47:GLN:HG3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/128 (98%)	125 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	102/101 (101%)	101 (99%)	1 (1%)	85 80

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CYQ	A	57[A]	-	4,5,11	3.53	3 (75%)	6,7,15	2.18	3 (50%)
1	CYQ	A	57[B]	-	4,5,11	3.46	3 (75%)	6,7,15	2.21	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CYQ	A	57[A]	-	-	0/0/3/11	0/0/0/0
1	CYQ	A	57[B]	-	-	0/0/3/11	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57[A]	CYQ	P-O2P	-4.66	1.45	1.54
1	A	57[A]	CYQ	P-O3P	-4.65	1.45	1.54
1	A	57[B]	CYQ	P-O2P	-4.64	1.45	1.54
1	A	57[B]	CYQ	P-O3P	-4.51	1.45	1.54
1	A	57[A]	CYQ	P-O1P	-2.27	1.45	1.50
1	A	57[B]	CYQ	P-O1P	-2.27	1.45	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	57[B]	CYQ	O3P-P-O2P	-3.38	97.65	108.35
1	A	57[A]	CYQ	O3P-P-O2P	-3.31	97.86	108.35
1	A	57[B]	CYQ	O2P-P-CD	2.68	113.17	106.61
1	A	57[A]	CYQ	O2P-P-CD	2.63	113.05	106.61
1	A	57[A]	CYQ	O3P-P-CD	2.62	113.03	106.61
1	A	57[B]	CYQ	O3P-P-CD	2.61	113.00	106.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	128/128 (100%)	0.14	4 (3%) 47 42	11, 20, 35, 46	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	128	GLY	4.1
1	A	125	GLU	2.7
1	A	129	MET	2.6
1	A	89	GLU	2.5

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CYQ	A	57[A]	6/12	0.19	2.78	17,19,20,20	6
1	CYQ	A	57[B]	6/12	0.19	2.78	17,19,20,20	6

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.