



Full wwPDB X-ray Structure Validation Report

Mar 1, 2014 – 03:01 AM GMT

PDB ID : 2C40
Title : CRYSTAL STRUCTURE OF INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FROM BACILLUS ANTHRACIS AT 2.2A RESOLUTION
Authors : Moroz, O.V.; Blagova, E.V.; Fogg, M.J.; Levnikov, V.M.; Brannigan, J.A.; Wilkinson, A.J.; Wilson, K.S.
Deposited on : 2005-10-13
Resolution : 2.20 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4867 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FAMILY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	3	0
			2380	1532	407	434	7			
1	B	300	Total	C	N	O	S	0	1	0
			2342	1507	396	432	7			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is SUGAR (RIBOSE) (three-letter code: RIB) (formula: C₅H₁₀O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			9	5	4		
3	B	1	Total	C	O	0	0
			9	5	4		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	77	Total	O	0	0
			77	77		
4	B	48	Total	O	0	0
			48	48		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FAMILY PROTEIN

Chain A: 



- Molecule 1: INOSINE-URIDINE PREFERRING NUCLEOSIDE HYDROLASE FAMILY PROTEIN

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	78.15Å 78.15Å 203.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	67.88 – 2.20 24.81 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (67.88-2.20) 99.7 (24.81-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.205 , 0.242 0.220 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.244	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.7	EDS
Estimated twinning fraction	0.045 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 37372 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4867	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, RIB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/2440 (0.0%)	0.89	7/3331 (0.2%)
1	B	0.82	2/2402 (0.1%)	0.87	5/3280 (0.2%)
All	All	0.82	3/4842 (0.1%)	0.88	12/6611 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	VAL	CB-CG1	-6.07	1.40	1.52
1	B	283	ASP	CG-OD1	5.64	1.38	1.25
1	B	283	ASP	CG-OD2	5.50	1.38	1.25

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	26	ASN	N-CA-CB	-7.10	97.83	110.60
1	A	286	ARG	NE-CZ-NH2	-7.07	116.76	120.30
1	A	276	GLN	CB-CA-C	-6.67	97.07	110.40
1	A	286	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	B	293	ASP	CB-CG-OD1	-6.04	112.87	118.30
1	B	54	ARG	NE-CZ-NH2	-5.98	117.31	120.30
1	A	147	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	A	15	LEU	CB-CG-CD1	5.42	120.21	111.00
1	A	45	MET	CG-SD-CE	-5.27	91.76	100.20
1	B	286	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	A	298	ARG	NE-CZ-NH1	5.08	122.84	120.30
1	B	13	ASP	CB-CG-OD2	-5.03	113.78	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2380	0	2312	53	0
1	B	2342	0	2264	37	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	9	0	7	0	0
3	B	9	0	6	0	0
4	A	77	0	0	5	0
4	B	48	0	0	4	0
All	All	4867	0	4589	89	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

All (89) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:113:LEU:HD23	1:B:120:THR:HG23	1.44	0.99
1:A:66:ASN:ND2	1:A:133:ARG:HH22	1.71	0.89
1:A:66:ASN:HD22	1:A:133:ARG:HH22	1.27	0.83
1:A:89:ILE:HD13	1:A:219:ILE:HD11	1.60	0.82
1:A:34:VAL:HB	1:A:45:MET:HE2	1.64	0.79
1:B:24:MET:CE	1:B:27:VAL:HG21	2.13	0.79
1:A:300:PHE:O	1:A:304:THR:HG23	1.83	0.78
1:B:113:LEU:HD23	1:B:120:THR:CG2	2.13	0.77
1:B:24:MET:HE2	1:B:27:VAL:HG21	1.65	0.77
1:B:66:ASN:ND2	1:B:133:ARG:HH22	1.83	0.77
1:B:300:PHE:O	1:B:304:THR:HG23	1.85	0.76
1:A:24:MET:HE3	1:A:27:VAL:HG21	1.68	0.76
1:A:45:MET:CE	1:A:64:ALA:HA	2.17	0.75
1:B:53:ASP:OD1	1:B:98:THR:HG23	1.88	0.73
1:A:181:VAL:HG12	1:A:269:VAL:HG21	1.72	0.72
1:A:24:MET:CE	1:A:27:VAL:HG21	2.20	0.72
1:A:53:ASP:OD1	1:A:98:THR:HG23	1.89	0.71
1:A:24:MET:HE3	1:A:27:VAL:CG2	2.20	0.71
1:A:185:TRP:O	1:A:286:ARG:HD2	1.92	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:113:LEU:CD2	1:B:120:THR:HG23	2.22	0.69
1:A:34:VAL:HB	1:A:45:MET:CE	2.23	0.67
1:A:45:MET:HE3	1:A:64:ALA:HA	1.77	0.66
1:B:185:TRP:O	1:B:286:ARG:HD2	1.96	0.65
1:A:262:VAL:HG11	1:A:289:HIS:HB3	1.76	0.65
1:A:53:ASP:OD1	1:A:98:THR:CG2	2.45	0.64
1:A:262:VAL:CG1	1:A:289:HIS:HB3	2.29	0.63
1:B:112:THR:O	1:B:116:THR:HG23	1.99	0.62
1:B:123:LEU:HG	1:B:254:VAL:HG21	1.82	0.61
1:A:45:MET:HE1	1:A:64:ALA:HA	1.82	0.61
1:A:300:PHE:O	1:A:304:THR:CG2	2.49	0.61
1:B:231:VAL:HG22	1:B:231:VAL:O	2.01	0.60
1:B:166:HIS:HE1	1:B:170:ALA:H	1.48	0.60
1:B:205:THR:HG22	1:B:208:ILE:HD12	1.83	0.60
1:A:113:LEU:HD22	1:A:120:THR:HG23	1.83	0.59
1:B:106:HIS:HD2	4:B:2023:HOH:O	1.86	0.59
1:A:115:GLN:CB	4:A:2029:HOH:O	2.50	0.59
1:A:45:MET:HE3	1:A:64:ALA:CB	2.34	0.58
1:A:231:VAL:O	1:A:231:VAL:HG13	2.04	0.56
1:A:298:ARG:HD2	4:A:2066:HOH:O	2.04	0.56
1:A:45:MET:HE3	1:A:64:ALA:CA	2.35	0.56
1:B:53:ASP:OD1	1:B:98:THR:CG2	2.53	0.55
1:B:83:TYR:HB3	1:B:230:ILE:HD13	1.88	0.55
1:A:120:THR:HG22	4:A:2030:HOH:O	2.06	0.53
1:B:66:ASN:HD21	1:B:133:ARG:HH22	1.54	0.53
1:A:166:HIS:HE1	1:A:170:ALA:H	1.57	0.52
1:B:66:ASN:C	1:B:66:ASN:HD22	2.13	0.52
1:B:66:ASN:ND2	1:B:133:ARG:NH2	2.56	0.51
1:A:112:THR:O	1:A:116:THR:HG23	2.10	0.51
1:A:120:THR:CG2	4:A:2030:HOH:O	2.58	0.51
1:B:185:TRP:HZ3	1:B:267:SER:HG	1.59	0.50
1:B:83:TYR:CB	1:B:230:ILE:HD13	2.42	0.49
1:A:123:LEU:HG	1:A:254:VAL:HG21	1.95	0.49
1:A:113:LEU:CD2	1:A:120:THR:HG23	2.43	0.49
1:B:6:PHE:CD2	1:B:123:LEU:HD13	2.48	0.48
1:B:13:ASP:HB3	1:B:247:ASP:HB2	1.96	0.48
1:B:199:THR:HG21	1:B:246:TRP:NE1	2.29	0.47
1:A:21:LEU:HD23	1:A:24:MET:CE	2.45	0.46
1:B:296:HIS:HD2	1:B:297:ASP:OD1	1.99	0.46
1:A:76:ASP:CB	4:A:2021:HOH:O	2.63	0.46
1:B:116:THR:HG21	4:B:2006:HOH:O	2.15	0.46
1:B:155:PHE:N	1:B:197:GLU:OE2	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:202:VAL:HG23	1:B:202:VAL:O	2.16	0.45
1:A:24:MET:HE3	1:A:27:VAL:CB	2.47	0.45
1:B:120:THR:HG21	4:B:2006:HOH:O	2.17	0.44
1:A:181:VAL:CG1	1:A:269:VAL:HG21	2.45	0.44
1:A:45:MET:HE3	1:A:64:ALA:HB2	1.99	0.44
1:A:45:MET:CE	1:A:64:ALA:CA	2.93	0.44
1:B:166:HIS:CE1	1:B:170:ALA:H	2.32	0.44
1:A:199:THR:HG21	1:A:244:TYR:HB3	1.99	0.44
1:B:166:HIS:CE1	1:B:170:ALA:HB3	2.53	0.43
1:B:24:MET:CE	1:B:27:VAL:CG2	2.91	0.43
1:B:209:ARG:NH1	4:B:2037:HOH:O	2.50	0.43
1:A:6:PHE:CD2	1:A:123:LEU:HD13	2.53	0.43
1:A:193:LEU:HD13	1:A:195:THR:HG22	2.01	0.43
1:A:54[A]:ARG:NH1	1:A:89:ILE:HD12	2.34	0.43
1:A:21:LEU:HA	1:A:24:MET:HE2	2.00	0.43
1:A:129:THR:HB	1:A:177:ASP:OD2	2.19	0.43
1:A:24:MET:HE1	1:A:27:VAL:HG21	1.97	0.43
1:A:83:TYR:HB2	1:A:230:ILE:HD13	2.00	0.42
1:A:199:THR:HG23	1:A:244:TYR:CD2	2.55	0.42
1:A:296:HIS:HD2	1:A:297:ASP:OD1	2.03	0.42
1:A:225:GLY:HA3	1:B:88:PRO:HG3	2.02	0.41
1:A:74:PRO:O	1:A:78:ARG:HG3	2.19	0.41
1:B:282:THR:HG22	1:B:284:ASP:H	1.85	0.41
1:A:54[A]:ARG:CZ	1:A:89:ILE:HD12	2.50	0.41
1:B:206:ILE:HD11	1:B:233:PRO:CB	2.52	0.40
1:A:194:ILE:N	1:A:194:ILE:HD12	2.36	0.40
1:A:231:VAL:CG1	1:A:231:VAL:O	2.68	0.40
1:A:21:LEU:HD23	1:A:24:MET:HE1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/312 (96%)	293 (97%)	8 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	297/312 (95%)	289 (97%)	8 (3%)	0	100	100
All	All	598/624 (96%)	582 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	247/269 (92%)	220 (89%)	27 (11%)	9	8
1	B	243/269 (90%)	219 (90%)	24 (10%)	11	10
All	All	490/538 (91%)	439 (90%)	51 (10%)	10	9

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	2	LYS
1	A	15	LEU
1	A	29	LEU
1	A	59	THR
1	A	66	ASN
1	A	89	ILE
1	A	98	THR
1	A	120	THR
1	A	123	LEU
1	A	129	THR
1	A	141	ILE
1	A	147	ARG
1	A	148	LEU
1	A	151	MET
1	A	183	ARG
1	A	185	TRP
1	A	204	LEU
1	A	205	THR
1	A	231	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	249	LEU
1	A	258	ASP
1	A	261	LYS
1	A	269	VAL
1	A	271	THR
1	A	283	ASP
1	A	304	THR
1	B	1	MET
1	B	13	ASP
1	B	15	LEU
1	B	29	LEU
1	B	59	THR
1	B	66	ASN
1	B	98	THR
1	B	120	THR
1	B	123	LEU
1	B	129	THR
1	B	133	ARG
1	B	147	ARG
1	B	148	LEU
1	B	151	MET
1	B	157	THR
1	B	186	GLU
1	B	193	LEU
1	B	197	GLU
1	B	204	LEU
1	B	205	THR
1	B	249	LEU
1	B	271	THR
1	B	298	ARG
1	B	304	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	66	ASN
1	A	106	HIS
1	A	166	HIS
1	A	296	HIS
1	B	7	ASN
1	B	66	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	106	HIS
1	B	166	HIS
1	B	296	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	RIB	A	1312	2	7,9,10	2.20	2 (28%)	8,12,14	2.42	3 (37%)
3	RIB	B	1312	2	7,9,10	2.02	2 (28%)	8,12,14	2.49	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RIB	A	1312	2	-	0/2/15/18	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	RIB	B	1312	2	-	0/2/15/18	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1312	RIB	C3'-C2'	4.35	1.59	1.53
3	B	1312	RIB	C3'-C2'	4.01	1.59	1.53
3	A	1312	RIB	O2'-C2'	3.29	1.50	1.43
3	B	1312	RIB	O2'-C2'	2.95	1.50	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1312	RIB	O2'-C2'-C3'	4.43	124.81	113.77
3	B	1312	RIB	C4'-C3'-C2'	4.21	106.87	101.77
3	A	1312	RIB	O2'-C2'-C3'	4.10	123.99	113.77
3	A	1312	RIB	C4'-C3'-C2'	3.60	106.13	101.77
3	A	1312	RIB	O4'-C4'-C3'	2.99	108.10	103.75
3	B	1312	RIB	O4'-C4'-C5'	-2.04	105.01	108.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/312 (96%)	0.31	16 (5%) 25 25	40, 43, 49, 54	0
1	B	300/312 (96%)	0.65	37 (12%) 5 4	39, 43, 49, 52	0
All	All	602/624 (96%)	0.48	53 (8%) 9 9	39, 43, 49, 54	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	272	TYR	4.7
1	B	285	GLY	4.2
1	B	283	ASP	4.0
1	B	124	PHE	3.6
1	B	128	LEU	3.6
1	B	203	PRO	3.6
1	B	282	THR	3.5
1	A	150	TRP	3.3
1	A	194	ILE	3.2
1	B	204	LEU	3.2
1	A	283	ASP	3.1
1	B	292	TYR	3.1
1	B	244	TYR	3.1
1	A	15	LEU	3.0
1	B	201	GLN	3.0
1	B	150	TRP	3.0
1	B	157	THR	3.0
1	A	193	LEU	3.0
1	A	217	LYS	3.0
1	B	202	VAL	2.9
1	A	18	LEU	2.8
1	B	233	PRO	2.8
1	B	116	THR	2.8
1	A	152	GLY	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	205	THR	2.8
1	B	114	LEU	2.7
1	B	123	LEU	2.7
1	A	16	VAL	2.7
1	B	232	PRO	2.6
1	B	136	TYR	2.6
1	B	1	MET	2.5
1	A	96	VAL	2.5
1	A	251	ALA	2.5
1	A	123	LEU	2.5
1	B	305	ARG	2.5
1	B	117	GLU	2.4
1	B	185	TRP	2.4
1	B	152	GLY	2.4
1	A	250	THR	2.3
1	B	58	ASN	2.3
1	B	127	PRO	2.3
1	B	215	GLU	2.2
1	A	124	PHE	2.2
1	A	174[A]	SER	2.2
1	B	125	THR	2.2
1	A	125	THR	2.1
1	B	281	GLU	2.1
1	B	126	GLY	2.1
1	B	59	THR	2.1
1	B	56	GLY	2.1
1	B	284	ASP	2.1
1	B	298	ARG	2.0
1	B	194	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	RIB	A	1312	9/10	0.27	1.91	42,48,51,52	0
3	RIB	B	1312	9/10	0.20	0.64	38,47,50,51	0
2	CA	B	1311	1/1	0.06	-2.61	33,33,33,33	0
2	CA	A	1311	1/1	0.05	-5.03	31,31,31,31	0

6.5 Other polymers

There are no such residues in this entry.