



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 04:26 PM GMT

PDB ID : 3C4C
Title : B-Raf Kinase in Complex with PLX4720
Authors : Zhang, K.Y.J.; Wang, W.
Deposited on : 2008-01-29
Resolution : 2.57 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2358 (2.60-2.56)
Ramachandran outliers	78287	2316 (2.60-2.56)
Sidechain outliers	78261	2316 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	280	
1	B	280	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4319 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B-Raf proto-oncogene serine/threonine-proteinkinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			2058	1301	366	378	13			
1	B	257	Total	C	N	O	S	0	0	0
			2054	1299	365	377	13			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	543	ALA	ILE	ENGINEERED	UNP P15056
A	544	SER	ILE	ENGINEERED	UNP P15056
A	551	LYS	ILE	ENGINEERED	UNP P15056
A	562	ARG	GLN	ENGINEERED	UNP P15056
A	588	ASN	LEU	ENGINEERED	UNP P15056
A	630	SER	LYS	ENGINEERED	UNP P15056
A	667	GLU	PHE	ENGINEERED	UNP P15056
A	673	SER	TYR	ENGINEERED	UNP P15056
A	688	ARG	ALA	ENGINEERED	UNP P15056
A	706	SER	LEU	ENGINEERED	UNP P15056
A	709	ARG	GLN	ENGINEERED	UNP P15056
A	713	GLU	SER	ENGINEERED	UNP P15056
A	716	GLU	LEU	ENGINEERED	UNP P15056
A	720	GLU	SER	ENGINEERED	UNP P15056
A	722	SER	-	EXPRESSION TAG	UNP P15056
A	723	GLY	-	EXPRESSION TAG	UNP P15056
B	543	ALA	ILE	ENGINEERED	UNP P15056
B	544	SER	ILE	ENGINEERED	UNP P15056
B	551	LYS	ILE	ENGINEERED	UNP P15056
B	562	ARG	GLN	ENGINEERED	UNP P15056
B	588	ASN	LEU	ENGINEERED	UNP P15056
B	630	SER	LYS	ENGINEERED	UNP P15056
B	667	GLU	PHE	ENGINEERED	UNP P15056
B	673	SER	TYR	ENGINEERED	UNP P15056
B	688	ARG	ALA	ENGINEERED	UNP P15056

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	706	SER	LEU	ENGINEERED	UNP P15056
B	709	ARG	GLN	ENGINEERED	UNP P15056
B	713	GLU	SER	ENGINEERED	UNP P15056
B	716	GLU	LEU	ENGINEERED	UNP P15056
B	720	GLU	SER	ENGINEERED	UNP P15056
B	722	SER	-	EXPRESSION TAG	UNP P15056
B	723	GLY	-	EXPRESSION TAG	UNP P15056

- Molecule 2 is N-{3-[(5-CHLORO-1H-PYRROLO[2,3-B]PYRIDIN-3-YL)CARBONYL]-2,4-DIFLUOROPHENYL}PROPANE-1-SULFONAMIDE (three-letter code: 324) (formula: C₁₇H₁₄ClF₂N₃O₃S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	S	0	0
			27	17	1	2	3	3	1		
2	B	1	Total	C	Cl	F	N	O	S	0	0
			27	17	1	2	3	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	87	Total	O	0	0
			87	87		
3	B	66	Total	O	0	0
			66	66		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: B-Raf proto-oncogene serine/threonine-proteinkinase

Chain A:



- Molecule 1: B-Raf proto-oncogene serine/threonine-proteinkinase

Chain B:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.57Å 105.50Å 110.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.57	Depositor
% Data completeness (in resolution range)	99.7 (50.00-2.57)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	REFMAC 5.1.25	Depositor
R, R_{free}	0.259 , 0.303	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4319	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 324

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2100	0.97	10/2830 (0.4%)
1	B	0.72	0/2096	0.93	7/2825 (0.2%)
All	All	0.73	0/4196	0.95	17/5655 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	3
All	All	0	4

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	587	ASP	CB-CG-OD2	7.66	125.20	118.30
1	B	704	ARG	NE-CZ-NH1	7.34	123.97	120.30
1	A	449	ASP	CB-CG-OD2	7.12	124.71	118.30
1	A	576	ASP	CB-CG-OD2	7.08	124.67	118.30
1	A	479	ASP	CB-CG-OD2	6.73	124.35	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	460	GLY	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	520	SER	Peptide
1	B	545	GLU	Peptide
1	B	615	GLY	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2058	0	2061	72	0
1	B	2054	0	2058	64	0
2	A	27	0	14	2	0
2	B	27	0	14	8	0
3	A	87	0	0	5	0
3	B	66	0	0	4	0
All	All	4319	0	4147	134	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:719:ARG:HA	1:A:720:GLU:O	1.62	1.00
1:B:619:TRP:O	3:B:100:HOH:O	1.91	0.88
1:B:688:ARG:HD3	1:B:717:LEU:CD2	2.07	0.83
1:B:688:ARG:HD3	1:B:717:LEU:HD21	1.65	0.78
1:A:490:PRO:O	3:A:781:HOH:O	2.03	0.77

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/280 (91%)	216 (85%)	26 (10%)	12 (5%)	4	4
1	B	253/280 (90%)	213 (84%)	28 (11%)	12 (5%)	4	4
All	All	507/560 (90%)	429 (85%)	54 (11%)	24 (5%)	4	4

5 of 24 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	627	MET
1	A	628	GLN
1	A	629	ASP
1	A	631	ASN
1	A	720	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/246 (92%)	186 (82%)	40 (18%)	3	4
1	B	226/246 (92%)	168 (74%)	58 (26%)	1	1
All	All	452/492 (92%)	354 (78%)	98 (22%)	1	2

5 of 98 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	458	THR
1	B	494	GLN
1	B	684	ASN
1	B	459	VAL
1	B	471	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	510	HIS
1	A	512	ASN
1	A	628	GLN
1	A	631	ASN
1	B	539	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	324	A	2	-	29,29,29	3.19	8 (27%)	43,43,43	2.98	19 (44%)
2	324	B	3	-	29,29,29	3.32	10 (34%)	43,43,43	2.58	18 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	324	A	2	-	-	0/17/17/17	0/1/3/3
2	324	B	3	-	-	0/17/17/17	0/1/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	324	C17-C18	12.59	1.49	1.39
2	A	2	324	C17-C18	11.00	1.47	1.39
2	A	2	324	C9-C2	8.59	1.49	1.42
2	B	3	324	C13-C14	6.42	1.48	1.38
2	B	3	324	C9-C2	5.70	1.47	1.42

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	324	C9-C2-C3	-8.65	101.43	107.91
2	A	2	324	O23-S22-O24	-7.96	104.68	119.30
2	B	3	324	C9-C8-N7	6.17	110.38	108.15
2	A	2	324	F20-C18-C17	5.77	126.01	118.57
2	B	3	324	C14-C13-C18	5.65	120.45	115.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.