



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 04:02 PM GMT

PDB ID : 2C5R
Title : THE STRUCTURE OF PHAGE PHI29 REPLICATION ORGANIZER PROTEIN P16.7 IN COMPLEX WITH DOUBLE STRANDED DNA
Authors : Albert, A.; Jimenez, M.; Munoz-Espin, D.; Asensio, J.L.; Hermoso, J.A.; Salas, M.; Meijer, W.J.J.
Deposited on : 2005-10-31
Resolution : 2.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1053 (2.90-2.90)
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	67	
1	B	67	
1	C	67	
1	D	67	
1	E	67	
1	F	67	
2	Y	8	
3	Z	8	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3535 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EARLY PROTEIN P16.7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			
1	B	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			
1	C	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			
1	D	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			
1	E	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			
1	F	64	Total	C	N	O	S	0	0	1
			526	329	91	105	1			

- Molecule 2 is a DNA chain called 5'-D(*TP*CP*CP*AP*CP*CP*GP*GP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	Y	8	Total	C	N	O	P	0	0	0
			161	76	29	48	8			

- Molecule 3 is a DNA chain called 5'-D(*CP*CP*GP*GP*TP*GP*GP*AP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Z	8	Total	C	N	O	P	0	0	0
			167	78	33	48	8			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total	O	0	0
			12	12		
4	B	11	Total	O	0	0
			11	11		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	3	Total 3	O 3	0	0
4	D	6	Total 6	O 6	0	0
4	E	9	Total 9	O 9	0	0
4	F	7	Total 7	O 7	0	0
4	Z	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EARLY PROTEIN P16.7




• Molecule 1: EARLY PROTEIN P16.7




• Molecule 1: EARLY PROTEIN P16.7




• Molecule 1: EARLY PROTEIN P16.7



• Molecule 1: EARLY PROTEIN P16.7



• Molecule 1: EARLY PROTEIN P16.7



- Molecule 2: 5'-D(*TP*CP*CP*AP*CP*CP*GP*GP)-3'

Chain Y: 



- Molecule 3: 5'-D(*CP*CP*GP*GP*TP*GP*GP*AP)-3'

Chain Z: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.50Å 72.11Å 127.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.90 45.66 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.8 (50.00-2.90) 95.8 (45.66-2.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.251 , 0.292 0.285 , 0.319	Depositor DCC
R_{free} test set	956 reflections (7.17%)	DCC
Wilson B-factor (Å ²)	92.6	Xtriage
Anisotropy	0.286	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 23.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 13346 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3535	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.10	0/531	1.30	3/718 (0.4%)
1	B	1.28	2/532 (0.4%)	1.24	4/720 (0.6%)
1	C	1.08	0/532	1.17	5/720 (0.7%)
1	D	1.19	1/532 (0.2%)	1.17	4/720 (0.6%)
1	E	1.14	0/532	1.11	1/720 (0.1%)
1	F	1.17	1/532 (0.2%)	1.26	6/720 (0.8%)
2	Y	1.61	8/179 (4.5%)	1.23	0/273
3	Z	1.56	3/187 (1.6%)	1.20	0/287
All	All	1.21	15/3557 (0.4%)	1.21	23/4878 (0.5%)

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	71	GLU	CG-CD	8.39	1.64	1.51
1	D	68	SER	CB-OG	8.16	1.52	1.42
3	Z	2	DC	C2-N3	7.05	1.41	1.35
1	B	68	SER	CB-OG	6.84	1.51	1.42
2	Y	10	DC	N3-C4	-6.61	1.29	1.33

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	LEU	CA-C-O	13.29	148.01	120.10
1	A	88	ASP	CB-CG-OD2	9.16	126.54	118.30
1	F	112	ARG	NE-CZ-NH1	8.11	124.35	120.30
1	B	112	ARG	NE-CZ-NH1	7.13	123.86	120.30
1	B	67	LEU	CA-CB-CG	6.94	131.26	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	526	0	522	24	0
1	B	526	0	522	20	0
1	C	526	0	522	28	0
1	D	526	0	522	19	0
1	E	526	0	522	22	0
1	F	526	0	522	19	0
2	Y	161	0	90	1	0
3	Z	167	0	90	8	0
4	A	12	0	0	2	0
4	B	11	0	0	2	0
4	C	3	0	0	3	0
4	D	6	0	0	0	0
4	E	9	0	0	3	0
4	F	7	0	0	2	0
4	Z	3	0	0	1	0
All	All	3535	0	3312	115	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 17.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:127:SER:O	1:A:129:LYS:N	1.64	1.29
4:B:2008:HOH:O	1:C:103:GLU:HG2	1.55	1.05
1:A:102:GLN:NE2	1:A:102:GLN:HA	1.64	1.04
1:A:128:LEU:C	1:A:129:LYS:N	2.15	1.00
1:A:102:GLN:CA	1:A:102:GLN:HE21	1.74	1.00

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	61/67 (91%)	54 (88%)	6 (10%)	1 (2%)	14	47
1	B	62/67 (92%)	53 (86%)	7 (11%)	2 (3%)	6	25
1	C	62/67 (92%)	53 (86%)	7 (11%)	2 (3%)	6	25
1	D	62/67 (92%)	53 (86%)	7 (11%)	2 (3%)	6	25
1	E	62/67 (92%)	53 (86%)	7 (11%)	2 (3%)	6	25
1	F	62/67 (92%)	55 (89%)	5 (8%)	2 (3%)	6	25
All	All	371/402 (92%)	321 (86%)	39 (10%)	11 (3%)	7	27

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	128	LEU
1	E	128	LEU
1	C	97	ARG
1	D	127	SER
1	D	128	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	60/64 (94%)	50 (83%)	10 (17%)	3	9
1	B	60/64 (94%)	50 (83%)	10 (17%)	3	9
1	C	60/64 (94%)	49 (82%)	11 (18%)	2	7
1	D	60/64 (94%)	48 (80%)	12 (20%)	2	6
1	E	60/64 (94%)	49 (82%)	11 (18%)	2	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	60/64 (94%)	52 (87%)	8 (13%)	6	16
All	All	360/384 (94%)	298 (83%)	62 (17%)	3	8

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	125	ARG
1	D	96	GLN
1	F	97	ARG
1	D	68	SER
1	D	102	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	106	ASN
1	D	106	ASN
1	F	82	ASN
1	C	102	GLN
1	E	111	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	64/67 (95%)	1.43	18 (28%) 1 1	46, 51, 57, 60	0
1	B	64/67 (95%)	1.48	16 (25%) 1 2	46, 51, 56, 58	0
1	C	64/67 (95%)	1.08	13 (20%) 1 2	47, 51, 55, 58	0
1	D	64/67 (95%)	1.42	17 (26%) 1 1	46, 51, 55, 55	0
1	E	64/67 (95%)	1.40	13 (20%) 1 2	47, 51, 57, 58	0
1	F	64/67 (95%)	1.21	12 (18%) 2 2	46, 51, 54, 55	0
2	Y	8/8 (100%)	3.13	5 (62%) 0 0	52, 52, 52, 53	0
3	Z	8/8 (100%)	5.78	8 (100%) 0 0	51, 52, 52, 53	0
All	All	400/418 (95%)	1.46	102 (25%) 1 2	46, 51, 55, 60	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	129	LYS	12.1
3	Z	1	DC	8.9
1	D	124	TYR	7.9
3	Z	3	DG	7.6
2	Y	16	DG	7.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.