



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 09:51 PM GMT

PDB ID : 3C5X
Title : Crystal structure of the precursor membrane protein- envelope protein heterodimer from the dengue 2 virus at low pH
Authors : Li, L.
Deposited on : 2008-02-01
Resolution : 2.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	3751 (2.20-2.20)
Ramachandran outliers	78287	3681 (2.20-2.20)
Sidechain outliers	78261	3682 (2.20-2.20)
RSRZ outliers	66119	2939 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	402	
2	C	130	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
6	BMA	C	1400	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3860 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope protein E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	390	Total	C	N	O	S	0	0	0
			3047	1926	525	569	27			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP O09234
A	-6	GLU	-	EXPRESSION TAG	UNP O09234
A	-5	ASN	-	EXPRESSION TAG	UNP O09234
A	-4	LEU	-	EXPRESSION TAG	UNP O09234
A	-3	TYR	-	EXPRESSION TAG	UNP O09234
A	-2	PHE	-	EXPRESSION TAG	UNP O09234
A	-1	GLN	-	EXPRESSION TAG	UNP O09234
A	0	GLY	-	EXPRESSION TAG	UNP O09234

- Molecule 2 is a protein called prM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	81	Total	C	N	O	S	0	0	0
			640	396	106	128	10			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	87	SER	ARG	ENGINEERED	UNP O09234
C	88	THR	ARG	ENGINEERED	UNP O09234
C	91	SER	ARG	ENGINEERED	UNP O09234

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	EXPRESSION TAG	UNP O09234
A	-6	GLU	-	EXPRESSION TAG	UNP O09234
A	-5	ASN	-	EXPRESSION TAG	UNP O09234
A	-4	LEU	-	EXPRESSION TAG	UNP O09234
A	-3	TYR	-	EXPRESSION TAG	UNP O09234
A	-2	PHE	-	EXPRESSION TAG	UNP O09234
A	-1	GLN	-	EXPRESSION TAG	UNP O09234
A	0	GLY	-	EXPRESSION TAG	UNP O09234


- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	2	Total	C	N	O	0	0
			28	16	2	10		

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	87	SER	ARG	ENGINEERED	UNP O09234
C	88	THR	ARG	ENGINEERED	UNP O09234
C	91	SER	ARG	ENGINEERED	UNP O09234

- Molecule 5 is ALPHA-D-ALTROPYRANOSE (three-letter code: SHD) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is SUGAR (BETA-D-MANNOSE) (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			11	6	5		
6	C	1	Total	C	O	0	0
			11	6	5		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	61	Total 61	O 61	0	0
7	C	23	Total 23	O 23	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Envelope protein E

Chain A: 



• Molecule 2: prM

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	71.21Å 108.59Å 108.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 43.21 – 2.21	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-2.20) 98.1 (43.21-2.21)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.74 (at 2.20Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.253 , 0.275 0.257 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	45.1	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 49.7	EDS
Estimated twinning fraction	0.016 for -h,l,k	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 42456 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3860	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, SHD, NAG, NDG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	1/3108 (0.0%)	0.72	7/4189 (0.2%)
2	C	0.53	1/652 (0.2%)	0.67	0/883
All	All	0.38	2/3760 (0.1%)	0.71	7/5072 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
4	C	1	0
All	All	1	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	49	ILE	C-N	-8.66	1.14	1.34
1	A	77	GLN	C-N	5.20	1.42	1.33

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	GLU	N-CA-C	13.72	148.06	111.00
1	A	196	MET	N-CA-CB	-10.48	91.74	110.60
1	A	195	GLU	CB-CA-C	-7.94	94.51	110.40
1	A	384	PRO	N-CA-C	5.92	127.50	112.10
1	A	196	MET	N-CA-C	-5.47	96.22	111.00

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	C	1396	NAG	C1

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	193	PHE	Peptide
1	A	194	ASN	Peptide
1	A	362	ASP	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3047	0	3047	130	0
2	C	640	0	609	20	0
3	A	28	0	25	5	0
4	C	28	0	24	12	0
5	C	11	0	7	2	0
6	C	22	0	20	0	0
7	A	61	0	0	6	0
7	C	23	0	0	0	0
All	All	3860	0	3732	157	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 21.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:195:GLU:O	1:A:195:GLU:CG	1.87	1.20
2:C:31:VAL:HG21	4:C:1397:NAG:C8	1.72	1.19
2:C:31:VAL:CG2	4:C:1397:NAG:H81	1.74	1.18
1:A:195:GLU:O	1:A:195:GLU:HG2	1.57	0.98
1:A:195:GLU:O	1:A:195:GLU:HG3	1.62	0.97

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	382/402 (95%)	365 (96%)	14 (4%)	3 (1%)	27	24
2	C	79/130 (61%)	74 (94%)	5 (6%)	0	100	100
All	All	461/532 (87%)	439 (95%)	19 (4%)	3 (1%)	30	28

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	195	GLU
1	A	187	PRO
1	A	364	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	339/348 (97%)	324 (96%)	15 (4%)	39	45
2	C	75/118 (64%)	70 (93%)	5 (7%)	23	24
All	All	414/466 (89%)	394 (95%)	20 (5%)	35	41

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	207	LEU
1	A	305	LYS
2	C	23	LEU
1	A	201	MET
1	A	202	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	242	ASN
1	A	256	GLN
1	A	316	GLN
1	A	233	GLN
1	A	276	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1398	3	12,14,15	0.61	0	15,19,21	0.84	0
3	NDG	A	1399	3	12,14,15	0.41	0	15,19,21	0.51	0
4	NAG	C	1396	2,4	12,14,15	0.55	0	15,19,21	1.57	4 (26%)
4	NAG	C	1397	5,4	12,14,15	0.51	0	15,19,21	1.64	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1398	3	-	0/6/23/26	0/1/1/1
3	NDG	A	1399	3	-	1/6/23/26	0/1/1/1
4	NAG	C	1396	2,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	C	1397	5,4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	1397	NAG	C3-C4-C5	-4.26	102.59	110.20
4	C	1396	NAG	O5-C5-C4	-2.58	107.38	110.65
4	C	1396	NAG	O5-C5-C6	2.58	109.69	106.98
4	C	1397	NAG	C6-C5-C4	2.45	118.91	113.00
4	C	1396	NAG	C4-C3-C2	-2.37	105.51	111.32

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	C	1396	NAG	C1

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1399	NDG	O7-C7-N2-C2

There are no ring outliers.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SHD	C	1398	4,6	10,11,12	1.54	3 (30%)	11,15,17	6.44	4 (36%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	BMA	C	1399	5	10,11,12	0.44	0	11,15,17	1.28	2 (18%)
6	BMA	C	1400	5	10,11,12	0.60	0	11,15,17	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SHD	C	1398	4,6	-	0/2/19/22	0/1/1/1
6	BMA	C	1399	5	-	0/2/19/22	0/1/1/1
6	BMA	C	1400	5	1/1/4/5	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1398	SHD	C3-C2	-2.85	1.46	1.52
5	C	1398	SHD	O2-C3	-2.61	1.36	1.43
5	C	1398	SHD	C4-C3	2.40	1.58	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1398	SHD	O1-C2-C3	-17.70	71.99	110.18
5	C	1398	SHD	O3-C4-C3	10.09	132.98	110.35
5	C	1398	SHD	O2-C3-C4	-5.47	98.08	110.35
6	C	1399	BMA	C6-C5-C4	-2.95	105.87	113.00
5	C	1398	SHD	C3-C4-C5	2.17	114.07	110.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	C	1400	BMA	C1

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	390/402 (97%)	0.49	23 (5%) 22 21	30, 52, 87, 113	0
2	C	81/130 (62%)	0.34	0 100 100	22, 41, 73, 97	0
All	All	471/532 (88%)	0.47	23 (4%) 26 28	22, 50, 85, 113	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	HIS	4.5
1	A	191	LEU	4.1
1	A	194	ASN	4.0
1	A	193	PHE	3.8
1	A	303	THR	3.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAG	C	1397	14/15	0.40	4.82	104,119,121,124	0
4	NAG	C	1396	14/15	0.18	1.10	64,69,76,93	0
3	NAG	A	1398	14/15	0.20	0.62	79,84,91,108	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NDG	A	1399	14/15	0.35	-	118,132,133,133	0

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	BMA	C	1400	11/12	0.52	8.82	119,120,121,122	0
6	BMA	C	1399	11/12	0.46	-	138,141,142,142	0
5	SHD	C	1398	11/12	0.37	-	124,127,130,135	0

6.5 Other polymers

There are no such residues in this entry.