



# wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 10:12 PM GMT

PDB ID : 2C6F  
Title : STRUCTURE OF HUMAN SOMATIC ANGIOTENSIN-I CONVERTING  
ENZYME N DOMAIN  
Authors : Corradi, H.R.; Schwager, S.L.U.; Nichinda, A.; Sturrock, E.D.; Acharya, K.R.  
Deposited on : 2005-11-09  
Resolution : 3.01 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1732 (3.04-3.00)
Ramachandran outliers	78287	1669 (3.04-3.00)
Sidechain outliers	78261	1672 (3.04-3.00)
RSRZ outliers	66119	1333 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	612	<div><div></div><div></div><div></div><div></div></div>
1	B	612	<div><div></div><div></div><div></div><div></div></div>

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NAG	A	695	-	X
3	NAG	A	697	-	X
3	NAG	B	692	-	X
3	NAG	B	695	-	X
6	ACT	A	800	-	X
6	ACT	A	801	-	X
6	ACT	B	801	-	X
7	GOL	A	2434	-	X
7	GOL	B	2433	X	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 9970 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ANGIOTENSIN-CONVERTINGENZYME, SOMATIC ISOFORM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	612	Total	C	N	O	S	6	0	0
			4884	3145	840	880	19			
1	B	611	Total	C	N	O	S	14	0	0
			4852	3124	830	879	19			

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Zn	0	0
			1	1		
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is ACETATE ION (three-letter code: ACT) (formula:  $\text{C}_2\text{H}_3\text{O}_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $\text{C}_3\text{H}_8\text{O}_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	11	Total	O	0	0
			11	11		
8	B	13	Total	O	0	0
			13	13		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ANGIOTENSIN-CONVERTINGENZYME, SOMATIC ISOFORM

Chain A: 



#### • Molecule 1: ANGIOTENSIN-CONVERTINGENZYME, SOMATIC ISOFORM

Chain B: 



V351	T352	K353	D354	Q355	L356	E362	H365	I366	Q367	Y368	Y369	Y372	K373	D374	Y377	R381	G382	A383	F387	H388	E389	F490	H491	V492	P493	M494	V495	S504	L507	Q508	F509	Q510	V414	D417	Y418	E419	S420	L426	K427	M428	A429	I433	F435	L436	P437	F438	G439	Y440
L441	V442	D443	Q444	W445	R453	P456	Y459	N460	W464	P475	P476	V477	T478	R479	N480	E481	T482	A486	K489	F490	H491	V492	P493	M494	V495	S504	L507	Q508	F509	Q510	D529	I530	Y531	R532	A536	K539	L540	R541	K542	V543	L544	S548	P551	W552	Q553			
E554	K557	D558	M559	V560	G561	L562	D563	A564	D566	A567	Q568	P569	L570	L571	K572	Y573	F574	Q575	W580	Q584	N585	Q586	E590	G593	N594	P595	E596	Y597	Q598	N606	Y607	P608	D612															



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.12Å 211.32Å 171.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.91 – 3.01 47.91 – 3.01	Depositor EDS
% Data completeness (in resolution range)	96.7 (47.91-3.01) 96.8 (47.91-3.01)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 3.01Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.224 , 0.273 0.222 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	45.6	Xtriage
Anisotropy	0.335	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 15.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 35643 reflections	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	9970	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.90% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, ACT, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.51	1/5042 (0.0%)	0.74	3/6885 (0.0%)
1	B	0.49	0/5010	0.86	6/6845 (0.1%)
All	All	0.50	1/10052 (0.0%)	0.80	9/13730 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	3	2
All	All	3	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	117	ASN	CB-CG	-5.47	1.38	1.51

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	PRO	CA-N-CD	-22.64	79.80	111.50
1	B	608	PRO	CA-N-CD	-18.89	85.05	111.50
1	B	607	TYR	CB-CG-CD1	-9.99	115.01	121.00
1	B	607	TYR	CB-CG-CD2	7.49	125.49	121.00
1	B	409	GLY	N-CA-C	5.92	127.91	113.10

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	79	ILE	CA
1	B	410	LEU	CA
1	B	530	ILE	CB

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	607	TYR	Peptide
1	B	275	PRO	Peptide
1	B	409	GLY	Peptide

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4884	0	4582	252	0
1	B	4852	0	4514	243	0
2	A	56	0	50	7	0
2	B	56	0	50	5	0
3	A	42	0	36	4	0
3	B	28	0	23	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	1	0
6	A	8	0	6	0	0
6	B	4	0	3	0	0
7	A	6	0	6	1	0
7	B	6	0	6	1	0
8	A	11	0	0	0	0
8	B	13	0	0	0	0
All	All	9970	0	9276	499	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 26.

The worst 5 of 499 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:3:PRO:HD3	1:B:6:GLN:NE2	1.36	1.40
1:B:282:THR:HG21	1:B:410:LEU:CB	1.60	1.31
1:A:279:VAL:HG21	1:A:410:LEU:CD1	1.60	1.29
1:B:282:THR:CG2	1:B:410:LEU:CB	2.18	1.20
1:A:279:VAL:CG1	1:A:410:LEU:HD13	1.73	1.17

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	610/612 (100%)	537 (88%)	60 (10%)	13 (2%)	11	45
1	B	609/612 (100%)	526 (86%)	73 (12%)	10 (2%)	14	54
All	All	1219/1224 (100%)	1063 (87%)	133 (11%)	23 (2%)	12	49

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	608	PRO
1	B	3	PRO
1	B	608	PRO
1	A	7	PRO
1	A	250	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	496/526 (94%)	448 (90%)	48 (10%)	12	41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	489/526 (93%)	437 (89%)	52 (11%)	10	35
All	All	985/1052 (94%)	885 (90%)	100 (10%)	11	37

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	582	GLN
1	B	129	LEU
1	B	553	GLN
1	A	587	GLN
1	B	66	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	6	GLN
1	B	18	GLN
1	B	575	GLN
1	A	545	GLN
1	A	575	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	691	1,2	12,14,15	0.54	0	15,19,21	0.67	0
2	NAG	A	693	1,2	12,14,15	0.54	0	15,19,21	0.66	0
2	NAG	A	694	2	12,14,15	0.42	0	15,19,21	0.65	0
2	NAG	A	696	2	12,14,15	0.41	0	15,19,21	0.65	0
2	NAG	B	691	1,2	12,14,15	0.53	0	15,19,21	0.67	0
2	NAG	B	693	1,2	12,14,15	2.53	1 (8%)	15,19,21	1.42	2 (13%)
2	NAG	B	694	2	12,14,15	2.52	1 (8%)	15,19,21	1.42	2 (13%)
2	NAG	B	696	2	12,14,15	0.42	0	15,19,21	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	691	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	693	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	694	2	-	0/6/23/26	0/1/1/1
2	NAG	A	696	2	-	0/6/23/26	0/1/1/1
2	NAG	B	691	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	693	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	694	2	-	0/6/23/26	0/1/1/1
2	NAG	B	696	2	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	693	NAG	C8-C7	-8.65	1.32	1.50
2	B	694	NAG	C8-C7	-8.60	1.32	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	693	NAG	C3-C2-N2	-4.44	105.00	111.76
2	B	694	NAG	C3-C2-N2	-4.43	105.01	111.76
2	B	694	NAG	O5-C5-C6	2.38	109.48	106.98
2	B	693	NAG	O5-C5-C6	2.35	109.44	106.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	GOL	A	2434	-	5,5,5	3.22	3 (60%)	5,5,5	1.73	1 (20%)
3	NAG	A	692	1	12,14,15	4.20	3 (25%)	15,19,21	1.81	4 (26%)
3	NAG	A	695	1	12,14,15	0.55	0	15,19,21	0.66	0
3	NAG	A	697	1	12,14,15	0.54	0	15,19,21	0.66	0
6	ACT	A	800	-	1,3,3	3.00	1 (100%)	0,3,3	0.00	-
6	ACT	A	801	-	1,3,3	3.47	1 (100%)	0,3,3	0.00	-
7	GOL	B	2433	-	5,5,5	3.25	3 (60%)	5,5,5	1.89	3 (60%)
3	NAG	B	692	1	12,14,15	4.06	2 (16%)	15,19,21	1.80	4 (26%)
3	NAG	B	695	1	12,14,15	0.54	0	15,19,21	0.66	0
6	ACT	B	801	-	1,3,3	3.01	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	2434	-	-	0/4/4/4	0/0/0/0
3	NAG	A	692	1	-	0/6/23/26	0/1/1/1
3	NAG	A	695	1	-	0/6/23/26	0/1/1/1
3	NAG	A	697	1	-	0/6/23/26	0/1/1/1
6	ACT	A	800	-	-	0/0/0/0	0/0/0/0
6	ACT	A	801	-	-	0/0/0/0	0/0/0/0
7	GOL	B	2433	-	-	0/4/4/4	0/0/0/0
3	NAG	B	692	1	-	0/6/23/26	0/1/1/1
3	NAG	B	695	1	-	0/6/23/26	0/1/1/1
6	ACT	B	801	-	-	0/0/0/0	0/0/0/0

The worst 5 of 14 bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	692	NAG	C8-C7	-13.38	1.22	1.50
3	B	692	NAG	C8-C7	-13.34	1.22	1.50
7	B	2433	GOL	C3-C2	-6.22	1.26	1.52
7	A	2434	GOL	C3-C2	-6.16	1.26	1.52
3	A	692	NAG	O7-C7	4.25	1.32	1.23

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	692	NAG	C3-C2-N2	-4.46	104.97	111.76
3	B	692	NAG	C3-C2-N2	-4.42	105.03	111.76
3	A	692	NAG	C8-C7-N2	3.51	122.98	116.11
3	B	692	NAG	C8-C7-N2	3.50	122.96	116.11
3	A	692	NAG	O7-C7-N2	-2.59	116.49	121.90

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	612/612 (100%)	-0.31	0 100 100	12, 36, 58, 83	3 (0%)
1	B	611/612 (99%)	-0.24	0 100 100	12, 38, 68, 84	6 (0%)
All	All	1223/1224 (99%)	-0.28	0 93 100	12, 37, 64, 84	9 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	694	14/15	0.33	17.37	79,80,81,81	0
2	NAG	B	691	14/15	0.32	12.97	55,57,62,69	0
2	NAG	A	691	14/15	0.33	6.32	56,60,68,72	0
2	NAG	A	694	14/15	0.28	1.55	69,72,73,75	0
2	NAG	B	693	14/15	0.23	0.55	61,65,67,68	0
2	NAG	A	693	14/15	0.23	-0.11	56,58,60,60	0
2	NAG	B	696	14/15	0.39	-	86,90,91,91	0
2	NAG	A	696	14/15	0.36	-	95,96,97,97	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	697	14/15	0.84	14.03	116,117,118,119	0
6	ACT	A	801	4/4	0.37	8.96	32,33,34,35	0
6	ACT	B	801	4/4	0.39	8.45	52,52,53,53	0
6	ACT	A	800	4/4	0.44	8.16	49,51,52,53	0
3	NAG	B	695	14/15	0.39	5.33	76,78,80,83	0
7	GOL	A	2434	6/6	0.24	4.93	29,31,32,33	0
3	NAG	A	695	14/15	0.33	4.86	75,76,79,79	0
7	GOL	B	2433	6/6	0.24	4.02	31,34,35,39	0
3	NAG	B	692	14/15	0.34	2.14	83,86,86,87	0
3	NAG	A	692	14/15	0.32	1.32	66,68,70,71	0
5	CL	B	703	1/1	0.12	-0.87	29,29,29,29	0
4	ZN	B	701	1/1	0.17	-1.30	29,29,29,29	0
5	CL	A	702	1/1	0.13	-1.36	37,37,37,37	0
4	ZN	A	701	1/1	0.13	-1.92	31,31,31,31	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.