



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 03:48 AM GMT

PDB ID : 3CB3  
Title : Crystal structure of L-Talarate dehydratase from Polaromonas sp. JS666 complexed with Mg and L-glucarate  
Authors : Fedorov, A.A.; Fedorov, E.V.; Yew, W.S.; Burley, S.K.; Gerlt, J.A.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2008-02-21  
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	4888 (2.00-2.00)
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	393	
1	B	393	
1	C	393	
1	D	393	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	LGT	B	601	-	X
3	LGT	C	601	-	X
3	LGT	D	601	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11715 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mandelate racemase/muconate lactonizing enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	0	0	0
			2867	1821	513	520	13			
1	B	361	Total	C	N	O	S	0	0	0
			2777	1766	495	503	13			
1	C	357	Total	C	N	O	S	0	0	0
			2750	1748	491	498	13			
1	D	358	Total	C	N	O	S	0	0	0
			2757	1753	492	499	13			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q12GE3
A	0	SER	-	EXPRESSION TAG	UNP Q12GE3
A	1	LEU	-	EXPRESSION TAG	UNP Q12GE3
A	384	GLU	-	EXPRESSION TAG	UNP Q12GE3
A	385	GLY	-	EXPRESSION TAG	UNP Q12GE3
A	386	HIS	-	EXPRESSION TAG	UNP Q12GE3
A	387	HIS	-	EXPRESSION TAG	UNP Q12GE3
A	388	HIS	-	EXPRESSION TAG	UNP Q12GE3
A	389	HIS	-	EXPRESSION TAG	UNP Q12GE3
A	390	HIS	-	EXPRESSION TAG	UNP Q12GE3
A	391	HIS	-	EXPRESSION TAG	UNP Q12GE3
B	-1	MET	-	EXPRESSION TAG	UNP Q12GE3
B	0	SER	-	EXPRESSION TAG	UNP Q12GE3
B	1	LEU	-	EXPRESSION TAG	UNP Q12GE3
B	384	GLU	-	EXPRESSION TAG	UNP Q12GE3
B	385	GLY	-	EXPRESSION TAG	UNP Q12GE3
B	386	HIS	-	EXPRESSION TAG	UNP Q12GE3
B	387	HIS	-	EXPRESSION TAG	UNP Q12GE3
B	388	HIS	-	EXPRESSION TAG	UNP Q12GE3
B	389	HIS	-	EXPRESSION TAG	UNP Q12GE3
B	390	HIS	-	EXPRESSION TAG	UNP Q12GE3

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	391	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	-1	MET	-	EXPRESSION TAG	UNP Q12GE3
C	0	SER	-	EXPRESSION TAG	UNP Q12GE3
C	1	LEU	-	EXPRESSION TAG	UNP Q12GE3
C	384	GLU	-	EXPRESSION TAG	UNP Q12GE3
C	385	GLY	-	EXPRESSION TAG	UNP Q12GE3
C	386	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	387	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	388	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	389	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	390	HIS	-	EXPRESSION TAG	UNP Q12GE3
C	391	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	-1	MET	-	EXPRESSION TAG	UNP Q12GE3
D	0	SER	-	EXPRESSION TAG	UNP Q12GE3
D	1	LEU	-	EXPRESSION TAG	UNP Q12GE3
D	384	GLU	-	EXPRESSION TAG	UNP Q12GE3
D	385	GLY	-	EXPRESSION TAG	UNP Q12GE3
D	386	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	387	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	388	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	389	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	390	HIS	-	EXPRESSION TAG	UNP Q12GE3
D	391	HIS	-	EXPRESSION TAG	UNP Q12GE3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	2	Total Mg 2 2	0	0
2	A	2	Total Mg 2 2	0	0
2	D	2	Total Mg 2 2	0	0
2	C	2	Total Mg 2 2	0	0

- Molecule 3 is L-GLUCARIC ACID (three-letter code: LGT) (formula: C<sub>6</sub>H<sub>10</sub>O<sub>8</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			14	6	8		
3	B	1	Total	C	O	0	0
			14	6	8		
3	C	1	Total	C	O	0	0
			14	6	8		
3	D	1	Total	C	O	0	0
			14	6	8		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	141	Total	O	0	0
			141	141		
4	B	123	Total	O	0	0
			123	123		
4	C	107	Total	O	0	0
			107	107		
4	D	129	Total	O	0	0
			129	129		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Mandelate racemase/muconate lactonizing enzyme

Chain A:




- Molecule 1: Mandelate racemase/muconate lactonizing enzyme


Chain B:



- Molecule 1: Mandelate racemase/muconate lactonizing enzyme


Chain C:





● Molecule 1: Mandelate racemase/muconate lactonizing enzyme

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	195.47Å 84.74Å 118.25Å 90.00° 126.04° 90.00°	Depositor
Resolution (Å)	24.91 – 2.00 39.51 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.7 (24.91-2.00) 94.6 (39.51-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.69 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.210 , 0.239 0.207 , 0.233	Depositor DCC
$R_{free}$ test set	5030 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 36.9	EDS
Estimated twinning fraction	0.022 for -h-2*1,-k,l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 103173 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	11715	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, LGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/2938	0.57	0/3991
1	B	0.33	0/2846	0.57	0/3866
1	C	0.32	0/2818	0.58	0/3826
1	D	0.33	0/2826	0.58	0/3837
All	All	0.33	0/11428	0.57	0/15520

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2867	0	2851	52	0
1	B	2777	0	2752	89	0
1	C	2750	0	2722	77	0
1	D	2757	0	2730	57	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	14	0	7	0	0
3	B	14	0	7	0	0
3	C	14	0	7	2	0
3	D	14	0	7	0	0
4	A	141	0	0	4	0
4	B	123	0	0	8	0
4	C	107	0	0	5	0
4	D	129	0	0	7	0
All	All	11715	0	11083	267	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

The worst 5 of 267 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:358:ARG:HH11	1:A:358:ARG:HB2	1.24	1.03
1:C:214:GLN:HE21	1:C:242:ALA:H	1.10	0.98
1:B:214:GLN:HE21	1:B:242:ALA:H	1.21	0.89
1:B:192:ILE:HD13	1:B:227:ILE:HG21	1.60	0.83
1:A:53:GLU:HG2	1:A:59:GLN:HG2	1.60	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/393 (94%)	364 (98%)	6 (2%)	1 (0%)	50	44
1	B	357/393 (91%)	346 (97%)	9 (2%)	2 (1%)	33	24
1	C	353/393 (90%)	341 (97%)	11 (3%)	1 (0%)	50	44
1	D	354/393 (90%)	347 (98%)	7 (2%)	0	100	100
All	All	1435/1572 (91%)	1398 (97%)	33 (2%)	4 (0%)	50	44

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	315	ALA
1	C	315	ALA
1	A	315	ALA
1	B	161	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	291/310 (94%)	287 (99%)	4 (1%)	78	81
1	B	281/310 (91%)	279 (99%)	2 (1%)	91	93
1	C	278/310 (90%)	275 (99%)	3 (1%)	84	86
1	D	279/310 (90%)	275 (99%)	4 (1%)	78	81
All	All	1129/1240 (91%)	1116 (99%)	13 (1%)	82	84

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	314	PHE
1	C	209	MET
1	D	190	LEU
1	B	209	MET
1	D	162	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	319	HIS
1	C	213	ASN
1	D	248	HIS
1	B	343	ASN
1	B	378	GLN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	LGT	A	601	2	13,13,13	1.31	1 (7%)	18,18,18	0.99	2 (11%)
3	LGT	B	601	2	13,13,13	1.38	2 (15%)	18,18,18	0.97	2 (11%)
3	LGT	C	601	2	13,13,13	1.34	2 (15%)	18,18,18	1.04	2 (11%)
3	LGT	D	601	2	13,13,13	1.23	1 (7%)	18,18,18	0.98	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	LGT	A	601	2	-	0/20/20/20	0/0/0/0
3	LGT	B	601	2	-	0/20/20/20	0/0/0/0
3	LGT	C	601	2	-	0/20/20/20	0/0/0/0
3	LGT	D	601	2	-	0/20/20/20	0/0/0/0

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	LGT	C2-C1	2.78	1.56	1.52
3	C	601	LGT	C5-C6	2.55	1.56	1.52
3	B	601	LGT	C5-C6	2.41	1.56	1.52
3	B	601	LGT	C2-C1	2.36	1.56	1.52
3	C	601	LGT	C2-C1	2.21	1.55	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	LGT	C3-C2-C1	2.64	112.54	109.21
3	C	601	LGT	C4-C5-C6	2.57	112.45	109.21
3	D	601	LGT	C4-C5-C6	2.42	112.26	109.21
3	B	601	LGT	C3-C2-C1	2.42	112.26	109.21
3	A	601	LGT	C3-C2-C1	2.35	112.18	109.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	373/393 (94%)	-0.30	1 (0%) 91 93	15, 25, 40, 51	0
1	B	361/393 (91%)	-0.04	13 (3%) 41 40	14, 26, 51, 62	0
1	C	357/393 (90%)	-0.03	13 (3%) 41 40	15, 27, 52, 61	0
1	D	358/393 (91%)	-0.32	2 (0%) 86 88	15, 25, 41, 56	0
All	All	1449/1572 (92%)	-0.17	29 (2%) 62 62	14, 26, 46, 62	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	158	HIS	5.4
1	C	157	LEU	4.4
1	B	158	HIS	3.8
1	A	102	TRP	3.8
1	B	102	TRP	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	LGT	D	601	14/14	0.21	2.07	29,38,47,48	0
3	LGT	C	601	14/14	0.24	2.03	34,41,49,51	0
3	LGT	B	601	14/14	0.26	2.00	37,43,47,48	0
2	MG	B	503	1/1	0.13	1.59	28,28,28,28	0
3	LGT	A	601	14/14	0.13	0.80	32,39,45,46	0
2	MG	C	502	1/1	0.10	0.38	32,32,32,32	0
2	MG	B	502	1/1	0.08	-1.08	31,31,31,31	0
2	MG	A	502	1/1	0.04	-1.55	21,21,21,21	0
2	MG	C	501	1/1	0.06	-1.85	36,36,36,36	0
2	MG	D	502	1/1	0.06	-1.88	29,29,29,29	0
2	MG	D	503	1/1	0.04	-2.50	21,21,21,21	0
2	MG	A	501	1/1	0.05	-4.52	23,23,23,23	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.