



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 27, 2014 – 12:22 AM GMT

PDB ID : 3CBF  
Title : Crystal structure of LysN, alpha-aminoadipate aminotransferase, from *Thermus thermophilus* HB27  
Authors : Tomita, T.; Miyazaki, T.; Miyagawa, T.; Fushinobu, S.; Kuzuyama, T.; Nishiyama, M.  
Deposited on : 2008-02-21  
Resolution : 1.67 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3587 (1.70-1.66)
Clashscore	79885	4225 (1.70-1.66)
Ramachandran outliers	78287	4144 (1.70-1.66)
Sidechain outliers	78261	4143 (1.70-1.66)
RSRZ outliers	66119	3587 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	397	
1	B	397	

## 2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 6639 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-aminodipate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	6	0
			3086	1987	542	548	9			
1	B	392	Total	C	N	O	S	0	3	0
			3070	1970	542	549	9			

- Molecule 2 is (2S)-2-[(3-HYDROXY-2-METHYL-5-[(PHOSPHONOOXY)METHYL]PYRIDIN-4-YL)METHYL]AMINO]HEXANEDIOICACID (three-letter code: N5F) (formula: C<sub>14</sub>H<sub>21</sub>N<sub>2</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			26	14	2	9	1		
2	B	1	Total	C	N	O	P	0	0
			26	14	2	9	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total 211	O 211	0	0
3	B	220	Total 220	O 220	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alpha-aminodipate aminotransferase

Chain A: 



- Molecule 1: Alpha-aminodipate aminotransferase

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.28Å 93.24Å 150.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.69 – 1.67 34.68 – 1.67	Depositor EDS
% Data completeness (in resolution range)	96.8 (34.69-1.67) 96.8 (34.68-1.67)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.93 (at 1.67Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.202 , 0.247 0.201 , 0.246	Depositor DCC
$R_{free}$ test set	4415 reflections (5.27%)	DCC
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtriage
Anisotropy	0.128	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 42.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 88140 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6639	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.50 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2938e-03.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: N5F

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.61	0/3170	0.75	4/4283 (0.1%)
1	B	0.62	0/3146	0.71	2/4252 (0.0%)
All	All	0.61	0/6316	0.73	6/8535 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	367	LEU	CA-CB-CG	6.92	131.21	115.30
1	A	80	ARG	NE-CZ-NH2	6.64	123.62	120.30
1	B	390	LEU	CA-CB-CG	5.81	128.66	115.30
1	A	80	ARG	NE-CZ-NH1	-5.81	117.40	120.30
1	A	61	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	B	188	LEU	CB-CG-CD2	-5.02	102.47	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3086	0	3182	46	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3070	0	3137	44	0
2	A	26	0	16	0	0
2	B	26	0	17	1	0
3	A	211	0	0	4	0
3	B	220	0	0	12	0
All	All	6639	0	6352	86	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (86) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:105:ASP:OD2	3:B:773:HOH:O	1.61	1.14
1:A:140:LEU:HD11	1:A:161:GLU:HG2	1.33	1.06
1:A:140:LEU:CD1	1:A:161:GLU:HG2	2.01	0.89
1:B:46:LEU:HD11	1:B:296:VAL:HG21	1.56	0.86
1:B:272[A]:MET:HB2	3:B:693:HOH:O	1.79	0.81
1:B:133:ARG:NH1	3:B:639:HOH:O	2.19	0.74
1:B:272[B]:MET:HB2	3:B:693:HOH:O	1.90	0.71
1:A:29:THR:HG22	1:A:34:ILE:HD11	1.71	0.71
1:B:314:LYS:HE3	3:B:757:HOH:O	1.90	0.70
1:A:274:ASN:H	1:A:274:ASN:HD22	1.41	0.69
1:A:223:ARG:HD2	3:A:581:HOH:O	1.93	0.68
1:B:31:ARG:HD2	3:B:658:HOH:O	1.95	0.66
1:B:107:VAL:HG11	1:B:200:VAL:HG21	1.78	0.66
1:A:168:LEU:CD1	1:A:188:LEU:HD21	2.27	0.65
1:A:261:GLN:HE21	1:B:17:ILE:HG22	1.61	0.64
1:A:112:LEU:HD21	1:A:118[A]:VAL:HG13	1.79	0.64
1:B:5:SER:N	3:B:779:HOH:O	2.31	0.62
1:B:343:ARG:NE	1:B:393:LEU:HD13	2.14	0.62
1:B:274:ASN:H	1:B:274:ASN:HD22	1.46	0.62
1:A:112:LEU:CD2	1:A:118[A]:VAL:HG13	2.33	0.59
1:A:168:LEU:HD13	1:A:188:LEU:HD21	1.85	0.58
1:A:134:LEU:HD11	1:B:109:LYS:HD3	1.85	0.58
1:B:172:PHE:CE1	1:B:179:LEU:HG	2.39	0.58
1:B:223:ARG:NH2	3:B:788:HOH:O	2.18	0.57
1:B:322:LYS:HE2	3:B:616:HOH:O	2.03	0.57
1:A:28:LEU:HD21	1:A:345:LEU:HD21	1.87	0.56
1:B:274:ASN:HD22	1:B:274:ASN:N	2.03	0.56
1:B:168:LEU:HD13	1:B:188:LEU:HD21	1.87	0.56
1:A:306:HIS:HD2	1:A:310:ARG:HH11	1.53	0.54
1:B:119:LEU:HD11	1:B:154:LEU:CD1	2.38	0.54

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:63:LYS:HE3	3:B:790:HOH:O	2.07	0.54
1:B:269:HIS:HD2	1:B:270:THR:O	1.91	0.53
1:A:274:ASN:N	1:A:274:ASN:HD22	2.07	0.53
1:B:22:ILE:HD12	1:B:126:MET:HG2	1.91	0.53
1:B:146:GLU:HG3	3:B:616:HOH:O	2.10	0.52
1:A:287:SER:O	1:A:291:GLU:HG3	2.10	0.51
1:A:343:ARG:NE	1:A:393:LEU:HD13	2.26	0.51
1:A:364:GLU:CD	1:A:364:GLU:H	2.14	0.51
1:A:118[A]:VAL:HG22	1:A:137:PRO:HB2	1.92	0.51
1:A:70:TYR:CZ	1:B:242:PRO:HB3	2.46	0.50
1:A:10:PHE:HB3	1:A:14:ALA:HB3	1.93	0.50
1:A:112:LEU:HD21	1:A:118[A]:VAL:CG1	2.42	0.50
1:A:29:THR:HA	1:A:34:ILE:HG12	1.95	0.49
1:A:40:GLY:HA3	1:A:325[B]:MET:HE1	1.95	0.49
1:B:376:ARG:NH1	1:B:376:ARG:HB3	2.27	0.49
1:B:376:ARG:HB3	1:B:376:ARG:HH11	1.78	0.49
1:A:305:LEU:HD11	1:A:321:PRO:HD2	1.93	0.49
1:B:187:ARG:NE	3:B:763:HOH:O	2.30	0.48
1:A:40:GLY:HA3	1:A:325[B]:MET:CE	2.44	0.47
1:B:120:LEU:O	1:B:141:THR:HA	2.13	0.47
1:B:70:TYR:HA	1:B:269:HIS:HE1	1.80	0.47
1:B:187:ARG:O	1:B:191:MET:HG3	2.14	0.47
1:B:28:LEU:HD21	1:B:345:LEU:HD21	1.96	0.46
1:B:343:ARG:HD2	1:B:393:LEU:HD22	1.98	0.45
1:B:10:PHE:HB3	1:B:14:ALA:HB3	1.98	0.45
1:A:261:GLN:HB2	1:B:17:ILE:CG2	2.47	0.45
1:A:269:HIS:HD2	1:A:270:THR:O	1.99	0.45
1:A:274:ASN:ND2	1:A:274:ASN:N	2.65	0.44
1:A:119:LEU:HD12	1:A:157:VAL:HG11	1.99	0.44
1:A:172:PHE:CE1	1:A:179[B]:LEU:HD13	2.52	0.44
1:B:43:ALA:HB2	1:B:300:LYS:HE3	2.01	0.43
1:A:256:LEU:HD13	1:A:260:VAL:HG23	2.01	0.43
2:B:501:N5F:N	2:B:501:N5F:OAN	2.52	0.43
1:B:70:TYR:HA	1:B:269:HIS:CE1	2.54	0.43
1:B:330:GLU:HG2	1:B:366:THR:HG22	2.01	0.42
1:A:31:ARG:HG2	3:A:699:HOH:O	2.18	0.42
1:A:289:ARG:NH1	3:A:694:HOH:O	2.35	0.42
1:A:169:ILE:HG12	1:A:202:ASP:HB3	2.01	0.42
1:B:274:ASN:ND2	1:B:274:ASN:N	2.66	0.42
1:B:10:PHE:HB3	1:B:14:ALA:CB	2.49	0.42
1:A:174:ASN:HA	1:A:175:PRO:HA	1.89	0.42
1:A:282:LEU:HG	1:A:286:PHE:CD1	2.54	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:121:GLU:HB3	1:A:144:ALA:HB2	2.02	0.42
1:B:256:LEU:O	1:B:260:VAL:HG23	2.20	0.42
1:A:80:ARG:HD2	1:A:94:VAL:O	2.19	0.41
1:A:134:LEU:HD11	1:B:109:LYS:CD	2.49	0.41
1:A:325[B]:MET:HB3	1:A:372:ALA:HB2	2.02	0.41
1:A:43:ALA:HA	1:A:44:PRO:HD3	1.94	0.41
1:B:343:ARG:HE	1:B:393:LEU:HD13	1.85	0.41
1:A:306:HIS:CD2	1:A:310:ARG:HH11	2.36	0.41
1:A:80:ARG:CD	1:A:94:VAL:O	2.69	0.41
1:A:300[A]:LYS:NZ	3:A:643:HOH:O	2.54	0.41
1:A:343:ARG:HE	1:A:393:LEU:HD13	1.85	0.40
1:B:282:LEU:HD22	1:B:286:PHE:CD1	2.57	0.40
1:A:274:ASN:ND2	1:A:274:ASN:H	2.13	0.40
1:B:182:LEU:HG	1:B:186:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/397 (100%)	384 (97%)	11 (3%)	1 (0%)	50	27
1	B	393/397 (99%)	382 (97%)	11 (3%)	0	100	100
All	All	789/794 (99%)	766 (97%)	22 (3%)	1 (0%)	59	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of

similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	317/316 (100%)	287 (90%)	30 (10%)	12 2
1	B	314/316 (99%)	300 (96%)	14 (4%)	38 13
All	All	631/632 (100%)	587 (93%)	44 (7%)	24 5

All (44) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	SER
1	A	30	GLN
1	A	34	ILE
1	A	45	GLU
1	A	46	LEU
1	A	79	LEU
1	A	118[A]	VAL
1	A	118[B]	VAL
1	A	120[A]	LEU
1	A	120[B]	LEU
1	A	134	LEU
1	A	158	LEU
1	A	162	ARG
1	A	175	PRO
1	A	179[A]	LEU
1	A	179[B]	LEU
1	A	256	LEU
1	A	274	ASN
1	A	282	LEU
1	A	290	LEU
1	A	316	VAL
1	A	322	LYS
1	A	333	LYS
1	A	335	LEU
1	A	340	LEU
1	A	345	LEU
1	A	364	GLU
1	A	367	LEU
1	A	386	LEU
1	A	394	LEU
1	B	85	GLU
1	B	120	LEU
1	B	134	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	158	LEU
1	B	160	ARG
1	B	179	LEU
1	B	274	ASN
1	B	314	LYS
1	B	333	LYS
1	B	335	LEU
1	B	345	LEU
1	B	367	LEU
1	B	390	LEU
1	B	394	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	101	GLN
1	A	135	GLN
1	A	261	GLN
1	A	269	HIS
1	A	274	ASN
1	A	302	GLN
1	A	306	HIS
1	A	360	ASN
1	B	30	GLN
1	B	101	GLN
1	B	135	GLN
1	B	190	GLN
1	B	261	GLN
1	B	269	HIS
1	B	274	ASN
1	B	360	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	N5F	A	500	-	26,26,26	1.12	1 (3%)	36,36,36	1.48	7 (19%)
2	N5F	B	501	-	26,26,26	1.35	2 (7%)	36,36,36	1.28	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	N5F	A	500	-	-	0/21/21/21	0/1/1/1
2	N5F	B	501	-	-	0/21/21/21	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	N5F	CAK-CAP	4.39	1.57	1.51
2	A	500	N5F	CAK-CAP	4.00	1.56	1.51
2	B	501	N5F	CA-N	2.88	1.51	1.47

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	N5F	CAK-N-CA	3.74	122.17	114.34
2	B	501	N5F	OAM-PAT-OAS	-3.30	97.53	106.65
2	A	500	N5F	OAM-PAT-OAS	-2.83	98.85	106.65
2	A	500	N5F	CAK-CAP-CAO	2.78	123.83	120.31

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	N5F	CAK-CAP-CAQ	2.41	121.91	119.70
2	A	500	N5F	OAB-CAI-CAH	-2.30	115.11	123.03
2	A	500	N5F	CAY-CAQ-CAP	2.15	119.73	118.10
2	A	500	N5F	CB-CA-C	2.10	115.22	110.71
2	A	500	N5F	CAQ-CAY-NAX	-2.04	120.18	123.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	392/397 (98%)	0.23	24 (6%) 21 23	13, 22, 35, 46	0
1	B	392/397 (98%)	0.11	14 (3%) 41 45	13, 21, 34, 41	0
All	All	784/794 (98%)	0.17	38 (4%) 29 32	13, 22, 35, 46	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	284	GLU	4.2
1	A	32	PRO	3.9
1	A	333	LYS	3.9
1	A	310	ARG	3.9
1	B	32	PRO	3.7
1	A	376	ARG	3.6
1	B	31	ARG	3.4
1	B	17	ILE	3.3
1	B	18	GLN	3.3
1	A	8	GLU	3.0
1	A	335	LEU	3.0
1	B	310	ARG	2.7
1	A	31	ARG	2.7
1	A	334	GLY	2.7
1	A	314	LYS	2.7
1	A	160	ARG	2.6
1	B	160	ARG	2.5
1	A	247	ALA	2.5
1	A	246	VAL	2.5
1	A	284	GLU	2.5
1	A	394	LEU	2.4
1	A	45	GLU	2.4
1	B	342	ARG	2.4
1	B	28	LEU	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	335	LEU	2.3
1	B	247	ALA	2.3
1	B	314	LYS	2.3
1	A	18	GLN	2.2
1	B	334	GLY	2.2
1	B	246	VAL	2.2
1	A	17	ILE	2.2
1	A	33	GLY	2.1
1	A	295	ARG	2.1
1	A	342	ARG	2.1
1	A	212	GLU	2.0
1	A	190	GLN	2.0
1	A	244	LEU	2.0
1	A	345	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	N5F	A	500	26/26	0.12	-0.07	14,18,20,21	0
2	N5F	B	501	26/26	0.10	-0.42	13,17,20,21	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.