



# wwPDB X-ray Structure Validation Summary Report i

Jun 19, 2014 – 05:20 PM EDT

PDB ID : 4CCJ  
Title : 60S ribosomal protein L8 histidine hydroxylase (NO66) in apo form  
Authors : Chowdhury, R.; Ge, W.; Clifton, I.J.; Schofield, C.J.  
Deposited on : 2013-10-23  
Resolution : 2.15 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : **FAILED**  
Xtriage (Phenix) : dev-1439  
EDS : stable23161  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23161

# 1 Overall quality at a glance

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1094 (2.18-2.14)
Clashscore	79885	1299 (2.18-2.14)
Ramachandran outliers	78287	1272 (2.18-2.14)
Sidechain outliers	78261	1272 (2.18-2.14)
RSRZ outliers	66119	1094 (2.18-2.14)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	467	
1	B	467	
1	C	467	
1	D	467	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	EDO	B	1640	-	X
2	EDO	C	1640	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15725 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BIFUNCTIONAL LYSINE-SPECIFIC DEMETHYLASE AND HISTIDYL-HYDROXYLASE NO66.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	458	Total 3718	C 2365	N 658	O 678	S 17	0	9	0
1	B	458	Total 3698	C 2353	N 656	O 672	S 17	0	7	0
1	C	458	Total 3690	C 2351	N 650	O 673	S 16	0	5	0
1	D	457	Total 3664	C 2334	N 647	O 667	S 16	0	4	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	MET	-	EXPRESSION TAG	UNP Q9H6W3
A	364	ALA	VAL	ENGINEERED MUTATION	UNP Q9H6W3
B	182	MET	-	EXPRESSION TAG	UNP Q9H6W3
B	364	ALA	VAL	ENGINEERED MUTATION	UNP Q9H6W3
C	182	MET	-	EXPRESSION TAG	UNP Q9H6W3
C	364	ALA	VAL	ENGINEERED MUTATION	UNP Q9H6W3
D	182	MET	-	EXPRESSION TAG	UNP Q9H6W3
D	364	ALA	VAL	ENGINEERED MUTATION	UNP Q9H6W3

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	329	Total O 329 329	0	0
3	B	288	Total O 288 288	0	0
3	C	173	Total O 173 173	0	0
3	D	153	Total O 153 153	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: BIFUNCTIONAL LYSINE-SPECIFIC DEMETHYLASE AND HISTIDYL-HYDROXYLASE NO66

Chain A: 




#### • Molecule 1: BIFUNCTIONAL LYSINE-SPECIFIC DEMETHYLASE AND HISTIDYL-HYDROXYLASE NO66


Chain B: 



#### • Molecule 1: BIFUNCTIONAL LYSINE-SPECIFIC DEMETHYLASE AND HISTIDYL-HYDROXYLASE NO66


Chain C: 





● Molecule 1: BIFUNCTIONAL LYSINE-SPECIFIC DEMETHYLASE AND HISTIDYL-HYDROXYLASE NO66

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.44Å 81.56Å 151.96Å 90.00° 94.66° 90.00°	Depositor
Resolution (Å)	63.72 – 2.15 63.72 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.1 (63.72-2.15) 94.9 (63.72-2.15)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.16Å)	Xtrriage
Refinement program	CNS 1.3	Depositor
R, $R_{free}$	0.202 , 0.210 0.221 , 0.225	Depositor DCC
$R_{free}$ test set	6443 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtrriage
Anisotropy	0.385	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 41.8	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 128267 reflections	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15725	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3840	0.62	0/5222
1	B	0.40	0/3814	0.64	1/5190 (0.0%)
1	C	0.36	0/3802	0.62	3/5174 (0.1%)
1	D	0.36	0/3770	0.57	1/5129 (0.0%)
All	All	0.37	0/15226	0.61	5/20715 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	4
1	D	0	2
All	All	0	6

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	611	LEU	CB-CA-C	5.58	120.81	110.20
1	D	411	ASP	N-CA-C	5.56	126.00	111.00
1	C	234	ASP	N-CA-CB	-5.51	100.67	110.60
1	B	259	GLY	N-CA-C	5.38	126.54	113.10
1	C	411	ASP	CB-CA-C	-5.10	100.21	110.40

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	517[A]	ARG	Mainchain
1	C	517[B]	ARG	Mainchain
1	C	530[A]	TRP	Mainchain
1	C	530[B]	TRP	Mainchain
1	D	187[A]	ARG	Mainchain

## 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3718	0	3652	61	0
1	B	3698	0	3637	67	0
1	C	3690	0	3605	92	0
1	D	3664	0	3579	90	0
2	B	4	0	6	1	0
2	C	4	0	6	0	0
2	D	4	0	6	0	0
3	A	329	0	0	4	0
3	B	288	0	0	5	0
3	C	173	0	0	1	0
3	D	153	0	0	1	0
All	All	15725	0	14491	285	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 10.

The worst 5 of 285 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:272:ARG:HH12	1:B:274:THR:HG22	1.20	1.01
1:B:455:PRO:HG2	1:B:458:PHE:HB3	1.43	1.01
1:C:263:ASP:HA	1:C:276:ASN:HD21	1.36	0.89
1:A:291:GLN:HE21	1:B:538:VAL:HG11	1.38	0.87
1:C:549:HIS:O	1:C:632:LEU:HD12	1.75	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/467 (100%)	459 (99%)	5 (1%)	1 (0%)	56	55
1	B	463/467 (99%)	450 (97%)	13 (3%)	0	100	100
1	C	461/467 (99%)	444 (96%)	17 (4%)	0	100	100
1	D	457/467 (98%)	443 (97%)	14 (3%)	0	100	100
All	All	1846/1868 (99%)	1796 (97%)	49 (3%)	1 (0%)	59	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	538	VAL

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	396/399 (99%)	389 (98%)	7 (2%)	71	77
1	B	394/399 (99%)	385 (98%)	9 (2%)	63	68
1	C	390/399 (98%)	385 (99%)	5 (1%)	80	87
1	D	386/399 (97%)	380 (98%)	6 (2%)	75	81
All	All	1566/1596 (98%)	1539 (98%)	27 (2%)	75	79

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	435	ILE
1	B	603	PRO
1	D	468	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	538	VAL
1	A	590	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	590	GLN
1	C	619	GLN
1	D	417	HIS
1	C	441	GLN
1	D	537	ASN

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

Mogul failed to run properly - this section will therefore be empty.

## 5.5 Carbohydrates [i](#)

Mogul failed to run properly - this section will therefore be empty.

## 5.6 Ligand geometry [i](#)

Mogul failed to run properly - this section will therefore be empty.

## 5.7 Other polymers [i](#)

Mogul failed to run properly - this section will therefore be empty.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	458/467 (98%)	0.07	6 (1%) 74 79	26, 38, 65, 80	3 (0%)
1	B	458/467 (98%)	0.06	7 (1%) 70 76	26, 41, 68, 81	2 (0%)
1	C	458/467 (98%)	0.35	27 (5%) 22 25	25, 51, 93, 100	3 (0%)
1	D	457/467 (97%)	0.24	20 (4%) 33 36	25, 57, 84, 100	1 (0%)
All	All	1831/1868 (98%)	0.18	60 (3%) 44 48	25, 45, 79, 100	9 (0%)

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	538	VAL	6.5
1	C	530[A]	TRP	5.7
1	C	537	ASN	5.2
1	D	616	VAL	5.0
1	C	542	LEU	4.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q < 0.9
2	EDO	B	1640	4/4	0.18	8.35	33,38,39,39	0
2	EDO	C	1640	4/4	0.15	3.12	42,47,48,50	0
2	EDO	D	1640	4/4	0.16	1.98	49,52,53,53	0

## 6.5 Other polymers

There are no such residues in this entry.