



# wwPDB X-ray Structure Validation Summary Report

Mar 12, 2014 – 04:58 PM GMT

PDB ID : 4CE7  
Title : Crystal structure of a novel unsaturated beta-glucuronyl hydrolase enzyme, belonging to family GH105, involved in ulvan degradation  
Authors : Nyvall Collen, P.; Jeudy, A.; Groisillier, A.; Coutinho, P.M.; Helbert, W.; Czjzek, M.  
Deposited on : 2013-11-09  
Resolution : 1.90 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1323  
EDS : trunk22714  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk22714

# 1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	370	
1	B	370	
1	C	370	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NO3	C	1363	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9354 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UNSATURATED 3S-RHAMNOGLYCURONYL HYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	345	Total	C	N	O	S	0	3	0
			2868	1841	479	535	13			
1	B	348	Total	C	N	O	S	0	2	0
			2873	1843	480	537	13			
1	C	342	Total	C	N	O	S	0	5	0
			2855	1832	477	533	13			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-6	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-5	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-4	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-3	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-2	HIS	-	EXPRESSION TAG	UNP L7P9J4
A	-1	GLY	-	EXPRESSION TAG	UNP L7P9J4
A	0	SER	-	EXPRESSION TAG	UNP L7P9J4
B	-7	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-6	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-5	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-4	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-3	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-2	HIS	-	EXPRESSION TAG	UNP L7P9J4
B	-1	GLY	-	EXPRESSION TAG	UNP L7P9J4
B	0	SER	-	EXPRESSION TAG	UNP L7P9J4
C	-7	HIS	-	EXPRESSION TAG	UNP L7P9J4
C	-6	HIS	-	EXPRESSION TAG	UNP L7P9J4
C	-5	HIS	-	EXPRESSION TAG	UNP L7P9J4
C	-4	HIS	-	EXPRESSION TAG	UNP L7P9J4
C	-3	HIS	-	EXPRESSION TAG	UNP L7P9J4
C	-2	HIS	-	EXPRESSION TAG	UNP L7P9J4

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	EXPRESSION TAG	UNP L7P9J4
C	0	SER	-	EXPRESSION TAG	UNP L7P9J4

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	C	1	Total	N	O	0	0
			4	1	3		
2	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	265	Total	O	0	0
			265	265		
3	B	260	Total	O	0	0
			260	260		
3	C	217	Total	O	0	0
			217	217		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: UNSATURATED 3S-RHAMNOGLYCURONYL HYDROLASE

Chain A: 




#### • Molecule 1: UNSATURATED 3S-RHAMNOGLYCURONYL HYDROLASE

Chain B: 



#### • Molecule 1: UNSATURATED 3S-RHAMNOGLYCURONYL HYDROLASE

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.98Å 93.33Å 156.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.19 – 1.90 34.81 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.19-1.90) 98.9 (34.81-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.162 , 0.207 0.164 , 0.209	Depositor DCC
$R_{free}$ test set	5313 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtriage
Anisotropy	0.040	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 39.0	EDS
Estimated twinning fraction	0.023 for k,h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 106771 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9354	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NO3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.09	7/2962 (0.2%)	1.03	14/4017 (0.3%)
1	B	1.07	6/2967 (0.2%)	0.95	9/4025 (0.2%)
1	C	1.08	7/2953 (0.2%)	1.02	16/4004 (0.4%)
All	All	1.08	20/8882 (0.2%)	1.00	39/12046 (0.3%)

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	TRP	CD2-CE2	7.56	1.50	1.41
1	B	145	TRP	CD2-CE2	7.33	1.50	1.41
1	A	223	TRP	CD2-CE2	7.13	1.50	1.41
1	C	79	TRP	CD2-CE2	7.02	1.49	1.41
1	C	266	TRP	CD2-CE2	6.75	1.49	1.41

The worst 5 of 39 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	ARG	NE-CZ-NH2	-17.46	111.57	120.30
1	C	183	ARG	NE-CZ-NH2	-15.09	112.76	120.30
1	A	255	ARG	NE-CZ-NH1	11.61	126.10	120.30
1	C	183	ARG	NE-CZ-NH1	10.38	125.49	120.30
1	A	255	ARG	CG-CD-NE	-8.84	93.24	111.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2868	0	52	6	0
1	B	2873	0	12	5	0
1	C	2855	0	31	5	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	8	0	0	0	0
3	A	265	0	0	3	0
3	B	260	0	0	2	1
3	C	217	0	0	2	1
All	All	9354	0	95	15	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

The worst 5 of 15 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:62:LYS:CE	3:A:2064:HOH:O	1.92	1.15
1:A:126:ASP:CB	3:A:2139:HOH:O	2.14	0.93
1:A:189:GLU:OE1	1:A:255:ARG:NH2	2.24	0.70
1:A:335:PRO:HD3	3:A:2263:HOH:O	1.92	0.69
1:B:189:GLU:OE1	1:B:255:ARG:NH2	2.25	0.69

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:B:2103:HOH:O	3:C:2178:HOH:O[3.655]	2.15	0.05

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries



of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/370 (93%)	339 (98%)	5 (2%)	0	100	100
1	B	348/370 (94%)	338 (97%)	10 (3%)	0	100	100
1	C	343/370 (93%)	335 (98%)	7 (2%)	1 (0%)	50	37
All	All	1035/1110 (93%)	1012 (98%)	22 (2%)	1 (0%)	59	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	302	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/317 (94%)	281 (94%)	17 (6%)	29	15
1	B	298/317 (94%)	288 (97%)	10 (3%)	49	36
1	C	297/317 (94%)	283 (95%)	14 (5%)	36	22
All	All	893/951 (94%)	852 (95%)	41 (5%)	37	23

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	127	ASN
1	B	255	ARG
1	C	265	LEU
1	B	135	ARG
1	B	137	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NO3	A	1362	-	3,3,3	2.98	2 (66%)	3,3,3	0.39	0
2	NO3	B	1362	-	3,3,3	2.73	2 (66%)	3,3,3	0.16	0
2	NO3	C	1362	-	3,3,3	2.92	2 (66%)	3,3,3	0.29	0
2	NO3	C	1363	-	3,3,3	2.91	2 (66%)	3,3,3	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NO3	A	1362	-	-	0/0/0/0	0/0/0/0
2	NO3	B	1362	-	-	0/0/0/0	0/0/0/0
2	NO3	C	1362	-	-	0/0/0/0	0/0/0/0
2	NO3	C	1363	-	-	0/0/0/0	0/0/0/0

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1362	NO3	O1-N	4.29	1.42	1.24
2	C	1362	NO3	O1-N	3.97	1.41	1.24
2	C	1363	NO3	O1-N	3.85	1.40	1.24
2	B	1362	NO3	O1-N	3.83	1.40	1.24
2	C	1363	NO3	O3-N	3.23	1.42	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	345/370 (93%)	-0.03	16 (4%) 31 31	16, 26, 45, 89	0
1	B	348/370 (94%)	-0.09	8 (2%) 57 59	17, 25, 41, 57	0
1	C	342/370 (92%)	-0.04	11 (3%) 45 46	15, 25, 48, 76	0
All	All	1035/1110 (93%)	-0.05	35 (3%) 43 44	15, 25, 46, 89	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	202	GLY	5.0
1	C	204	ASP	4.2
1	A	204	ASP	4.0
1	A	336	GLU	3.7
1	C	205	GLU	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NO3	C	1363	4/4	0.13	3.76	61,61,65,70	0
2	NO3	B	1362	4/4	0.13	1.27	64,69,70,74	0
2	NO3	A	1362	4/4	0.14	0.81	58,62,67,70	0
2	NO3	C	1362	4/4	0.10	-0.23	56,64,64,66	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.