



# Full wwPDB X-ray Structure Validation Report

Feb 26, 2014 – 06:19 PM GMT

PDB ID : 2CFA  
Title : Structure of viral flavin-dependant thymidylate synthase ThyX  
Authors : Graziani, S.; Bernauer, J.; Skouloubris, S.; Graille, M.; Zhou, C.-Z.; Marchand, C.; Decottignies, P.; van Tilbeurgh, H.; Myllykallio, H.; Liebl, U.  
Deposited on : 2006-02-17  
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	217	
2	B	217	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3201 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1460	935	252	265	8	0	0	0


- Molecule 2 is a protein called THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	179	1454	932	250	264	8	1	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	77	ILE	LEU	CONFLICT	UNP O41156

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total	O	0	0
			106	106		
4	B	75	Total	O	0	0
			75	75		


### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: THYMIDYLATE SYNTHASE



#### • Molecule 2: THYMIDYLATE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.26Å 76.99Å 93.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 33.65 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.6 (20.00-2.30) 98.6 (33.65-2.30)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.92 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.213 , 0.262 0.207 , 0.259	Depositor DCC
$R_{free}$ test set	1124 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.2	Xtriage
Anisotropy	0.795	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	2 of 22506 reflections (0.009%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3201	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CME, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1482	0.60	0/2000
2	B	0.49	0/1473	0.60	1/1983 (0.1%)
All	All	0.49	0/2955	0.60	1/3983 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	151	LYS	N-CA-CB	7.91	124.83	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1460	0	1456	22	0
2	B	1454	0	1456	34	0
3	A	53	0	31	1	0
3	B	53	0	31	2	0
4	A	106	0	0	2	0
4	B	75	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3201	0	2974	55	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (55) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:98:PRO:C	2:B:208:SER:N	1.74	1.41
1:A:159:PRO:O	1:A:162:THR:HG22	1.50	1.11
2:B:76:VAL:HG11	2:B:168:MET:CE	2.01	0.91
2:B:34:ASN:HB3	2:B:36:ILE:HG22	1.50	0.91
2:B:76:VAL:CG1	2:B:168:MET:HE2	2.01	0.90
2:B:76:VAL:HG11	2:B:168:MET:HE2	1.57	0.86
2:B:66:LEU:HD11	2:B:168:MET:HE3	1.63	0.80
2:B:76:VAL:CG1	2:B:168:MET:CE	2.59	0.79
2:B:66:LEU:CD1	2:B:168:MET:HE3	2.13	0.79
1:A:67:LYS:HD3	1:A:165:THR:HB	1.68	0.75
1:A:1:MET:N	1:A:67:LYS:O	2.20	0.74
2:B:66:LEU:HD11	2:B:168:MET:CE	2.23	0.68
2:B:76:VAL:HG11	2:B:168:MET:HE1	1.76	0.67
2:B:182:ARG:HA	2:B:187:THR:HG21	1.77	0.65
1:A:182:ARG:HA	1:A:187:THR:HG21	1.84	0.60
2:B:66:LEU:CD1	2:B:168:MET:CE	2.80	0.59
2:B:180:GLU:HG3	4:B:2057:HOH:O	2.02	0.59
1:A:29:VAL:O	1:A:29:VAL:HG22	2.03	0.58
2:B:73:ALA:O	2:B:77:ILE:HG12	2.03	0.57
2:B:13:VAL:HG22	2:B:16:VAL:HB	1.85	0.57
2:B:177:HIS:HD2	4:B:2040:HOH:O	1.86	0.57
2:B:174:ASP:OD2	4:B:2055:HOH:O	2.16	0.57
2:B:84:PHE:CE2	2:B:168:MET:HG3	2.40	0.57
1:A:28:ARG:C	1:A:30:SER:H	2.08	0.57
2:B:63:THR:HB	2:B:169:SER:OG	2.05	0.56
2:B:60:ALA:HB3	2:B:172:ILE:HG12	1.87	0.55
2:B:63:THR:CG2	4:B:2008:HOH:O	2.56	0.53
2:B:66:LEU:HD12	2:B:168:MET:HE3	1.89	0.52
2:B:76:VAL:CG1	2:B:168:MET:HE1	2.37	0.51
1:A:28:ARG:O	1:A:31:ASN:OD1	2.27	0.51
1:A:9:THR:HG23	2:B:7:SER:OG	2.10	0.51
2:B:154:ALA:O	2:B:157:ILE:HG12	2.11	0.51
1:A:10:LYS:HD3	2:B:10:LYS:HD3	1.93	0.51
1:A:70:ARG:HD2	1:A:164:THR:HB	1.93	0.50
1:A:7:SER:OG	2:B:9:THR:HG23	2.11	0.50

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:86:GLU:HG2	2:B:166:ILE:HG22	1.93	0.50
2:B:7:SER:HB3	2:B:63:THR:HG23	1.94	0.49
2:B:36:ILE:HG13	2:B:39:LYS:HD2	1.96	0.48
3:A:1219:FAD:H2'	3:A:1219:FAD:H9	1.95	0.48
1:A:182:ARG:HD3	1:A:187:THR:HG21	1.96	0.48
1:A:86:GLU:HG2	1:A:166:ILE:HG22	1.95	0.47
1:A:39:LYS:HB2	4:A:2018:HOH:O	2.15	0.47
1:A:176:ILE:O	1:A:180:GLU:HB2	2.16	0.46
1:A:204:LYS:HB3	1:A:204:LYS:HE3	1.82	0.46
3:B:1217:FAD:H1'2	3:B:1217:FAD:O4'	2.19	0.43
1:A:158:LEU:HB3	1:A:162:THR:HG21	2.01	0.42
3:B:1217:FAD:H9	3:B:1217:FAD:H2'	2.00	0.42
1:A:20:GLU:OE1	4:A:2012:HOH:O	2.22	0.41
1:A:189:ARG:HD3	1:A:189:ARG:N	2.34	0.41
1:A:61:PHE:CE1	1:A:169:SER:HB3	2.55	0.41
1:A:84:PHE:O	2:B:55:SER:HB2	2.19	0.41
2:B:76:VAL:HG12	2:B:168:MET:HE2	1.92	0.41
1:A:50:ARG:HD3	1:A:51:HIS:CE1	2.56	0.41
2:B:52:LYS:HE3	4:B:2032:HOH:O	2.21	0.41
2:B:143:LYS:HB3	4:B:2052:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/217 (80%)	168 (97%)	4 (2%)	1 (1%)	33	39
2	B	169/217 (78%)	169 (100%)	0	0	100	100
All	All	342/434 (79%)	337 (98%)	4 (1%)	1 (0%)	50	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/190 (81%)	138 (90%)	16 (10%)	10	11
2	B	154/190 (81%)	139 (90%)	15 (10%)	12	13
All	All	308/380 (81%)	277 (90%)	31 (10%)	11	12

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	35	GLN
1	A	44	LEU
1	A	50	ARG
1	A	77	LEU
1	A	125	TRP
1	A	132	LYS
1	A	158	LEU
1	A	160	LEU
1	A	162	THR
1	A	165	THR
1	A	172	ILE
1	A	173	ARG
1	A	187	THR
1	A	189	ARG
1	A	194	LEU
2	B	13	VAL
2	B	36	ILE
2	B	44	LEU
2	B	45	LEU
2	B	63	THR
2	B	131	GLU
2	B	132	LYS
2	B	152	GLU
2	B	155	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	158	LEU
2	B	166	ILE
2	B	187	THR
2	B	189	ARG
2	B	194	LEU
2	B	200	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	34	ASN
1	A	35	GLN
1	A	85	GLN
1	A	196	ASN
1	A	217	HIS
2	B	17	ASN
2	B	34	ASN
2	B	136	GLN
2	B	177	HIS
2	B	185	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CME	A	48	1	9,9,10	5.98	2 (22%)	7,9,11	1.14	1 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	CME	B	48	2	9,9,10	5.38	1 (11%)	7,9,11	2.90	4 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	48	1	-	0/6/8/10	0/0/0/0
2	CME	B	48	2	-	0/6/8/10	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	48	CME	O-C	17.71	1.23	1.11
2	B	48	CME	O-C	15.99	1.22	1.11
1	A	48	CME	CA-C	2.16	1.52	1.48

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	48	CME	C-CA-N	-4.81	109.03	113.83
2	B	48	CME	CB-SG-SD	4.33	112.52	103.87
2	B	48	CME	CE-SD-SG	2.59	116.88	103.50
1	A	48	CME	CE-SD-SG	2.54	116.65	103.50
2	B	48	CME	CB-CA-N	2.42	114.37	110.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	A	1219	-	58,58,58	1.05	4 (6%)	85,89,89	1.89	13 (15%)
3	FAD	B	1217	-	58,58,58	1.07	5 (8%)	85,89,89	2.13	20 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	1219	-	-	0/34/50/50	0/1/6/6
3	FAD	B	1217	-	-	0/34/50/50	0/1/6/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1219	FAD	C2A-N3A	3.42	1.38	1.32
3	B	1217	FAD	C2A-N3A	3.16	1.38	1.32
3	A	1219	FAD	C1'-C2'	3.12	1.54	1.51
3	A	1219	FAD	C5X-N5	2.60	1.39	1.35
3	B	1217	FAD	C1'-N10	2.56	1.51	1.48
3	B	1217	FAD	C1'-C2'	2.38	1.53	1.51
3	A	1219	FAD	C2A-N1A	2.37	1.38	1.33
3	B	1217	FAD	C5X-N5	2.30	1.38	1.35
3	B	1217	FAD	C2A-N1A	2.15	1.38	1.33

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1217	FAD	N3A-C2A-N1A	-11.38	119.19	128.71
3	A	1219	FAD	N3A-C2A-N1A	-11.09	119.44	128.71
3	B	1217	FAD	O4B-C1B-N9A	6.72	114.69	108.44
3	A	1219	FAD	O4B-C1B-N9A	5.45	113.51	108.44
3	B	1217	FAD	O4'-C4'-C5'	-5.44	98.95	110.12
3	B	1217	FAD	C2-N1-C10	5.32	120.33	114.98

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1219	FAD	C2-N1-C10	4.66	119.67	114.98
3	A	1219	FAD	N3A-C4A-N9A	4.05	132.75	125.43
3	B	1217	FAD	C1'-N10-C9A	4.02	122.79	118.87
3	B	1217	FAD	N3A-C4A-N9A	3.47	131.69	125.43
3	A	1219	FAD	C1'-N10-C9A	3.42	122.20	118.87
3	B	1217	FAD	C4X-C10-N1	-3.38	119.35	122.73
3	B	1217	FAD	C4X-N5-C5X	2.84	119.88	116.69
3	A	1219	FAD	C4X-N5-C5X	2.64	119.65	116.69
3	B	1217	FAD	C4-N3-C2	-2.60	120.05	125.39
3	A	1219	FAD	C4X-C10-N1	-2.57	120.16	122.73
3	B	1217	FAD	N7A-C8A-N9A	-2.53	107.22	114.36
3	B	1217	FAD	C5X-C9A-N10	2.51	119.28	116.80
3	A	1219	FAD	C4-N3-C2	-2.51	120.24	125.39
3	A	1219	FAD	C5A-C4A-N3A	-2.47	120.31	125.70
3	B	1217	FAD	C4A-C5A-N7A	-2.43	107.44	109.52
3	A	1219	FAD	C9A-N10-C10	-2.38	119.43	121.77
3	B	1217	FAD	C8A-N9A-C4A	2.37	108.71	106.90
3	A	1219	FAD	C2A-N3A-C4A	2.36	120.72	114.01
3	B	1217	FAD	O2P-P-O3P	2.32	116.16	105.14
3	B	1217	FAD	O3P-P-O5'	-2.30	93.10	103.41
3	B	1217	FAD	C1B-N9A-C4A	-2.25	122.74	126.64
3	A	1219	FAD	C5X-C9A-N10	2.24	119.01	116.80
3	A	1219	FAD	N7A-C8A-N9A	-2.22	108.08	114.36
3	B	1217	FAD	C2A-N3A-C4A	2.17	120.19	114.01
3	B	1217	FAD	C5A-C4A-N3A	-2.17	120.97	125.70
3	B	1217	FAD	O2P-P-O5'	2.12	119.17	108.51
3	B	1217	FAD	C9A-N10-C10	-2.07	119.73	121.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	180/217 (82%)	0.51	17 (9%) 9 14	18, 33, 84, 94	0
2	B	179/217 (82%)	0.51	19 (10%) 7 11	18, 33, 89, 92	1 (0%)
All	All	359/434 (82%)	0.51	36 (10%) 8 13	18, 33, 88, 94	1 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	PRO	7.1
1	A	31	ASN	6.7
2	B	145	LEU	5.5
1	A	89	GLN	5.4
1	A	124	THR	5.1
1	A	33	GLU	5.0
2	B	149	ILE	4.8
1	A	34	ASN	4.6
2	B	126	TRP	4.6
2	B	148	GLY	4.4
1	A	148	GLY	4.3
2	B	146	GLU	4.0
1	A	35	GLN	4.0
2	B	127	ALA	3.8
1	A	126	TRP	3.7
2	B	150	ALA	3.3
1	A	30	SER	3.2
2	B	143	LYS	3.2
2	B	125	TRP	3.1
1	A	1	MET	3.1
2	B	189	ARG	2.9
1	A	217	HIS	2.8
1	A	39	LYS	2.8
1	A	127	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	139	GLU	2.5
2	B	88	SER	2.5
1	A	125	TRP	2.4
1	A	156	PHE	2.3
2	B	134	TYR	2.3
2	B	135	ALA	2.2
2	B	1	MET	2.2
2	B	141	TYR	2.2
2	B	33	GLU	2.1
1	A	139	GLU	2.1
2	B	155	ARG	2.1
2	B	159	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CME	B	48	10/11	0.14	0.08	24,25,37,40	0
1	CME	A	48	10/11	0.11	-0.72	32,33,44,47	0

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	FAD	A	1219	53/53	0.14	-0.06	15,21,25,27	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FAD	B	1217	53/53	0.13	-0.41	16,25,32,32	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.