



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 08:54 PM GMT

PDB ID : 3CLS  
Title : Crystal structure of the R236C mutant of ETF from *Methylophilus methylotrophus*  
Authors : Katona, G.; Leys, D.  
Deposited on : 2008-03-20  
Resolution : 1.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	C	264	
2	D	321	

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5053 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Electron transfer flavoprotein subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	254	1950	1222	327	387	14	0	4	0


- Molecule 2 is a protein called Electron transfer flavoprotein subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	318	2334	1468	392	466	8	0	3	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	236	CYS	ARG	ENGINEERED	UNP P53571

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	307	Total	O	0	0
			307	307		
5	D	386	Total	O	0	0
			386	386		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Electron transfer flavoprotein subunit beta

Chain C: 



- Molecule 2: Electron transfer flavoprotein subunit alpha

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	117.53Å 117.53Å 84.85Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.65 19.68 – 1.65	Depositor EDS
% Data completeness (in resolution range)	95.6 (20.00-1.65) 95.6 (19.68-1.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 1.66Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.163 , 0.191 0.182 , 0.211	Depositor DCC
$R_{free}$ test set	3836 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 35.7	EDS
Estimated twinning fraction	0.030 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 76167 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5053	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	C	0.88	2/1974 (0.1%)	0.93	2/2671 (0.1%)
2	D	0.88	4/2372 (0.2%)	0.92	3/3232 (0.1%)
All	All	0.88	6/4346 (0.1%)	0.92	5/5903 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	214	GLU	CB-CG	7.54	1.66	1.52
2	D	214	GLU	CG-CD	7.05	1.62	1.51
1	C	125	GLN	CG-CD	6.30	1.65	1.51
1	C	162	ARG	CB-CG	-5.98	1.36	1.52
2	D	279	LYS	CD-CE	5.80	1.65	1.51
2	D	279	LYS	CE-NZ	5.28	1.62	1.49

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	279	LYS	CD-CE-NZ	7.36	128.62	111.70
1	C	242	MET	CG-SD-CE	-5.98	90.64	100.20
2	D	15	ARG	NE-CZ-NH2	5.50	123.05	120.30
2	D	100	LEU	CB-CG-CD2	5.44	120.24	111.00
1	C	117	PHE	CB-CG-CD2	-5.14	117.20	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1950	0	12	11	0
2	D	2334	0	0	8	0
3	C	23	0	12	0	0
4	D	53	0	31	1	0
5	C	307	0	0	7	0
5	D	386	0	0	5	0
All	All	5053	0	55	18	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (18) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:137[A]:GLN:NE2	5:D:1493:HOH:O	2.04	0.91
1:C:24:MET:CE	5:C:1560:HOH:O	2.34	0.74
1:C:261:PHE:O	1:C:262:LYS:HB2	1.86	0.73
2:D:252:GLN:NE2	5:D:1701:HOH:O	2.26	0.67
2:D:87[B]:ASN:OD1	5:D:1442:HOH:O	2.15	0.65
2:D:44:GLN:NE2	5:D:1392:HOH:O	2.33	0.60
1:C:98:ILE:CG1	1:C:226[B]:MET:CE	2.80	0.60
1:C:226[A]:MET:CE	2:D:112:TYR:O	2.51	0.57
1:C:240:ALA:CB	1:C:242:MET:CE	2.84	0.56
1:C:102:ARG:CZ	5:C:1566:HOH:O	2.56	0.53
1:C:102:ARG:NE	5:C:1566:HOH:O	2.45	0.49
2:D:186:LYS:NZ	5:D:1698:HOH:O	2.47	0.47
2:D:236:CYS:SG	4:D:1319:FAD:H1'2	2.55	0.47
1:C:102:ARG:CG	5:C:1562:HOH:O	2.64	0.46
1:C:102:ARG:NH2	5:C:1566:HOH:O	2.49	0.44
1:C:32:MET:CG	5:C:1525:HOH:O	2.66	0.44
1:C:202:THR:N	5:C:1271:HOH:O	2.53	0.42
2:D:163:LEU:N	2:D:163:LEU:CD2	2.83	0.41

There are no symmetry-related clashes.



## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	254/264 (96%)	249 (98%)	4 (2%)	1 (0%)	43	19
2	D	319/321 (99%)	312 (98%)	7 (2%)	0	100	100
All	All	573/585 (98%)	561 (98%)	11 (2%)	1 (0%)	56	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	145	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	204/216 (94%)	198 (97%)	6 (3%)	55	22
2	D	252/259 (97%)	248 (98%)	4 (2%)	75	51
All	All	456/475 (96%)	446 (98%)	10 (2%)	64	35

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	55	THR
1	C	102	ARG
1	C	156	ASN
1	C	162	ARG
1	C	202	THR
1	C	262	LYS
2	D	19	LEU
2	D	60	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	236	CYS
2	D	279	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AMP	C	1262	-	25,25,25	1.50	6 (24%)	38,38,38	2.18	11 (28%)
4	FAD	D	1319	-	58,58,58	1.27	5 (8%)	85,89,89	1.69	9 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AMP	C	1262	-	-	0/10/26/26	0/1/3/3
4	FAD	D	1319	-	-	0/34/50/50	0/1/6/6

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	1319	FAD	C2B-C1B	-4.17	1.47	1.53
4	D	1319	FAD	C1'-N10	3.67	1.52	1.48
3	C	1262	AMP	P-O3P	-3.01	1.43	1.54
3	C	1262	AMP	C4-N9	-3.00	1.33	1.37
4	D	1319	FAD	C5X-N5	2.85	1.39	1.35
3	C	1262	AMP	O4'-C1'	-2.65	1.37	1.41
4	D	1319	FAD	C2A-N3A	2.56	1.37	1.32
3	C	1262	AMP	C5-C4	2.53	1.46	1.40
4	D	1319	FAD	C2A-N1A	2.20	1.38	1.33
3	C	1262	AMP	C2-N1	2.19	1.38	1.33
3	C	1262	AMP	C5-N7	-2.18	1.32	1.40

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1319	FAD	N3A-C2A-N1A	-10.50	119.93	128.71
3	C	1262	AMP	N3-C2-N1	-7.13	122.75	128.71
3	C	1262	AMP	N3-C4-N9	5.23	134.88	125.43
3	C	1262	AMP	C8-N9-C4	4.60	110.41	106.90
4	D	1319	FAD	C2-N1-C10	3.99	119.00	114.98
3	C	1262	AMP	C1'-N9-C4	-3.35	120.84	126.64
4	D	1319	FAD	C4B-O4B-C1B	-3.07	106.42	109.75
4	D	1319	FAD	N3A-C4A-N9A	2.97	130.79	125.43
3	C	1262	AMP	N6-C6-N1	2.84	124.94	119.36
4	D	1319	FAD	C4A-C5A-N7A	-2.83	107.10	109.52
3	C	1262	AMP	C5-C4-N9	-2.59	103.43	107.16
4	D	1319	FAD	C2A-N3A-C4A	2.38	120.79	114.01
3	C	1262	AMP	P-O5'-C5'	2.36	125.01	118.19
4	D	1319	FAD	O4'-C4'-C3'	2.34	114.86	109.05
3	C	1262	AMP	C5'-C4'-C3'	-2.33	105.89	115.21
4	D	1319	FAD	O4B-C1B-N9A	-2.27	106.33	108.44
3	C	1262	AMP	O3P-P-O1P	2.25	117.78	110.44
3	C	1262	AMP	O5'-P-O1P	-2.17	100.34	106.71
3	C	1262	AMP	C5-C6-N6	-2.11	115.95	120.72
4	D	1319	FAD	C5A-C4A-N3A	-2.09	121.15	125.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	C	254/264 (96%)	-0.13	9 (3%) 42 42	0, 10, 30, 48	0
2	D	318/321 (99%)	-0.14	10 (3%) 47 48	2, 12, 33, 48	0
All	All	572/585 (97%)	-0.13	19 (3%) 44 44	0, 11, 31, 48	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	192	GLY	5.7
2	D	32	GLY	5.3
1	C	191	TYR	4.5
1	C	193	SER	4.4
1	C	202	THR	4.1
2	D	191	VAL	3.6
1	C	21	GLU	3.6
2	D	179	VAL	3.2
2	D	226	ASP	3.1
2	D	31	SER	3.0
1	C	55	THR	2.6
2	D	197	ILE	2.4
2	D	147	GLY	2.4
1	C	54	ASP	2.3
2	D	111	GLY	2.3
1	C	192	ALA	2.1
1	C	156	ASN	2.0
2	D	228	ALA	2.0
1	C	16	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FAD	D	1319	53/53	0.07	0.04	3,5,9,10	0
3	AMP	C	1262	23/23	0.06	-0.48	2,3,4,6	0

### 6.5 Other polymers

There are no such residues in this entry.