



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 12:02 AM GMT

PDB ID : 1CMX
Title : STRUCTURAL BASIS FOR THE SPECIFICITY OF UBIQUITIN C-
TERMINAL HYDROLASES
Authors : Johnston, S.C.; Riddle, S.M.; Cohen, R.E.; Hill, C.P.
Deposited on : 1999-05-12
Resolution : 2.25 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	235	
1	C	235	
2	B	76	
2	D	76	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4142 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (UBIQUITIN YUH1-UBAL).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	10	0	0
			1684	1072	274	335	3			
1	C	210	Total	C	N	O	S	24	0	0
			1649	1052	266	328	3			

- Molecule 2 is a protein called PROTEIN (UBIQUITIN YUH1-UBAL).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	76	Total	C	N	O	S	10	0	0
			601	378	105	117	1			
2	D	17	Total	C	N	O		6	0	0
			133	83	29	21				

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	38	Total	O	0	0
			38	38		
3	B	16	Total	O	0	0
			16	16		
3	C	18	Total	O	0	0
			18	18		
3	D	3	Total	O	0	0
			3	3		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN (UBIQUITIN YUH1-UBAL)

Chain A: 



- Molecule 1: PROTEIN (UBIQUITIN YUH1-UBAL)

Chain C: 




- Molecule 2: PROTEIN (UBIQUITIN YUH1-UBAL)

Chain B: 



- Molecule 2: PROTEIN (UBIQUITIN YUH1-UBAL)

Chain D: 



GLU	
SER	
THR	
LEU	
HIS	
L769	
	G776

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	199.30Å 199.30Å 36.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.25 28.77 – 2.23	Depositor EDS
% Data completeness (in resolution range)	99.0 (30.00-2.25) 97.7 (28.77-2.23)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.22Å)	Xtriage
Refinement program	X-PLOR 98.0	Depositor
R, R_{free}	0.248 , 0.285 0.243 , 0.281	Depositor DCC
R_{free} test set	1237 reflections (4.83%)	DCC
Wilson B-factor (Å ²)	46.1	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.4	EDS
Estimated twinning fraction	0.022 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 26106 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4142	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/1719	0.86	0/2340
1	C	0.63	0/1684	0.83	1/2293 (0.0%)
2	B	0.85	0/603	0.89	0/811
2	D	0.65	0/126	0.89	0/163
All	All	0.70	0/4132	0.86	1/5607 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	440	THR	N-CA-CB	6.76	123.14	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	0	1634	53	0
1	C	1649	0	1600	39	0
2	B	601	0	625	15	0
2	D	133	0	151	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	38	0	0	0	0
3	B	16	0	0	2	0
3	C	18	0	0	1	0
3	D	3	0	0	0	0
All	All	4142	0	4010	99	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (99) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:439:LEU:O	1:C:440:THR:HG23	1.65	0.95
1:A:41:GLU:HG2	1:A:44:LEU:HD12	1.62	0.80
1:A:83:LYS:NZ	1:A:185:LEU:HD22	1.98	0.79
1:A:48:LEU:HD21	1:A:202:ILE:CD1	2.18	0.73
1:A:40:THR:HG23	1:C:440:THR:HB	1.71	0.73
1:C:458:LEU:HD11	1:C:565:LEU:HB3	1.71	0.72
1:A:42:PRO:HG2	1:A:43:GLU:H	1.56	0.71
1:C:529:PHE:HB3	1:C:534:THR:HG21	1.73	0.70
1:C:538:VAL:O	1:C:542:ILE:HG13	1.91	0.69
1:A:224:LEU:HD21	2:B:311:LYS:HD3	1.76	0.67
1:C:515:PHE:HZ	1:C:534:THR:HG22	1.61	0.67
2:B:361:ILE:HD13	2:B:367:LEU:HD21	1.79	0.63
1:A:83:LYS:HZ1	1:A:185:LEU:HD22	1.63	0.62
1:A:83:LYS:HZ3	1:A:185:LEU:HD22	1.66	0.60
1:A:48:LEU:HD21	1:A:202:ILE:HD13	1.83	0.59
1:A:79:VAL:HG12	1:A:173:GLU:HG3	1.85	0.59
1:C:428:LYS:HB3	1:C:430:GLU:OE1	2.03	0.58
1:A:39:LEU:HD22	1:A:228:MET:HE1	1.84	0.58
1:C:439:LEU:O	1:C:440:THR:CG2	2.48	0.58
1:A:10:PRO:HD3	1:A:88:ASN:ND2	2.19	0.58
1:A:39:LEU:HD22	1:A:228:MET:CE	2.34	0.57
1:A:40:THR:O	1:A:40:THR:HG22	2.05	0.57
1:A:37:TYR:HE1	1:A:44:LEU:HD22	1.69	0.56
1:C:439:LEU:HD11	1:C:628:MET:CE	2.35	0.56
1:A:31:TRP:CE2	1:A:105:LEU:HD22	2.41	0.56
1:A:8:VAL:CG1	1:A:149:PHE:CD2	2.89	0.56
2:B:323:ILE:HB	2:B:352:ASP:HA	1.89	0.55
1:A:10:PRO:HD3	1:A:88:ASN:HD22	1.71	0.54
1:A:40:THR:HG21	1:C:440:THR:HG22	1.88	0.54
1:C:434:PHE:CD1	1:C:448:LEU:HD22	2.43	0.54
1:A:8:VAL:HG11	1:A:149:PHE:CE2	2.43	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:434:PHE:HZ	1:C:452:VAL:HG22	1.72	0.53
1:A:24:LYS:HD3	1:A:145:ASN:OD1	2.08	0.53
1:A:40:THR:HG22	1:C:608:ARG:NH1	2.23	0.53
1:C:458:LEU:HD12	1:C:566:HIS:O	2.08	0.53
2:B:325:ASN:ND2	3:B:871:HOH:O	2.34	0.53
1:C:408:VAL:HG11	1:C:549:PHE:CD2	2.44	0.52
1:C:408:VAL:HG12	1:C:409:VAL:N	2.25	0.52
2:B:345:PHE:HB3	2:B:350:LEU:HD21	1.92	0.52
1:A:177:ILE:HD11	1:A:204:GLN:OE1	2.09	0.52
1:C:441:GLU:CD	1:C:442:PRO:HD2	2.31	0.51
1:A:8:VAL:HG11	1:A:149:PHE:CD2	2.45	0.51
1:C:519:GLN:OE1	1:C:529:PHE:CE1	2.64	0.51
1:A:36:ILE:HG21	1:A:228:MET:HE2	1.92	0.51
1:A:218:ALA:HB2	1:A:226:PHE:CE1	2.46	0.51
1:A:170:TYR:CE2	1:A:179:GLU:HG3	2.47	0.50
1:C:408:VAL:HG12	1:C:409:VAL:H	1.77	0.50
1:C:537:PHE:O	1:C:541:VAL:HG23	2.10	0.50
1:A:44:LEU:HD11	2:B:368:HIS:NE2	2.27	0.49
1:A:83:LYS:HZ3	1:A:185:LEU:CD2	2.25	0.49
1:C:540:ASN:O	1:C:544:GLU:HG3	2.11	0.49
2:B:322:THR:HA	2:B:355:THR:HA	1.93	0.49
1:C:450:ARG:NE	1:C:600:ASP:OD2	2.44	0.49
1:A:38:SER:HB3	1:A:41:GLU:HB2	1.95	0.49
1:A:38:SER:HB3	1:A:41:GLU:CB	2.42	0.49
1:A:119:GLN:NE2	1:A:133:THR:HG21	2.28	0.48
1:C:480:ILE:HD12	1:C:500:SER:O	2.13	0.48
1:C:439:LEU:HD11	1:C:628:MET:HE3	1.96	0.47
2:B:301:MET:HE1	3:B:838:HOH:O	2.13	0.47
1:A:141:VAL:O	1:A:145:ASN:ND2	2.37	0.47
2:B:339:ASP:OD1	2:B:340:GLN:HG3	2.15	0.47
1:C:431:TRP:CE2	1:C:505:LEU:HD22	2.50	0.47
1:A:23:HIS:HD2	1:A:27:LEU:O	1.96	0.47
1:A:139:LEU:O	1:A:143:LYS:HG3	2.15	0.47
1:C:425:LEU:HD21	1:C:541:VAL:HG11	1.97	0.46
1:A:215:MET:O	1:A:218:ALA:HB3	2.14	0.46
1:A:52:VAL:HG21	1:A:202:ILE:HD12	1.98	0.46
1:C:519:GLN:OE1	1:C:529:PHE:CD1	2.69	0.45
1:A:48:LEU:HA	1:A:49:PRO:HD3	1.73	0.45
1:A:79:VAL:O	1:A:127:ASN:HB2	2.16	0.45
1:C:448:LEU:HA	1:C:449:PRO:HD3	1.83	0.45
1:C:549:PHE:HB2	3:C:824:HOH:O	2.17	0.45
1:A:42:PRO:CG	1:A:43:GLU:H	2.29	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:119:GLN:HE22	1:A:133:THR:HG21	1.80	0.44
2:B:356:LEU:HD22	2:B:361:ILE:HD12	1.99	0.44
1:C:458:LEU:HA	1:C:566:HIS:O	2.17	0.44
2:B:333:LYS:HB3	2:B:333:LYS:HE2	1.69	0.44
1:C:529:PHE:CB	1:C:534:THR:HG21	2.46	0.43
1:C:451:PRO:CG	1:C:453:LYS:HE2	2.48	0.43
1:A:177:ILE:HD11	1:A:204:GLN:CD	2.39	0.43
1:A:39:LEU:HD22	1:A:228:MET:SD	2.59	0.43
1:A:37:TYR:CE1	1:A:44:LEU:HD22	2.52	0.42
1:A:156:ALA:HA	1:A:157:PRO:HD3	1.85	0.42
1:A:137:PHE:O	1:A:141:VAL:HG23	2.20	0.42
2:B:304:PHE:HA	2:B:313:ILE:O	2.20	0.42
1:A:87:LYS:NZ	1:A:160:THR:O	2.50	0.42
1:A:233:PRO:O	1:A:234:ASN:HB3	2.20	0.42
2:B:306:LYS:HZ1	1:C:605:GLU:HG3	1.85	0.42
1:C:460:PRO:HG2	1:C:623:VAL:HG11	2.02	0.41
1:C:515:PHE:CZ	1:C:534:THR:HG22	2.48	0.41
1:A:41:GLU:HG2	1:A:44:LEU:CD1	2.43	0.41
1:A:83:LYS:NZ	1:A:185:LEU:CD2	2.76	0.41
2:B:342:ARG:HD3	2:B:349:GLN:OE1	2.21	0.41
1:A:195:ASP:HA	1:A:196:PRO:HD2	1.72	0.41
1:A:40:THR:CG2	1:C:608:ARG:CZ	2.99	0.41
2:B:323:ILE:HG22	2:B:327:LYS:HE3	2.01	0.41
1:A:40:THR:OG1	1:C:440:THR:HG21	2.20	0.40
1:C:408:VAL:HG11	1:C:549:PHE:CE2	2.56	0.40
1:C:484:GLN:HA	1:C:493:TYR:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/235 (89%)	191 (91%)	14 (7%)	5 (2%)	9	4
1	C	206/235 (88%)	193 (94%)	9 (4%)	4 (2%)	12	6

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	74/76 (97%)	72 (97%)	2 (3%)	0	100	100
2	D	11/76 (14%)	9 (82%)	2 (18%)	0	100	100
All	All	501/622 (80%)	465 (93%)	27 (5%)	9 (2%)	13	6

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	THR
1	C	440	THR
1	C	524	SER
1	C	527	ASN
1	A	61	ILE
1	A	124	SER
1	A	127	ASN
1	A	42	PRO
1	C	442	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/209 (90%)	177 (94%)	12 (6%)	25	24
1	C	185/209 (88%)	174 (94%)	11 (6%)	28	27
2	B	68/68 (100%)	65 (96%)	3 (4%)	39	43
2	D	14/68 (21%)	14 (100%)	0	100	100
All	All	456/554 (82%)	430 (94%)	26 (6%)	29	29

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ASN
1	A	38	SER
1	A	61	ILE
1	A	79	VAL
1	A	83	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	126	LYS
1	A	130	ASP
1	A	131	ASP
1	A	136	GLN
1	A	146	VAL
1	A	163	THR
1	A	197	THR
2	B	342	ARG
2	B	354	ARG
2	B	363	LYS
1	C	438	SER
1	C	483	LYS
1	C	526	LYS
1	C	531	ASP
1	C	536	GLN
1	C	540	ASN
1	C	543	LYS
1	C	583	ARG
1	C	585	LEU
1	C	586	SER
1	C	594	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
2	B	340	GLN
1	C	420	ASN
1	C	519	GLN
2	D	740	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLZ	B	376	1,2	3,3,3	0.83	0	1,2,2	1.83	0
2	GLZ	D	776	1,2	3,3,3	0.70	0	1,2,2	1.18	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLZ	B	376	1,2	-	0/0/1/1	0/0/0/0
2	GLZ	D	776	1,2	-	0/0/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/235 (91%)	0.36	19 (8%) 10 12	23, 48, 92, 100	2 (0%)
1	C	210/235 (89%)	1.09	45 (21%) 1 1	25, 58, 100, 100	5 (2%)
2	B	76/76 (100%)	-0.34	0 100 100	25, 38, 57, 63	2 (2%)
2	D	17/76 (22%)	0.74	2 (11%) 5 6	47, 62, 87, 92	1 (5%)
All	All	517/622 (83%)	0.57	66 (12%) 4 5	23, 50, 98, 100	10 (1%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	522	THR	10.0
1	C	552	GLY	9.3
1	C	521	ASP	8.9
1	C	524	SER	8.3
1	C	560	THR	7.5
1	C	523	SER	7.2
1	C	618	ALA	6.4
1	C	525	SER	6.2
1	C	443	GLU	6.1
1	A	123	SER	5.6
1	A	39	LEU	5.1
1	C	407	ALA	5.0
1	C	440	THR	5.0
1	C	527	ASN	4.6
1	C	408	VAL	4.4
1	C	446	ALA	4.4
1	C	442	PRO	4.3
1	C	551	THR	4.2
1	C	553	GLN	4.2
1	A	40	THR	4.1
1	C	439	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	585	LEU	3.7
1	A	124	SER	3.7
1	C	597	THR	3.6
1	A	43	GLU	3.5
1	C	526	LYS	3.5
1	C	563	THR	3.4
1	A	56	VAL	3.3
1	C	532	VAL	3.2
1	C	564	ASN	3.2
1	A	78	ASP	3.2
1	C	619	ASN	3.1
1	C	559	ALA	3.0
1	C	586	SER	3.0
1	C	622	ASP	2.9
1	A	155	GLU	2.9
1	A	199	THR	2.8
2	D	743	LEU	2.7
1	A	95	ILE	2.7
1	C	456	VAL	2.6
1	C	447	PHE	2.6
1	C	495	ILE	2.5
1	A	197	THR	2.5
1	C	496	LEU	2.5
1	A	94	ALA	2.5
1	C	494	ALA	2.5
1	C	555	GLU	2.5
1	A	195	ASP	2.4
1	A	190	TYR	2.4
1	C	561	ALA	2.3
1	A	125	SER	2.3
1	A	169	THR	2.3
2	D	706	LYS	2.3
1	A	208	ARG	2.3
1	C	529	PHE	2.3
1	C	558	GLU	2.3
1	A	55	ILE	2.2
1	C	457	LEU	2.2
1	C	621	GLU	2.1
1	C	444	LEU	2.1
1	A	229	LEU	2.1
1	C	508	PRO	2.1
1	C	530	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	599	THR	2.0
1	C	627	ALA	2.0
1	C	617	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GLZ	D	776	4/4	0.12	-0.92	57,61,62,66	0
2	GLZ	B	376	4/4	0.08	-1.37	26,32,32,32	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.