



# Full wwPDB X-ray Structure Validation Report i

Aug 15, 2014 – 04:31 AM EDT

PDB ID : 4CRI  
Title : Crystal Structure of 53BP1 tandem tudor domains in complex with methylated K810 Rb peptide  
Authors : Krojer, T.; Johansson, C.; Gileadi, C.; Fedorov, O.; Carr, S.; La Thangue, N.B.; Vollmar, M.; Crawley, L.; von Delft, F.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Oppermann, U.  
Deposited on : 2014-02-26  
Resolution : 2.35 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable23489  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable23489

# 1 Overall quality at a glance

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	176	
1	B	176	
2	C	17	
2	D	17	

## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1991 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TUMOR SUPPRESSOR P53-BINDING PROTEIN 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	120	Total 944	C 605	N 157	O 179	S 3	0	0	1
1	B	120	Total 922	C 589	N 152	O 178	S 3	0	0	1

- Molecule 2 is a protein called RB1 PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	7	Total 60	C 43	N 8	O 9	0	0	0
2	D	6	Total 42	C 28	N 7	O 7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	818	CYS	-	EXPRESSION TAG	UNP P78495
D	818	CYS	-	EXPRESSION TAG	UNP P78495

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total 17	O 17	0	0
3	B	6	Total 6	O 6	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TUMOR SUPPRESSOR P53-BINDING PROTEIN 1

Chain A:



- Molecule 1: TUMOR SUPPRESSOR P53-BINDING PROTEIN 1

Chain B:



- Molecule 2: RB1 PROTEIN

Chain C:



- Molecule 2: RB1 PROTEIN

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.79Å 105.79Å 156.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.51 – 2.35 39.51 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.51-2.35) 99.9 (39.51-2.35)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 2.34Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.194 , 0.230 0.194 , 0.230	Depositor DCC
$R_{free}$ test set	1159 reflections (5.24%)	DCC
Wilson B-factor (Å <sup>2</sup> )	59.6	Xtrriage
Anisotropy	0.211	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.4	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 22130 reflections	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1991	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	1/963 (0.1%)	0.65	1/1295 (0.1%)
1	B	0.47	1/941 (0.1%)	0.67	0/1272
2	C	0.57	0/50	0.73	0/68
2	D	0.35	0/31	0.64	0/42
All	All	0.50	2/1985 (0.1%)	0.66	1/2677 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1603	GLY	C-N	-5.55	1.23	1.34
1	A	1603	GLY	C-N	-5.16	1.24	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1517	LEU	CB-CG-CD2	-5.88	101.00	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	944	0	925	8	0
1	B	922	0	871	14	0
2	C	60	0	68	4	0
2	D	42	0	41	4	0
3	A	17	0	0	1	1
3	B	6	0	0	0	0
All	All	1991	0	1905	23	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (23) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:1594:ASN:OD1	1:B:1597:ARG:NH2	2.15	0.79
1:B:1573:GLU:HB2	1:B:1578:ARG:HG2	1.64	0.77
1:B:1549:GLU:HA	2:D:805:TYR:HA	1.65	0.76
1:A:1573:GLU:HB2	1:A:1578:ARG:HG2	1.78	0.65
1:B:1552:TYR:HA	2:D:806:ILE:HG12	1.86	0.58
1:B:1548:SER:H	2:D:806:ILE:HG22	1.70	0.57
1:A:1595:ARG:NH1	3:A:2017:HOH:O	2.33	0.56
1:B:1492:VAL:HG12	1:B:1501:PHE:HB3	1.88	0.54
1:A:1521:ASP:OD2	2:C:810:MLY:HH22	2.08	0.53
1:B:1534:LEU:HD22	1:B:1602:LEU:HD11	1.93	0.51
1:B:1514:LYS:NZ	2:C:810:MLY:O	2.44	0.50
1:A:1507:THR:O	1:A:1508:ARG:HG2	2.11	0.50
1:B:1554:SER:OG	1:B:1555:ALA:N	2.49	0.45
1:B:1548:SER:HB3	1:B:1551:GLU:O	2.19	0.43
2:C:807:SER:HB2	2:C:808:PRO:HD2	2.01	0.43
1:A:1495:TRP:CD1	2:C:810:MLY:HD2	2.54	0.43
1:B:1500:TYR:CD2	1:B:1589:SER:HA	2.55	0.42
1:A:1492:VAL:HG12	1:A:1501:PHE:HB3	2.02	0.42
1:B:1517:LEU:HD13	1:B:1527:VAL:HB	2.02	0.42
1:B:1551:GLU:O	2:D:806:ILE:HG21	2.20	0.41
1:A:1534:LEU:HD22	1:A:1602:LEU:HD11	2.02	0.41
1:A:1534:LEU:HB2	1:A:1602:LEU:CD1	2.50	0.41
1:B:1501:PHE:HB2	1:B:1588:LEU:HB2	2.01	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:A:2015:HOH:O	3:A:2015:HOH:O[12_565]	1.63	0.57

## 5.3 Torsion angles

### 5.3.1 Protein backbone i

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	118/176 (67%)	115 (98%)	2 (2%)	1 (1%)	27	31
1	B	118/176 (67%)	114 (97%)	4 (3%)	0	100	100
2	C	5/17 (29%)	5 (100%)	0	0	100	100
2	D	4/17 (24%)	4 (100%)	0	0	100	100
All	All	245/386 (64%)	238 (97%)	6 (2%)	1 (0%)	43	52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1603	GLY

### 5.3.2 Protein sidechains i

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/146 (67%)	98 (100%)	0	100	100
1	B	93/146 (64%)	90 (97%)	3 (3%)	51	67
2	C	6/15 (40%)	5 (83%)	1 (17%)	3	2
2	D	3/15 (20%)	3 (100%)	0	100	100
All	All	200/322 (62%)	196 (98%)	4 (2%)	68	83

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1496	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	1530	LYS
1	B	1535	CYS
2	C	804	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	MLY	C	810	2	10,10,11	5.95	2 (20%)	9,11,13	2.57	1 (11%)
2	MLY	D	810	2	10,10,11	5.56	1 (10%)	9,11,13	2.21	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLY	C	810	2	-	0/7/9/11	0/0/0/0
2	MLY	D	810	2	-	0/7/9/11	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	810	MLY	O-C	18.59	1.24	1.11

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	810	MLY	O-C	17.42	1.23	1.11
2	C	810	MLY	CA-C	2.51	1.54	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	810	MLY	C-CA-N	-7.38	106.45	113.83
2	D	810	MLY	C-CA-N	-6.03	107.80	113.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	120/176 (68%)	0.34	5 (4%) 35 38	42, 66, 98, 122	0
1	B	120/176 (68%)	0.43	8 (6%) 17 20	52, 77, 126, 145	0
2	C	7/17 (41%)	1.10	0 100 100	47, 51, 63, 81	0
2	D	6/17 (35%)	0.41	0 100 100	79, 96, 115, 130	0
All	All	253/386 (65%)	0.41	13 (5%) 27 30	42, 71, 115, 145	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1556	GLY	4.6
1	B	1552	TYR	4.0
1	B	1604	PRO	3.5
1	A	1510	VAL	3.2
1	A	1604	PRO	3.2
1	A	1534	LEU	3.1
1	B	1485	SER	3.0
1	A	1486	PHE	2.9
1	A	1485	SER	2.9
1	B	1576	GLY	2.7
1	B	1559	LYS	2.3
1	B	1555	ALA	2.3
1	B	1575	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MLY	C	810	11/12	0.25	1.87	49,55,64,66	0
2	MLY	D	810	11/12	0.19	1.17	65,79,85,85	0

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

There are no ligands in this entry.

### 6.5 Other polymers

There are no such residues in this entry.