



Full wwPDB X-ray Structure Validation Report (i)

Mar 1, 2014 – 03:12 AM GMT

PDB ID : 1CV2

Title : Hydrolytic haloalkane dehalogenase libn from sphingomonas paucimobilis UT26 AT 1.6 Å resolution

Authors : Marek, J.; Vevodova, J.; Damborsky, J.; Smatanova, I.; Svensson, L.A.; Newman, J.; Nagata, Y.; Takagi, M.

Deposited on : 1999-08-22

Resolution : 1.58 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.15 2013

Xtriage (Phenix) : dev-1323

EDS : stable22639

Percentile statistics : 21963

Refmac : 5.8.0049

CCP4 : 6.3.0 (Settle)

Ideal geometry (proteins) : Engh & Huber (2001)


Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)

Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance (i)

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2778 (1.60-1.56)
Clashscore	79885	3207 (1.60-1.56)
Ramachandran outliers	78287	3107 (1.60-1.56)
Sidechain outliers	78261	3104 (1.60-1.56)
RSRZ outliers	66119	2778 (1.60-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	296	<div style="width: 100%;"><div style="width: 100%; background-color: green;"></div></div>

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2750 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HALOALKANE DEHALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	293	2301	1466	409	416	10	0	0	0

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	449	449	449	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HALOALKANE DEHALOGENASE

Chain A:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	50.26 Å 71.67 Å 72.70 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.58 23.75 – 1.58	Depositor EDS
% Data completeness (in resolution range)	94.2 (20.00-1.58) 94.2 (23.75-1.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	5.46 (at 1.58 Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R , R_{free}	0.152 , 0.211 0.149 , 0.201	Depositor DCC
R_{free} test set	1745 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	8.6	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 70.4	EDS
Estimated twinning fraction	0.015 for -h,l,k	Xtriage
L-test for twinning	$< L > = 0.48$, $< L^2 > = 0.31$	Xtriage
Outliers	0 of 34649 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2750	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.48	0/2364	1.27	20/3214 (0.6%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	295	PRO	C-N-CA	16.51	162.99	121.70
1	A	40	THR	N-CA-CB	-11.47	88.52	110.30
1	A	202	ARG	NE-CZ-NH2	-7.52	116.54	120.30
1	A	20	ARG	NE-CZ-NH1	7.12	123.86	120.30
1	A	292	ARG	CD-NE-CZ	6.88	133.23	123.60
1	A	292	ARG	NE-CZ-NH2	-6.77	116.91	120.30
1	A	44	LEU	CA-CB-CG	6.68	130.66	115.30
1	A	294	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	A	19	ARG	CA-C-N	6.07	130.56	117.20
1	A	80	TYR	CB-CG-CD1	5.92	124.55	121.00
1	A	108	ASP	CB-CG-OD1	5.86	123.57	118.30
1	A	240	PHE	CB-CG-CD1	5.84	124.89	120.80
1	A	202	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	A	201	ARG	NE-CZ-NH1	-5.63	117.48	120.30
1	A	254	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	70	ASP	CB-CG-OD1	5.47	123.22	118.30
1	A	227	TYR	CA-CB-CG	5.46	123.77	113.40
1	A	40	THR	CB-CA-C	5.23	125.72	111.60
1	A	191	ARG	CA-CB-CG	5.06	124.54	113.40
1	A	258	ARG	NE-CZ-NH1	-5.02	117.79	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2234	41	0
2	A	449	0	0	17	0
All	All	2750	0	2234	41	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

All (41) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:242:ASN:HD21	1:A:254:ARG:HD2	1.40	0.85
1:A:267:THR:HG23	2:A:1124:HOH:O	1.86	0.76
1:A:242:ASN:HB3	1:A:267:THR:HG22	1.72	0.70
1:A:36:HIS:HE1	1:A:63:LEU:H	1.39	0.68
1:A:6:LYS:HD3	1:A:7:PRO:O	1.93	0.68
1:A:72:LEU:HD13	1:A:79:ARG:HD2	1.76	0.67
1:A:20:ARG:HD2	2:A:1377:HOH:O	1.96	0.66
1:A:40:THR:HG21	2:A:1016:HOH:O	1.97	0.65
1:A:36:HIS:CE1	1:A:63:LEU:H	2.15	0.63
1:A:146:GLN:HB3	2:A:1299:HOH:O	2.02	0.58
1:A:256:PHE:O	1:A:259:THR:HG22	2.03	0.58
1:A:36:HIS:HD2	1:A:37:GLY:O	1.88	0.56
1:A:266:ILE:HG13	2:A:1153:HOH:O	2.08	0.54
1:A:19:ARG:O	1:A:70:ASP:HB2	2.08	0.53
1:A:36:HIS:HB2	1:A:40:THR:CG2	2.38	0.53
1:A:242:ASN:ND2	1:A:254:ARG:HD2	2.17	0.52
1:A:146:GLN:HG2	2:A:1211:HOH:O	2.08	0.52
1:A:51:HIS:HD2	2:A:1277:HOH:O	1.91	0.51
1:A:85:HIS:HE1	2:A:1010:HOH:O	1.95	0.50
1:A:85:HIS:HD2	2:A:1011:HOH:O	1.93	0.50
1:A:146:GLN:HG2	2:A:1194:HOH:O	2.13	0.49
1:A:78:GLU:OE2	1:A:78:GLU:HA	2.13	0.48
1:A:174:LEU:HB3	1:A:175:PRO:HD3	1.96	0.48
1:A:216:THR:HB	2:A:1164:HOH:O	2.13	0.48
1:A:72:LEU:CD1	1:A:79:ARG:HD2	2.42	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:44:LEU:HD22	1:A:275:GLN:HB2	1.96	0.46
1:A:157:GLN:NE2	2:A:1324:HOH:O	2.50	0.45
1:A:19:ARG:O	1:A:70:ASP:OD2	2.34	0.45
1:A:51:HIS:HE1	1:A:280:ASP:OD1	2.00	0.44
1:A:4:GLY:N	2:A:1352:HOH:O	2.50	0.44
1:A:20:ARG:NH1	2:A:1408:HOH:O	2.50	0.44
1:A:267:THR:HG22	2:A:1171:HOH:O	2.18	0.44
1:A:270:GLY:HA3	1:A:274:ILE:HG22	2.01	0.42
1:A:157:GLN:HG2	2:A:1315:HOH:O	2.20	0.42
1:A:259:THR:HG21	2:A:1052:HOH:O	2.19	0.42
1:A:106:VAL:HB	1:A:111:SER:HA	2.02	0.42
1:A:242:ASN:ND2	1:A:258:ARG:NH2	2.68	0.41
1:A:17:LYS:HE2	1:A:87:ASP:HB3	2.01	0.41
1:A:178:ILE:CG1	1:A:273:PHE:HB2	2.50	0.41
1:A:260:TRP:HA	1:A:261:PRO:HD3	1.93	0.40
1:A:75:SER:HB3	1:A:199:GLU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	291/296 (98%)	281 (97%)	9 (3%)	1 (0%)	50 23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	20	ARG

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	230/236 (98%)	220 (96%)	10 (4%)	40 11

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	LYS
1	A	39	PRO
1	A	40	THR
1	A	44	LEU
1	A	146	GLN
1	A	169	PHE
1	A	191	ARG
1	A	207	TRP
1	A	225	ARG
1	A	259	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	HIS
1	A	51	HIS
1	A	85	HIS
1	A	146	GLN

5.3.3 RNA (i)

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/296 (98%)	-0.39	1 (0%) 91 94	5, 9, 25, 43	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	296	ALA	4.5

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.