



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 26, 2014 – 02:59 PM GMT

PDB ID : 2CVZ  
Title : Structure of hydroxyisobutyrate dehydrogenase from thermus thermophilus HB8  
Authors : Lokanath, N.K.; Kunishima, N.; Miyano, M.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)  
Deposited on : 2005-06-14  
Resolution : 1.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3513 (1.80-1.80)
Clashscore	79885	4461 (1.80-1.80)
Ramachandran outliers	78287	4404 (1.80-1.80)
Sidechain outliers	78261	4403 (1.80-1.80)
RSRZ outliers	66119	3515 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	289	
1	B	289	
1	C	289	
1	D	289	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9812 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-hydroxyisobutyrate dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	Se	0	0	0
			2201	1399	393	403	1	5			
1	B	288	Total	C	N	O	S	Se	0	0	0
			2201	1399	393	403	1	5			
1	C	289	Total	C	N	O	S	Se	0	0	0
			2209	1404	394	404	1	6			
1	D	288	Total	C	N	O	S	Se	0	0	0
			2201	1399	393	403	1	5			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
A	12	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
A	16	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
A	131	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
A	240	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
A	262	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	12	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	16	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	131	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	240	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
B	262	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	1	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	12	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	16	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	131	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	240	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
C	262	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
D	1	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
D	12	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
D	16	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	131	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
D	240	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6
D	262	MSE	MET	MODIFIED RESIDUE	UNP Q5SLQ6

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	230	Total	O	0	0
			230	230		
3	B	267	Total	O	0	0
			267	267		
3	C	168	Total	O	0	0
			168	168		
3	D	191	Total	O	0	0
			191	191		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-hydroxyisobutyrate dehydrogenase

Chain A:




- Molecule 1: 3-hydroxyisobutyrate dehydrogenase

Chain B:




- Molecule 1: 3-hydroxyisobutyrate dehydrogenase

Chain C:



- Molecule 1: 3-hydroxyisobutyrate dehydrogenase

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	85.88Å 106.37Å 168.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 28.95 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.0 (30.00-1.80) 97.1 (28.95-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.33 (at 1.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.182 , 0.200 0.182 , 0.200	Depositor DCC
$R_{free}$ test set	13907 reflections (10.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.7	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 35.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 139226 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9812	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/2239	0.62	0/3027
1	B	0.30	0/2239	0.62	0/3027
1	C	0.28	0/2247	0.59	0/3037
1	D	0.29	0/2239	0.58	0/3027
All	All	0.29	0/8964	0.60	0/12118

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2201	0	2239	24	0
1	B	2201	0	2239	29	0
1	C	2209	0	2251	44	0
1	D	2201	0	2239	25	0
2	A	48	0	26	0	0
2	B	48	0	26	0	0
2	C	48	0	26	1	0
3	A	230	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	267	0	0	2	0
3	C	168	0	0	2	0
3	D	191	0	0	1	0
All	All	9812	0	9046	111	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 6.

All (111) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:252:LEU:HD23	1:D:255:LEU:HD23	1.51	0.90
1:B:282:ARG:NH1	1:C:266:GLU:OE1	2.13	0.81
1:C:221:ARG:HD3	1:C:289:ARG:HD3	1.69	0.74
1:B:192:SER:HB3	1:B:195:LYS:HB2	1.72	0.71
1:C:40:GLN:HG3	1:C:45:SER:O	1.95	0.67
1:C:120:THR:O	1:C:124:GLU:HG3	1.96	0.65
1:C:1:MSE:HG2	1:C:2:GLU:H	1.62	0.65
1:C:130:VAL:HG12	1:C:132:LEU:HD22	1.77	0.65
1:C:189:GLN:NE2	1:D:242:VAL:HG13	2.11	0.65
1:A:120:THR:O	1:A:124:GLU:HG3	1.99	0.63
1:D:114:ALA:HB2	1:D:132:LEU:HD23	1.81	0.62
1:C:95:GLU:HG3	1:C:99:ARG:HH12	1.64	0.62
1:C:116:VAL:HG12	1:C:130:VAL:HG22	1.82	0.61
1:C:1:MSE:CG	1:C:2:GLU:H	2.17	0.57
1:B:99:ARG:HD3	3:B:1546:HOH:O	2.03	0.57
1:C:189:GLN:HE22	1:D:242:VAL:HG13	1.70	0.57
1:B:77:TYR:HB3	1:B:78:PRO:HD3	1.86	0.56
1:D:7:ILE:HD13	1:D:72:VAL:HG11	1.88	0.56
1:B:98:ARG:O	1:B:102:GLU:HG3	2.05	0.56
1:C:63:LEU:O	1:C:90:THR:HG23	2.05	0.56
1:C:23:ARG:HD3	3:C:1416:HOH:O	2.05	0.55
1:D:67:ARG:O	1:D:71:GLU:HG3	2.07	0.55
1:C:207:SER:HA	1:D:207:SER:HA	1.90	0.54
1:C:95:GLU:HG3	1:C:99:ARG:NH1	2.22	0.54
1:A:77:TYR:HB3	1:A:78:PRO:HD3	1.90	0.54
1:B:102:GLU:HG2	1:B:105:ARG:NH2	2.22	0.53
1:C:7:ILE:HB	1:C:61:THR:HG22	1.90	0.53
1:C:197:LEU:HD13	1:C:215:PRO:HG3	1.91	0.52
1:A:172:LEU:HD13	1:A:172:LEU:C	2.30	0.52
1:B:40:GLN:HG3	1:B:45:SER:O	2.09	0.52
1:C:251:PRO:HB2	1:D:255:LEU:HD21	1.92	0.52
1:B:289:ARG:OXT	1:B:289:ARG:HG2	2.10	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:1:MSE:HG2	1:C:2:GLU:N	2.25	0.52
1:B:77:TYR:O	1:B:107:LYS:HE3	2.09	0.52
1:A:85:TYR:CE2	1:A:141:ARG:HG2	2.44	0.51
1:C:14:TYR:HB3	1:C:15:PRO:CD	2.41	0.51
1:B:63:LEU:O	1:B:90:THR:HG23	2.09	0.51
1:A:252:LEU:HD23	1:B:255:LEU:HD23	1.91	0.51
1:A:197:LEU:HD13	1:A:215:PRO:HG3	1.92	0.51
1:D:77:TYR:HB3	1:D:78:PRO:HD3	1.92	0.51
1:B:137:GLU:H	1:B:137:GLU:CD	2.13	0.50
1:C:218:VAL:O	1:C:221:ARG:HD2	2.12	0.50
1:D:215:PRO:HG2	1:D:216:GLN:OE1	2.12	0.49
1:D:206:ARG:HD2	1:D:211:GLU:OE1	2.12	0.49
1:C:14:TYR:OH	1:C:38:ARG:HG2	2.12	0.49
1:C:143:ARG:N	1:C:144:PRO:HD2	2.27	0.49
1:D:143:ARG:N	1:D:144:PRO:HD2	2.27	0.49
1:B:14:TYR:HB3	1:B:15:PRO:CD	2.43	0.48
1:A:207:SER:HA	1:B:207:SER:HA	1.96	0.48
1:D:94:PRO:O	1:D:98:ARG:HG3	2.14	0.48
1:B:113:ASP:HB3	1:B:162:HIS:NE2	2.29	0.48
1:B:116:VAL:HG12	1:B:130:VAL:HG22	1.95	0.48
1:A:236:LEU:O	1:A:240:MSE:HG2	2.14	0.47
1:D:37:LEU:O	1:D:41:GLU:HG3	2.15	0.47
1:A:282:ARG:HD3	3:A:1485:HOH:O	2.14	0.47
1:B:172:LEU:HD13	1:B:172:LEU:C	2.36	0.47
1:B:113:ASP:N	1:B:113:ASP:OD2	2.46	0.46
1:A:255:LEU:HD23	1:B:252:LEU:HD23	1.98	0.46
1:C:1:MSE:CG	1:C:2:GLU:N	2.79	0.45
1:D:40:GLN:HG3	1:D:45:SER:O	2.17	0.45
1:C:84:THR:HG22	1:C:85:TYR:N	2.31	0.45
1:A:7:ILE:HD13	1:A:72:VAL:HG11	1.98	0.45
1:D:141:ARG:HD3	3:D:448:HOH:O	2.17	0.45
1:A:95:GLU:HG3	1:A:99:ARG:NH1	2.32	0.45
1:D:34:GLU:HG3	1:D:35:LYS:H	1.82	0.45
1:A:115:PRO:HG2	1:A:131:MSE:HG3	1.99	0.45
1:D:174:VAL:HG21	1:D:253:LEU:HD23	1.99	0.45
1:C:3:LYS:HD2	3:C:1431:HOH:O	2.18	0.44
1:C:172:LEU:C	1:C:172:LEU:HD13	2.37	0.44
1:C:112:LEU:CD2	1:C:134:GLY:HA3	2.48	0.44
1:C:236:LEU:O	1:C:240:MSE:HG2	2.17	0.44
1:C:217:ARG:HD2	1:C:274:VAL:HG21	2.00	0.44
1:C:252:LEU:HD23	1:D:255:LEU:CD2	2.36	0.44
1:A:14:TYR:HB3	1:A:15:PRO:CD	2.48	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:173:ALA:HB1	1:B:260:TYR:CE2	2.53	0.44
1:A:114:ALA:HB2	1:A:132:LEU:HD23	2.00	0.43
1:B:216:GLN:N	1:B:216:GLN:OE1	2.44	0.43
1:B:136:GLU:HG2	1:B:137:GLU:OE2	2.17	0.43
1:B:141:ARG:HD2	3:B:1417:HOH:O	2.17	0.43
1:D:70:TYR:O	1:D:74:GLU:HG3	2.18	0.43
1:C:5:ALA:HA	1:C:27:LEU:O	2.18	0.43
1:D:14:TYR:HB3	1:D:15:PRO:CD	2.48	0.43
1:A:255:LEU:CD2	1:B:252:LEU:HD23	2.49	0.43
1:B:143:ARG:N	1:B:144:PRO:HD2	2.33	0.43
1:B:5:ALA:HA	1:B:27:LEU:O	2.19	0.43
1:C:14:TYR:HB3	1:C:15:PRO:HD3	2.01	0.42
1:D:218:VAL:O	1:D:221:ARG:HD2	2.19	0.42
1:D:7:ILE:HG23	1:D:29:TRP:CE3	2.54	0.42
1:C:3:LYS:HG2	1:C:25:PRO:HB2	2.02	0.42
1:C:173:ALA:HB1	1:C:260:TYR:CE2	2.55	0.42
1:D:258:GLU:O	1:D:262:MSE:HG3	2.20	0.42
1:A:113:ASP:HB3	1:A:162:HIS:NE2	2.35	0.42
1:B:7:ILE:HB	1:B:61:THR:HG22	2.01	0.42
1:C:207:SER:O	1:C:211:GLU:HG3	2.19	0.42
1:C:31:ARG:HB3	2:C:1303:NDP:O1X	2.20	0.42
1:A:67:ARG:O	1:A:71:GLU:HG3	2.20	0.41
1:C:34:GLU:H	1:C:34:GLU:CD	2.24	0.41
1:A:115:PRO:HG2	1:A:131:MSE:CG	2.50	0.41
1:A:218:VAL:O	1:A:221:ARG:HD2	2.20	0.41
1:C:7:ILE:HD11	1:C:53:VAL:HG11	2.01	0.41
1:C:214:ILE:HB	1:C:215:PRO:HD3	2.02	0.41
1:A:207:SER:O	1:A:211:GLU:HG3	2.21	0.41
1:C:48:VAL:HB	1:C:49:PRO:HD2	2.03	0.41
1:D:236:LEU:O	1:D:240:MSE:HG2	2.21	0.40
1:C:48:VAL:HB	1:C:52:ARG:HD3	2.02	0.40
1:B:184:LEU:C	1:B:184:LEU:HD13	2.42	0.40
1:A:77:TYR:O	1:A:107:LYS:HE2	2.21	0.40
1:B:85:TYR:CE2	1:B:141:ARG:HG2	2.56	0.40
1:C:66:THR:HG22	1:C:70:TYR:CD2	2.57	0.40
1:A:51:GLU:CD	1:A:51:GLU:H	2.24	0.40
1:A:143:ARG:N	1:A:144:PRO:HD2	2.37	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/289 (99%)	281 (98%)	5 (2%)	0	100	100
1	B	286/289 (99%)	279 (98%)	7 (2%)	0	100	100
1	C	287/289 (99%)	280 (98%)	7 (2%)	0	100	100
1	D	286/289 (99%)	284 (99%)	2 (1%)	0	100	100
All	All	1145/1156 (99%)	1124 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/216 (102%)	218 (99%)	3 (1%)	78	69
1	B	221/216 (102%)	218 (99%)	3 (1%)	78	69
1	C	222/216 (103%)	217 (98%)	5 (2%)	63	46
1	D	221/216 (102%)	218 (99%)	3 (1%)	78	69
All	All	885/864 (102%)	871 (98%)	14 (2%)	75	63

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	ARG
1	A	194	GLU
1	A	234	LYS
1	B	113	ASP
1	B	188	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	282	ARG
1	C	113	ASP
1	C	131	MSE
1	C	234	LYS
1	C	255	LEU
1	C	282	ARG
1	D	34	GLU
1	D	141	ARG
1	D	234	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	175	ASN
1	C	273	HIS
1	D	201	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NDP	A	1302	-	52,52,52	1.77	7 (13%)	80,80,80	1.47	10 (12%)
2	NDP	B	1301	-	52,52,52	1.80	9 (17%)	80,80,80	1.49	9 (11%)
2	NDP	C	1303	-	52,52,52	1.78	7 (13%)	80,80,80	1.45	12 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1302	-	-	0/35/77/77	0/3/5/5
2	NDP	B	1301	-	-	0/35/77/77	0/3/5/5
2	NDP	C	1303	-	-	0/35/77/77	0/3/5/5

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1303	NDP	C2N-C3N	5.66	1.45	1.34
2	A	1302	NDP	C2N-C3N	5.63	1.45	1.34
2	B	1301	NDP	C2N-C3N	5.59	1.45	1.34
2	C	1303	NDP	C4N-C5N	-5.55	1.36	1.49
2	B	1301	NDP	C4N-C5N	-5.55	1.36	1.49
2	A	1302	NDP	C4N-C5N	-5.42	1.37	1.49
2	C	1303	NDP	C4N-C3N	-4.94	1.40	1.50
2	B	1301	NDP	C4N-C3N	-4.90	1.40	1.50
2	A	1302	NDP	C4N-C3N	-4.85	1.40	1.50
2	A	1302	NDP	C8A-N9A	3.59	1.41	1.36
2	C	1303	NDP	C8A-N9A	3.58	1.41	1.36
2	B	1301	NDP	C8A-N9A	3.55	1.41	1.36
2	C	1303	NDP	C2N-N1N	3.22	1.42	1.36
2	B	1301	NDP	C2N-N1N	3.21	1.42	1.36
2	A	1302	NDP	C2N-N1N	3.19	1.42	1.36
2	C	1303	NDP	C6N-C5N	2.47	1.38	1.33
2	B	1301	NDP	C6N-C5N	2.39	1.38	1.33
2	B	1301	NDP	C4A-N3A	2.27	1.39	1.35
2	B	1301	NDP	C2B-C1B	-2.26	1.48	1.52
2	A	1302	NDP	C6N-C5N	2.25	1.38	1.33
2	B	1301	NDP	C3B-C2B	-2.20	1.47	1.53
2	C	1303	NDP	C4A-N3A	2.17	1.38	1.35
2	A	1302	NDP	C4A-N3A	2.14	1.38	1.35

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1303	NDP	C3N-C2N-N1N	-4.79	116.25	123.05
2	B	1301	NDP	C3N-C2N-N1N	-4.77	116.28	123.05
2	A	1302	NDP	C3N-C2N-N1N	-4.75	116.31	123.05
2	B	1301	NDP	N3A-C2A-N1A	-4.46	124.98	128.71
2	A	1302	NDP	N3A-C2A-N1A	-4.34	125.08	128.71
2	C	1303	NDP	N3A-C2A-N1A	-4.22	125.18	128.71
2	B	1301	NDP	C4B-O4B-C1B	-4.14	105.25	109.75
2	A	1302	NDP	C5N-C4N-C3N	4.06	123.59	112.60
2	B	1301	NDP	C5N-C4N-C3N	4.06	123.59	112.60
2	C	1303	NDP	C5N-C4N-C3N	4.04	123.54	112.60
2	A	1302	NDP	C4B-O4B-C1B	-3.74	105.69	109.75
2	C	1303	NDP	C4B-O4B-C1B	-3.48	105.96	109.75
2	B	1301	NDP	O4B-C1B-N9A	3.06	111.29	108.44
2	A	1302	NDP	O4B-C1B-N9A	2.99	111.22	108.44
2	B	1301	NDP	O7N-C7N-N7N	-2.53	116.55	122.93
2	A	1302	NDP	C8A-N9A-C4A	-2.51	104.98	106.90
2	A	1302	NDP	O7N-C7N-N7N	-2.50	116.61	122.93
2	C	1303	NDP	O7N-C7N-N7N	-2.50	116.61	122.93
2	C	1303	NDP	C8A-N9A-C4A	-2.50	104.99	106.90
2	A	1302	NDP	C4N-C3N-C2N	-2.43	118.72	121.68
2	B	1301	NDP	C4N-C3N-C2N	-2.38	118.78	121.68
2	B	1301	NDP	C8A-N9A-C4A	-2.36	105.10	106.90
2	C	1303	NDP	C4N-C3N-C2N	-2.28	118.91	121.68
2	C	1303	NDP	O4B-C1B-N9A	2.24	110.53	108.44
2	C	1303	NDP	O4D-C1D-C2D	-2.15	101.66	106.60
2	C	1303	NDP	C4N-C5N-C6N	-2.12	118.71	122.61
2	B	1301	NDP	C4N-C5N-C6N	-2.11	118.74	122.61
2	A	1302	NDP	C4N-C5N-C6N	-2.10	118.75	122.61
2	C	1303	NDP	O4B-C1B-C2B	2.09	108.90	106.95
2	A	1302	NDP	O2A-PA-O1A	2.01	123.46	112.21
2	C	1303	NDP	O2A-PA-O1A	2.00	123.41	112.21

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	288/289 (99%)	-0.33	1 (0%) 91 90	12, 17, 30, 42	0
1	B	288/289 (99%)	-0.29	2 (0%) 84 83	12, 17, 31, 44	0
1	C	289/289 (100%)	-0.11	2 (0%) 84 83	15, 22, 40, 57	0
1	D	288/289 (99%)	-0.22	3 (1%) 79 76	14, 21, 34, 42	0
All	All	1153/1156 (99%)	-0.24	8 (0%) 84 83	12, 20, 34, 57	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MSE	8.2
1	B	289	ARG	3.0
1	D	270	ASP	2.7
1	C	2	GLU	2.6
1	A	77	TYR	2.5
1	D	289	ARG	2.2
1	D	140	GLU	2.0
1	B	2	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NDP	B	1301	48/48	0.08	-0.38	12,17,21,24	0
2	NDP	C	1303	48/48	0.09	-0.46	18,22,27,28	0
2	NDP	A	1302	48/48	0.07	-0.60	13,16,19,21	0

## 6.5 Other polymers

There are no such residues in this entry.