



Full wwPDB X-ray Structure Validation Report

Jul 17, 2014 – 06:06 PM EDT

PDB ID : 4CWM
Title : High-glycosylation crystal structure of the bifunctional endonuclease (AtBFN2) from Arabidopsis thaliana
Authors : Yu, T.-F.; Maestre-Reyna, M.; Ko, C.-Y.; Ko, T.-P.; Sun, Y.-J.; Lin, T.-Y.; Shaw, J.-F.; Wang, A.H.-J.
Deposited on : 2014-04-03
Resolution : 2.09 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.16 November 2013
Xtriage (Phenix) : dev-1439
EDS : stable23161
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable23161

1 Overall quality at a glance

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3012 (2.10-2.10)
Clashscore	79885	3649 (2.10-2.10)
Ramachandran outliers	78287	3610 (2.10-2.10)
Sidechain outliers	78261	3611 (2.10-2.10)
RSRZ outliers	66119	3013 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	269	
1	B	269	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4817 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDONUCLEASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	1
			2049	1295	350	394	10			
1	B	257	Total	C	N	O	S	0	0	1
			2049	1295	350	394	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	HIS	-	EXPRESSION TAG	UNP Q9C9G4
B	264	HIS	-	EXPRESSION TAG	UNP Q9C9G4

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	5	Total	C	N	O	0	0
			61	34	2	25		
2	B	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	3	Total	Zn	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total 3	Zn 3	0	0

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	3	Total 39	C 22	N 2	O 15	0	0
5	B	3	Total 39	C 22	N 2	O 15	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	233	Total 233	O 233	0	0
6	B	224	Total 224	O 224	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ENDONUCLEASE 2

Chain A: 



• Molecule 1: ENDONUCLEASE 2

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	45.38Å 52.71Å 61.10Å 71.34° 78.57° 76.74°	Depositor
Resolution (Å)	23.28 – 2.09 23.27 – 2.09	Depositor EDS
% Data completeness (in resolution range)	97.3 (23.28-2.09) 97.4 (23.27-2.09)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.09Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.197 , 0.268 0.203 , 0.274	Depositor DCC
R_{free} test set	1506 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	23.5	Xtriage
Anisotropy	0.454	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 29732 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4817	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.34 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.9116e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2098	0.70	0/2847
1	B	0.58	0/2098	0.72	0/2847
All	All	0.58	0/4196	0.71	0/5694

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2049	0	1947	33	0
1	B	2049	0	1947	25	0
2	A	61	0	52	4	0
2	B	61	0	52	4	0
3	A	56	0	50	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	B	78	0	68	2	0
6	A	233	0	0	9	0
6	B	224	0	0	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4817	0	4116	59	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (59) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:41:CYS:HB2	6:A:2010:HOH:O	1.73	0.88
1:A:216:ALA:HB1	1:A:242:ILE:HD11	1.58	0.82
1:B:41:CYS:HB2	6:B:2008:HOH:O	1.83	0.78
1:A:201:LYS:O	1:A:203:THR:N	2.20	0.74
1:B:175:GLU:O	1:B:179:ASP:OD2	2.06	0.74
1:A:91:ASN:HD22	2:A:301:NAG:H83	1.53	0.73
1:B:91:ASN:HD22	2:B:301:NAG:H83	1.53	0.72
1:A:77:GLU:HG2	6:A:2099:HOH:O	1.89	0.71
1:B:88:ALA:HA	2:B:301:NAG:H81	1.75	0.69
1:B:26:LYS:HB2	6:B:2027:HOH:O	1.97	0.64
1:A:88:ALA:HA	2:A:301:NAG:H81	1.81	0.63
1:A:61:ASN:HB3	1:A:134:ALA:HB2	1.81	0.63
1:A:176:GLY:CA	6:A:2160:HOH:O	2.47	0.62
1:A:71:ASN:HD22	1:A:81:LYS:NZ	1.97	0.62
1:A:20:THR:HG22	6:A:2030:HOH:O	1.99	0.62
1:A:201:LYS:C	1:A:203:THR:N	2.53	0.61
1:B:201:LYS:C	1:B:203:THR:N	2.55	0.59
5:B:331:NAG:H83	6:B:2110:HOH:O	2.02	0.59
1:B:176:GLY:C	6:B:2146:HOH:O	2.41	0.58
1:B:77:GLU:HG2	6:B:2087:HOH:O	2.06	0.56
1:B:191:ASP:OD1	1:B:191:ASP:C	2.45	0.55
1:B:93:THR:HB	1:B:261:ILE:HD11	1.89	0.55
1:B:132:SER:OG	1:B:138:GLY:HA2	2.07	0.54
1:A:91:ASN:HD22	2:A:301:NAG:C8	2.20	0.54
1:A:71:ASN:HD22	1:A:81:LYS:HZ3	1.56	0.53
1:A:176:GLY:C	6:A:2160:HOH:O	2.48	0.52
1:A:48:LYS:O	1:A:52:HIS:HA	2.12	0.49
1:A:244:TYR:OH	6:A:2157:HOH:O	2.20	0.49
1:B:176:GLY:CA	6:B:2146:HOH:O	2.60	0.49
1:B:55:SER:HB2	1:B:56:PRO:HD3	1.95	0.49
1:B:54:SER:HA	1:B:57:LEU:HD12	1.96	0.48
1:B:191:ASP:OD1	1:B:195:ARG:NH1	2.47	0.48
1:A:132:SER:OG	1:A:138:GLY:HA2	2.14	0.48
1:A:216:ALA:HB1	1:A:242:ILE:CD1	2.35	0.48
1:A:107:SER:OG	1:A:108:GLN:N	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:57:LEU:O	1:B:85:VAL:HA	2.14	0.46
1:B:82:GLY:O	1:B:87:GLY:HA3	2.15	0.46
1:A:53:TRP:CG	2:A:302:NAG:H5	2.51	0.45
1:A:8:ILE:HD13	1:A:177:MET:CE	2.47	0.45
1:A:3:LYS:HD3	1:A:7:GLU:OE2	2.17	0.45
1:B:188:GLU:HG2	6:B:2153:HOH:O	2.16	0.45
1:B:176:GLY:O	6:B:2146:HOH:O	2.21	0.45
1:A:60:ILE:HG13	1:A:85:VAL:HG12	1.99	0.45
1:B:107:SER:OG	1:B:108:GLN:N	2.50	0.44
1:A:185:ILE:HG23	1:A:193:VAL:HG21	2.00	0.44
1:A:9:ILE:HD13	1:A:123:GLY:HA2	1.99	0.43
1:B:181:LEU:HD23	5:B:361:NAG:H81	2.00	0.43
1:A:112:THR:O	1:A:116:LEU:HG	2.19	0.43
1:B:88:ALA:HA	2:B:301:NAG:C8	2.45	0.43
1:B:149:ARG:HD3	6:B:2133:HOH:O	2.19	0.43
1:B:185:ILE:HG23	1:B:193:VAL:HG21	2.00	0.42
1:A:32:SER:HB2	6:A:2045:HOH:O	2.19	0.42
1:A:242:ILE:O	1:A:246:ARG:HG2	2.19	0.42
1:A:239:ARG:O	1:A:242:ILE:HG22	2.20	0.42
1:A:89:ILE:O	1:A:93:THR:HG23	2.19	0.41
1:A:230:THR:HG22	1:A:232:GLU:CG	2.51	0.41
1:B:91:ASN:HD22	2:B:301:NAG:C8	2.26	0.41
1:A:187:THR:C	6:A:2166:HOH:O	2.59	0.41
1:A:171:ASN:ND2	6:A:2156:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	251/269 (93%)	238 (95%)	13 (5%)	0	100	100
1	B	251/269 (93%)	238 (95%)	13 (5%)	0	100	100
All	All	502/538 (93%)	476 (95%)	26 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/226 (95%)	206 (96%)	9 (4%)	40	38
1	B	215/226 (95%)	208 (97%)	7 (3%)	50	51
All	All	430/452 (95%)	414 (96%)	16 (4%)	45	45

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LYS
1	A	32	SER
1	A	75	LYS
1	A	108	GLN
1	A	118	VAL
1	A	127	GLN
1	A	201	LYS
1	A	223	LYS
1	A	242	ILE
1	B	75	LYS
1	B	118	VAL
1	B	149	ARG
1	B	150	LYS
1	B	191	ASP
1	B	194	LYS
1	B	203	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	ASN
1	A	71	ASN
1	A	171	ASN
1	B	61	ASN
1	B	71	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	301	1,2	12,14,15	1.07	1 (8%)	15,19,21	1.63	5 (33%)
2	NAG	A	302	2	12,14,15	0.74	1 (8%)	15,19,21	2.17	4 (26%)
2	BMA	A	303	2	10,11,12	0.83	0	11,15,17	1.08	1 (9%)
2	MAN	A	304	2	10,11,12	0.88	1 (10%)	11,15,17	0.99	0
2	MAN	A	305	2	10,11,12	0.80	1 (10%)	11,15,17	1.03	1 (9%)
3	NAG	A	331	1,3	12,14,15	0.84	1 (8%)	15,19,21	1.16	1 (6%)
3	NAG	A	332	3	12,14,15	0.70	1 (8%)	15,19,21	1.46	2 (13%)
3	NAG	A	361	1,3	12,14,15	0.94	1 (8%)	15,19,21	1.44	1 (6%)
3	NAG	A	362	3	12,14,15	0.70	0	15,19,21	1.45	3 (20%)
2	NAG	B	301	1,2	12,14,15	1.00	1 (8%)	15,19,21	1.98	6 (40%)
2	NAG	B	302	2	12,14,15	0.74	0	15,19,21	1.43	2 (13%)
2	BMA	B	303	2	10,11,12	0.84	0	11,15,17	1.25	1 (9%)
2	MAN	B	304	2	10,11,12	0.94	1 (10%)	11,15,17	2.20	4 (36%)
2	MAN	B	305	2	10,11,12	0.82	1 (10%)	11,15,17	0.85	0
5	NAG	B	331	1,5	12,14,15	0.76	1 (8%)	15,19,21	1.37	3 (20%)
5	NAG	B	332	5	12,14,15	0.76	1 (8%)	15,19,21	1.25	2 (13%)
5	BMA	B	333	5	10,11,12	0.81	0	11,15,17	1.17	2 (18%)
5	NAG	B	361	1,5	12,14,15	1.01	2 (16%)	15,19,21	0.92	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	362	5	12,14,15	0.84	1 (8%)	15,19,21	1.47	4 (26%)
5	BMA	B	363	5	10,11,12	1.05	1 (10%)	11,15,17	1.31	3 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	301	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	302	2	-	0/6/23/26	0/1/1/1
2	BMA	A	303	2	-	0/2/19/22	0/1/1/1
2	MAN	A	304	2	-	0/2/19/22	0/1/1/1
2	MAN	A	305	2	-	0/2/19/22	0/1/1/1
3	NAG	A	331	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	332	3	-	0/6/23/26	0/1/1/1
3	NAG	A	361	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	362	3	-	0/6/23/26	0/1/1/1
2	NAG	B	301	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	302	2	-	0/6/23/26	0/1/1/1
2	BMA	B	303	2	-	0/2/19/22	0/1/1/1
2	MAN	B	304	2	-	0/2/19/22	0/1/1/1
2	MAN	B	305	2	-	0/2/19/22	0/1/1/1
5	NAG	B	331	1,5	-	0/6/23/26	0/1/1/1
5	NAG	B	332	5	-	0/6/23/26	0/1/1/1
5	BMA	B	333	5	-	0/2/19/22	0/1/1/1
5	NAG	B	361	1,5	-	0/6/23/26	0/1/1/1
5	NAG	B	362	5	-	0/6/23/26	0/1/1/1
5	BMA	B	363	5	-	0/2/19/22	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	NAG	O5-C5	-3.34	1.40	1.45
2	B	301	NAG	O5-C5	-3.13	1.40	1.45
3	A	331	NAG	O5-C5	-2.81	1.41	1.45
5	B	361	NAG	O5-C5	-2.60	1.41	1.45
3	A	361	NAG	O5-C5	-2.60	1.41	1.45
5	B	363	BMA	O5-C5	-2.60	1.41	1.45
2	B	304	MAN	O5-C5	-2.45	1.41	1.45
5	B	331	NAG	O5-C5	-2.43	1.41	1.45
5	B	362	NAG	O5-C5	-2.40	1.41	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	332	NAG	O5-C5	-2.13	1.42	1.45
2	B	305	MAN	O5-C5	-2.11	1.42	1.45
5	B	332	NAG	O5-C5	-2.09	1.42	1.45
2	A	302	NAG	O5-C5	-2.07	1.42	1.45
2	A	305	MAN	O5-C5	-2.04	1.42	1.45
5	B	361	NAG	C2-N2	-2.03	1.44	1.46
2	A	304	MAN	O5-C5	-2.02	1.42	1.45

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	NAG	O5-C5-C6	6.76	114.08	106.98
2	B	304	MAN	C4-C3-C2	5.24	116.46	110.61
3	A	361	NAG	O5-C5-C6	4.14	111.33	106.98
2	B	301	NAG	C2-N2-C7	3.74	127.92	123.39
3	A	332	NAG	O5-C5-C6	3.73	110.89	106.98
2	A	301	NAG	C2-N2-C7	3.61	127.76	123.39
2	B	301	NAG	C8-C7-N2	3.37	122.49	116.12
3	A	332	NAG	O5-C5-C4	-3.21	106.58	110.65
5	B	331	NAG	C3-C2-N2	3.12	116.31	111.62
2	B	302	NAG	O5-C5-C6	3.09	110.22	106.98
5	B	362	NAG	C3-C2-N2	-2.89	107.28	111.62
2	A	301	NAG	C8-C7-N2	2.79	121.40	116.12
5	B	362	NAG	O5-C5-C4	2.71	114.10	110.65
2	B	301	NAG	O5-C5-C4	2.70	114.08	110.65
2	B	304	MAN	C3-C4-C5	2.67	114.97	110.17
5	B	362	NAG	C2-N2-C7	-2.65	120.19	123.39
5	B	331	NAG	C8-C7-N2	2.62	121.06	116.12
2	B	301	NAG	O7-C7-C8	-2.57	117.12	122.04
5	B	332	NAG	C4-C3-C2	2.55	116.09	110.74
5	B	333	BMA	C3-C4-C5	-2.54	105.62	110.17
2	A	301	NAG	O4-C4-C5	-2.53	102.56	109.25
2	A	305	MAN	C4-C3-C2	2.51	113.41	110.61
2	A	302	NAG	O3-C3-C2	2.49	114.23	109.16
2	A	302	NAG	O7-C7-C8	-2.47	117.31	122.04
5	B	333	BMA	O5-C5-C6	2.40	109.50	106.98
5	B	363	BMA	O4-C4-C3	-2.38	105.04	110.36
2	B	301	NAG	O4-C4-C5	-2.35	103.05	109.25
3	A	362	NAG	C3-C4-C5	-2.34	105.97	110.17
5	B	363	BMA	C3-C4-C5	2.30	114.29	110.17
2	A	301	NAG	O7-C7-C8	-2.24	117.75	122.04
3	A	362	NAG	C2-N2-C7	-2.23	120.70	123.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	332	NAG	C3-C2-N2	-2.21	108.31	111.62
2	A	302	NAG	C2-N2-C7	2.20	126.06	123.39
3	A	331	NAG	C8-C7-N2	2.19	120.25	116.12
2	B	304	MAN	O5-C5-C6	2.19	109.27	106.98
2	A	303	BMA	O5-C5-C6	2.18	109.27	106.98
5	B	362	NAG	O4-C4-C3	-2.12	105.62	110.36
5	B	331	NAG	O7-C7-C8	-2.12	117.98	122.04
2	B	303	BMA	O5-C5-C6	2.12	109.20	106.98
2	B	302	NAG	O5-C5-C4	2.08	113.29	110.65
5	B	363	BMA	O5-C5-C6	2.05	109.13	106.98
2	A	301	NAG	O5-C5-C4	2.05	113.25	110.65
2	B	301	NAG	C3-C4-C5	2.04	113.84	110.17
2	B	304	MAN	O3-C3-C2	-2.02	106.08	109.74
3	A	362	NAG	O7-C7-C8	-2.01	118.19	122.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/269 (95%)	0.17	7 (2%) 52 57	15, 25, 41, 57	0
1	B	257/269 (95%)	0.19	3 (1%) 75 80	14, 25, 43, 59	0
All	All	514/538 (95%)	0.18	10 (1%) 64 68	14, 25, 42, 59	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	173	ALA	4.3
1	B	107	SER	3.4
1	A	172	SER	3.4
1	B	172	SER	3.0
1	A	32	SER	2.8
1	A	107	SER	2.8
1	B	33	ALA	2.3
1	A	98	SER	2.3
1	A	174	LEU	2.3
1	A	79	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	BMA	A	303	11/12	0.18	13.19	29,37,45,51	0
5	BMA	B	333	11/12	0.21	5.12	64,67,70,70	0
2	BMA	B	303	11/12	0.15	3.39	38,48,57,63	0
2	MAN	A	304	11/12	0.17	2.00	31,40,42,44	0
3	NAG	A	332	14/15	0.20	1.75	46,50,53,57	0
3	NAG	A	361	14/15	0.14	0.87	24,29,36,38	0
2	MAN	B	304	11/12	0.18	0.83	51,55,58,61	0
3	NAG	A	331	14/15	0.13	0.82	30,36,42,49	0
5	NAG	B	361	14/15	0.13	0.29	23,30,35,38	0
2	NAG	B	301	14/15	0.13	0.10	18,22,24,24	0
5	NAG	B	331	14/15	0.12	-0.08	30,35,38,44	0
2	NAG	B	302	14/15	0.16	-0.18	24,27,32,37	0
2	NAG	A	301	14/15	0.11	-0.40	19,21,24,24	0
5	NAG	B	332	14/15	0.12	-0.74	41,45,49,58	0
2	NAG	A	302	14/15	0.14	-0.77	25,27,31,31	0
2	MAN	A	305	11/12	0.19	-	43,55,58,61	0
3	NAG	A	362	14/15	0.26	-	43,49,51,52	0
5	BMA	B	363	11/12	0.15	-	48,51,53,56	0
5	NAG	B	362	14/15	0.21	-	40,45,51,52	0
2	MAN	B	305	11/12	0.24	-	58,65,66,66	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ZN	B	433	1/1	0.07	-1.91	17,17,17,17	0
4	ZN	B	432	1/1	0.07	-2.44	20,20,20,20	0
4	ZN	A	433	1/1	0.05	-3.08	15,15,15,15	0
4	ZN	A	431	1/1	0.02	-4.43	22,22,22,22	0
4	ZN	A	432	1/1	0.04	-4.74	18,18,18,18	0
4	ZN	B	431	1/1	0.03	-6.96	22,22,22,22	0

6.5 Other polymers ⓘ

There are no such residues in this entry.