



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2014 – 12:33 AM GMT

PDB ID : 3CXL
Title : Crystal structure of human chimerin 1 (CHN1)
Authors : Shen, L.; Buck, M.; Tong, Y.; Tempel, W.; MacKenzie, F.; Arrowsmith, C.H.;
Edwards, A.M.; Bountra, C.; Wilkstrom, M.; Bochkarev, A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2008-04-24
Resolution : 2.60 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : **FAILED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	A	463	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3170 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-chimerin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	402	3167	2020	543	580	24	0	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	MET	-	EXPRESSION TAG	UNP P15882
A	-2	HIS	-	EXPRESSION TAG	UNP P15882
A	-1	HIS	-	EXPRESSION TAG	UNP P15882
A	0	HIS	-	EXPRESSION TAG	UNP P15882
A	1	HIS	-	EXPRESSION TAG	UNP P15882
A	2	HIS	-	EXPRESSION TAG	UNP P15882
A	3	HIS	-	EXPRESSION TAG	UNP P15882
A	4	SER	-	EXPRESSION TAG	UNP P15882
A	5	SER	-	EXPRESSION TAG	UNP P15882
A	6	GLY	-	EXPRESSION TAG	UNP P15882
A	7	ARG	-	EXPRESSION TAG	UNP P15882
A	8	GLU	-	EXPRESSION TAG	UNP P15882
A	9	ASN	-	EXPRESSION TAG	UNP P15882
A	10	LEU	-	EXPRESSION TAG	UNP P15882
A	11	TYR	-	EXPRESSION TAG	UNP P15882
A	12	PHE	-	EXPRESSION TAG	UNP P15882
A	13	GLN	-	EXPRESSION TAG	UNP P15882
A	14	GLY	-	EXPRESSION TAG	UNP P15882

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is UNKNOWN ATOM OR ION (three-letter code: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	X	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS failed to run properly.

- Molecule 1: N-chimerin

Chain A: 



4 Data and refinement statistics i

EDS failed to run properly - this section will therefore be incomplete.

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, α , β , γ	152.19Å 152.19Å 69.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.51 – 2.60	Depositor
% Data completeness (in resolution range)	99.5 (28.51-2.60)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.266 , 0.300	Depositor
Wilson B-factor (Å ²)	66.7	Xtriage
Anisotropy	0.058	Xtriage
Estimated twinning fraction	0.348 for h,-h-k,-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 28308 reflections	Xtriage
Total number of atoms	3170	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	1/3237 (0.0%)	0.57	1/4384 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	391	CYS	CB-SG	-5.25	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	162	PRO	N-CA-CB	5.63	110.06	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3167	0	3047	49	0
2	A	2	0	0	0	0
3	A	1	0	0	0	0
All	All	3170	0	3047	49	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 8.

All (49) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:60:ASP:O	1:A:64:ILE:HG23	1.83	0.79
1:A:142:ILE:HD13	1:A:223:ALA:HB2	1.70	0.73
1:A:401:LEU:O	1:A:404:VAL:HG22	1.92	0.69
1:A:401:LEU:O	1:A:405:THR:HG23	1.99	0.61
1:A:429:SER:OG	1:A:437:ALA:HB1	1.99	0.61
1:A:46:TYR:CD1	1:A:113:ILE:HD11	2.37	0.59
1:A:134:ILE:HA	1:A:137:MET:HG3	1.86	0.58
1:A:295:ARG:HD2	1:A:336:ILE:HG23	1.86	0.57
1:A:420:ILE:HA	1:A:445:ARG:CD	2.35	0.57
1:A:262:VAL:O	1:A:262:VAL:HG22	2.07	0.54
1:A:46:TYR:HD1	1:A:113:ILE:HD11	1.71	0.54
1:A:404:VAL:HG23	1:A:418:LEU:HD11	1.89	0.54
1:A:284:VAL:HG22	1:A:319:PHE:CG	2.43	0.54
1:A:86:LEU:HD11	1:A:121:LEU:HD21	1.90	0.53
1:A:134:ILE:HA	1:A:137:MET:CG	2.39	0.53
1:A:405:THR:HG22	1:A:418:LEU:HD12	1.90	0.53
1:A:420:ILE:HA	1:A:445:ARG:HD2	1.91	0.52
1:A:97:LEU:HD23	1:A:105:VAL:HG12	1.92	0.51
1:A:20:LEU:HD13	1:A:227:TRP:CZ3	2.46	0.50
1:A:265:VAL:HG11	1:A:284:VAL:HG23	1.93	0.50
1:A:139:ILE:HD12	1:A:139:ILE:C	2.32	0.49
1:A:288:CYS:O	1:A:292:ILE:HG12	2.13	0.48
1:A:259:LEU:O	1:A:262:VAL:HG13	2.14	0.47
1:A:274:VAL:HG13	1:A:279:THR:O	2.14	0.47
1:A:404:VAL:CG2	1:A:418:LEU:HD11	2.45	0.47
1:A:235:LYS:NZ	1:A:242:ASN:OD1	2.40	0.47
1:A:225:PHE:CE2	1:A:227:TRP:HA	2.51	0.46
1:A:213:PHE:CE1	1:A:226:MET:HE3	2.51	0.46
1:A:213:PHE:CE1	1:A:226:MET:CE	2.99	0.45
1:A:267:SER:HA	1:A:349:ASP:O	2.16	0.45
1:A:336:ILE:HD12	1:A:337:ASN:N	2.32	0.45
1:A:368:LYS:O	1:A:370:MET:HE2	2.17	0.45
1:A:420:ILE:HD13	1:A:445:ARG:HD2	1.99	0.44
1:A:49:GLU:O	1:A:51:HIS:ND1	2.46	0.44
1:A:415:ALA:HB3	1:A:453:LYS:NZ	2.33	0.44
1:A:213:PHE:CZ	1:A:226:MET:HE1	2.53	0.43
1:A:400:HIS:O	1:A:404:VAL:HG13	2.18	0.43
1:A:55:SER:O	1:A:58:ALA:HB3	2.20	0.42
1:A:355:ILE:HD12	1:A:427:MET:HG2	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:146:VAL:HG12	1:A:147:GLY:N	2.35	0.42
1:A:270:LEU:HD23	1:A:390:HIS:CG	2.55	0.42
1:A:35:CYS:SG	1:A:37:CYS:HB2	2.61	0.41
1:A:284:VAL:HG22	1:A:319:PHE:CD1	2.56	0.41
1:A:433:ASP:HB3	1:A:436:ALA:HB3	2.02	0.41
1:A:206:HIS:CE1	1:A:255:CYS:HA	2.55	0.41
1:A:355:ILE:HD12	1:A:427:MET:CG	2.51	0.41
1:A:289:ILE:CD1	1:A:400:HIS:CD2	3.04	0.40
1:A:230:ILE:O	1:A:231:ALA:C	2.60	0.40
1:A:417:ASN:OD1	1:A:418:LEU:N	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	396/463 (86%)	387 (98%)	8 (2%)	1 (0%)	50 77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	328/413 (79%)	321 (98%)	7 (2%)	66 90

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	CYS
1	A	61	GLN
1	A	91	GLN
1	A	262	VAL
1	A	283	MET
1	A	295	ARG
1	A	438	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is unknown and 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS failed to run properly - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS failed to run properly - this section will therefore be empty.

6.4 Ligands ⓘ

EDS failed to run properly - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS failed to run properly - this section will therefore be empty.