



Full wwPDB X-ray Structure Validation Report

Mar 1, 2014 – 04:04 AM GMT

PDB ID : 3D0G
Title : Crystal structure of spike protein receptor-binding domain from the 2002-2003 SARS coronavirus human strain complexed with human-civet chimeric receptor ACE2
Authors : Li, F.
Deposited on : 2008-05-01
Resolution : 2.80 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2295 (2.80-2.80)
Ramachandran outliers	78287	2252 (2.80-2.80)
Sidechain outliers	78261	2254 (2.80-2.80)
RSRZ outliers	66119	1802 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	597	
1	B	597	
2	E	179	
2	F	179	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NDG	B	617	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 12643 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Angiotensin-convertingenzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	597	Total	C	N	O	S	0	0	0
			4864	3110	803	922	29			
1	B	597	Total	C	N	O	S	0	0	0
			4864	3110	803	922	29			

- Molecule 2 is a protein called Spike protein S1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	173	Total	C	N	O	S	0	0	0
			1397	906	227	258	6			
2	F	173	Total	C	N	O	S	0	0	0
			1397	906	227	258	6			

- Molecule 3 is SUGAR (2-(ACETYLAMINO)-2-DEOXY-A-D-GLUCOPYRANOSE) (three-letter code: NDG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	8	1	6		
3	B	1	Total	C	N	O	0	0
			15	8	1	6		
3	B	1	Total	C	N	O	0	0
			15	8	1	6		
3	B	1	Total	C	N	O	0	0
			15	8	1	6		
3	F	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		
5	A	1	Total	Zn	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total 1	Cl 1	0	0
6	A	1	Total 1	Cl 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	21	Total 21	O 21	0	0
7	F	6	Total 6	O 6	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Angiotensin-convertingenzyme 2



• Molecule 1: Angiotensin-convertingenzyme 2




• Molecule 2: Spike protein S1





● Molecule 2: Spike protein S1

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.01Å 119.76Å 108.80Å 90.00° 96.22° 90.00°	Depositor
Resolution (Å)	47.14 – 2.80 47.16 – 2.79	Depositor EDS
% Data completeness (in resolution range)	93.8 (47.14-2.80) 88.7 (47.16-2.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.214 , 0.279 0.210 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	67.8	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 48072 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12643	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG, NDG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/5000	0.60	0/6796
1	B	0.52	0/5000	0.63	2/6796 (0.0%)
2	E	0.52	0/1442	0.63	0/1963
2	F	0.51	0/1442	0.64	0/1963
All	All	0.51	0/12884	0.62	2/17518 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	456	LEU	CA-CB-CG	5.75	128.52	115.30
1	B	421	ILE	CB-CA-C	-5.23	101.14	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4864	0	0	16	0
1	B	4864	0	0	12	0
2	E	1397	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	1397	0	0	5	0
3	A	15	0	0	0	0
3	B	45	0	0	2	0
3	F	15	0	0	0	0
4	E	15	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	B	21	0	0	1	0
7	F	6	0	0	0	0
All	All	12643	0	0	39	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (39) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:457:GLU:OE2	1:A:460:ARG:NH1	2.16	0.78
1:B:183:TYR:OH	1:B:509:ASP:OD1	2.16	0.63
2:E:416:PHE:N	2:E:416:PHE:CD1	2.67	0.61
1:A:183:TYR:OH	1:A:509:ASP:OD1	2.20	0.58
1:B:546:ASN:ND2	3:B:618:NDG:O1L	2.37	0.58
1:A:560:LEU:O	1:A:563:SER:CB	2.52	0.57
1:A:125:THR:O	1:A:129:THR:OG1	2.23	0.56
1:A:493:HIS:CD2	1:A:499:ASP:OD1	2.61	0.54
1:B:546:ASN:ND2	3:B:618:NDG:C1	2.71	0.54
2:E:436:TYR:OH	2:E:484:TYR:OH	2.26	0.53
1:A:511:SER:O	1:A:514:ARG:NH1	2.43	0.52
2:E:329:PHE:O	2:E:330:ASN:CB	2.57	0.52
1:A:38:GLU:OE2	1:A:353:LYS:NZ	2.45	0.50
1:A:269:ASP:OD2	1:A:272:GLY:N	2.45	0.49
2:F:416:PHE:CD1	2:F:416:PHE:N	2.82	0.48
1:A:557:MET:C	1:A:557:MET:SD	2.92	0.48
1:B:401:HIS:CB	7:B:904:HOH:O	2.62	0.48
2:E:436:TYR:O	2:E:480:ASP:OD2	2.32	0.47
2:E:329:PHE:N	2:E:329:PHE:CD2	2.83	0.47
1:B:482:ARG:NH2	1:B:489:GLU:OE1	2.48	0.47
2:F:415:ASP:O	2:F:416:PHE:C	2.53	0.46
2:E:390:LYS:O	2:E:393:ASP:N	2.49	0.46
2:F:329:PHE:O	2:F:330:ASN:CB	2.64	0.46
1:A:225:ASP:O	1:A:229:THR:CG2	2.64	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:F:396:GLN:OE1	2:F:403:GLY:O	2.34	0.46
1:A:378:HIS:ND1	1:A:401:HIS:ND1	2.64	0.45
1:A:273:ARG:O	1:A:449:THR:OG1	2.34	0.45
1:B:47:SER:O	1:B:51:ASN:OD1	2.36	0.44
1:B:397:ASN:OD1	1:B:399:GLY:N	2.50	0.44
2:F:329:PHE:N	2:F:329:PHE:CD2	2.86	0.44
1:A:551:GLY:O	1:A:555:PHE:N	2.52	0.43
1:A:460:ARG:NH2	1:A:510:TYR:O	2.52	0.43
1:B:166:GLU:OE1	1:B:493:HIS:CE1	2.71	0.43
1:A:476:LYS:NZ	1:A:479:GLU:OE1	2.53	0.42
1:B:378:HIS:ND1	1:B:401:HIS:ND1	2.68	0.42
1:A:166:GLU:OE1	1:A:493:HIS:CE1	2.73	0.42
1:B:269:ASP:OD2	1:B:272:GLY:N	2.53	0.42
1:B:239:HIS:O	1:B:240:LEU:C	2.58	0.41
1:B:407:ILE:CD1	1:B:522:GLN:O	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	595/597 (100%)	540 (91%)	51 (9%)	4 (1%)	30 69
1	B	595/597 (100%)	546 (92%)	40 (7%)	9 (2%)	15 46
2	E	169/179 (94%)	138 (82%)	23 (14%)	8 (5%)	4 11
2	F	169/179 (94%)	141 (83%)	22 (13%)	6 (4%)	5 17
All	All	1528/1552 (98%)	1365 (89%)	136 (9%)	27 (2%)	13 39

All (27) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	VAL
2	E	404	VAL
2	F	369	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	370	SER
1	B	536	GLU
2	E	416	PHE
2	F	368	GLY
2	F	416	PHE
1	B	20	THR
1	B	87	GLU
1	B	212	VAL
1	B	213	ASP
2	E	358	SER
2	E	368	GLY
2	F	465	LYS
1	B	340	GLN
2	E	330	ASN
2	E	402	THR
2	E	465	LYS
1	A	289	PRO
1	A	339	VAL
2	E	401	GLN
1	B	84	PRO
1	B	339	VAL
1	B	54	ILE
1	A	147	GLY
2	F	391	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	527/527 (100%)	474 (90%)	53 (10%)	11	30
1	B	527/527 (100%)	473 (90%)	54 (10%)	11	29
2	E	151/157 (96%)	129 (85%)	22 (15%)	5	13
2	F	151/157 (96%)	129 (85%)	22 (15%)	5	13
All	All	1356/1368 (99%)	1205 (89%)	151 (11%)	9	25

All (151) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	GLU
1	A	29	LEU
1	A	55	THR
1	A	60	GLN
1	A	78	THR
1	A	81	GLN
1	A	85	LEU
1	A	86	GLN
1	A	90	ASN
1	A	100	LEU
1	A	103	ASN
1	A	114	LYS
1	A	129	THR
1	A	148	LEU
1	A	156	LEU
1	A	198	ASP
1	A	212	VAL
1	A	240	LEU
1	A	283	VAL
1	A	292	ASP
1	A	293	VAL
1	A	334	THR
1	A	343	VAL
1	A	364	VAL
1	A	368	ASP
1	A	371	THR
1	A	375	GLU
1	A	381	TYR
1	A	385	TYR
1	A	392	LEU
1	A	401	HIS
1	A	410	LEU
1	A	418	LEU
1	A	439	LEU
1	A	444	LEU
1	A	449	THR
1	A	455	MET
1	A	456	LEU
1	A	460	ARG
1	A	476	LYS
1	A	488	VAL
1	A	503	LEU
1	A	514	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	518	ARG
1	A	526	GLN
1	A	549	GLU
1	A	557	MET
1	A	562	LYS
1	A	573	VAL
1	A	585	LEU
1	A	595	LEU
1	A	597	ASP
1	A	600	LYS
1	B	29	LEU
1	B	81	GLN
1	B	89	GLN
1	B	98	GLN
1	B	100	LEU
1	B	103	ASN
1	B	114	LYS
1	B	121	ASN
1	B	125	THR
1	B	137	ASN
1	B	148	LEU
1	B	156	LEU
1	B	187	LYS
1	B	198	ASP
1	B	212	VAL
1	B	240	LEU
1	B	271	TRP
1	B	290	ASN
1	B	291	ILE
1	B	309	LYS
1	B	313	LYS
1	B	333	LEU
1	B	334	THR
1	B	343	VAL
1	B	357	ARG
1	B	360	MET
1	B	375	GLU
1	B	381	TYR
1	B	385	TYR
1	B	392	LEU
1	B	401	HIS
1	B	410	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	416	LYS
1	B	418	LEU
1	B	421	ILE
1	B	427	ASP
1	B	429	GLN
1	B	432	ASN
1	B	439	LEU
1	B	444	LEU
1	B	455	MET
1	B	456	LEU
1	B	488	VAL
1	B	493	HIS
1	B	496	THR
1	B	503	LEU
1	B	526	GLN
1	B	529	LEU
1	B	549	GLU
1	B	553	LYS
1	B	557	MET
1	B	560	LEU
1	B	585	LEU
1	B	593	THR
2	E	329	PHE
2	E	332	THR
2	E	333	LYS
2	E	337	VAL
2	E	338	TYR
2	E	364	PHE
2	E	369	VAL
2	E	395	ARG
2	E	412	LEU
2	E	415	ASP
2	E	416	PHE
2	E	417	MET
2	E	423	TRP
2	E	433	THR
2	E	447	LYS
2	E	455	ILE
2	E	467	CYS
2	E	468	THR
2	E	472	LEU
2	E	480	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	481	TYR
2	E	484	TYR
2	F	329	PHE
2	F	333	LYS
2	F	337	VAL
2	F	341	GLU
2	F	344	LYS
2	F	359	THR
2	F	360	PHE
2	F	364	PHE
2	F	369	VAL
2	F	372	THR
2	F	373	LYS
2	F	390	LYS
2	F	395	ARG
2	F	399	PRO
2	F	412	LEU
2	F	415	ASP
2	F	416	PHE
2	F	417	MET
2	F	426	ARG
2	F	427	ASN
2	F	455	ILE
2	F	485	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NDG	A	616	-	15,15,15	0.47	0	21,21,21	1.21	2 (9%)
3	NDG	B	616	-	15,15,15	0.41	0	21,21,21	0.74	0
3	NDG	B	617	-	15,15,15	0.73	0	21,21,21	1.44	1 (4%)
3	NDG	B	618	-	15,15,15	0.55	0	21,21,21	2.11	3 (14%)
4	NAG	E	91	-	15,15,15	0.52	0	21,21,21	1.63	5 (23%)
3	NDG	F	91	-	15,15,15	0.47	0	21,21,21	1.06	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDG	A	616	-	-	0/6/26/26	0/1/1/1
3	NDG	B	616	-	-	0/6/26/26	0/1/1/1
3	NDG	B	617	-	-	0/6/26/26	0/1/1/1
3	NDG	B	618	-	-	0/6/26/26	0/1/1/1
4	NAG	E	91	-	-	0/6/26/26	0/1/1/1
3	NDG	F	91	-	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	618	NDG	O-C1-C2	7.79	117.51	109.61
3	B	617	NDG	O-C1-C2	4.26	113.93	109.61
4	E	91	NAG	C1-O5-C5	3.78	120.17	113.40
4	E	91	NAG	O5-C1-C2	3.56	113.22	109.61
3	B	618	NDG	C1-O-C5	3.18	119.09	113.40
3	F	91	NDG	O-C1-C2	2.87	112.52	109.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	616	NDG	C4-C3-C2	2.78	114.39	110.44
3	B	618	NDG	C1-C2-C3	2.31	113.90	110.59
4	E	91	NAG	O5-C5-C4	2.24	113.91	109.76
4	E	91	NAG	C1-C2-C3	2.23	113.78	110.59
3	A	616	NDG	C3-C4-C5	2.19	114.12	110.20
3	F	91	NDG	C4-C3-C2	2.17	113.53	110.44
4	E	91	NAG	C4-C3-C2	2.00	113.28	110.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	597/597 (100%)	0.43	50 (8%) 11 9	41, 69, 119, 141	0
1	B	597/597 (100%)	0.37	37 (6%) 20 19	37, 65, 119, 139	0
2	E	173/179 (96%)	0.71	24 (13%) 4 3	54, 79, 123, 135	0
2	F	173/179 (96%)	0.39	9 (5%) 26 26	55, 80, 122, 134	0
All	All	1540/1552 (99%)	0.43	120 (7%) 13 11	37, 70, 121, 141	0

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	VAL	8.2
1	A	105	SER	7.4
1	B	86	GLN	7.3
2	E	501	PHE	5.9
1	A	290	ASN	5.7
1	B	89	GLN	5.2
1	B	615	ASP	4.7
1	A	54	ILE	4.6
1	A	108	LEU	4.6
1	A	425	SER	4.4
1	A	338	ASN	4.4
1	A	104	GLY	4.3
1	A	107	VAL	4.2
1	A	291	ILE	4.2
1	B	339	VAL	4.2
1	A	424	LEU	4.1
1	B	140	GLU	4.1
1	A	428	PHE	4.0
2	E	369	VAL	4.0
1	B	97	LEU	3.9
2	E	356	TYR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	342	ALA	3.8
1	A	364	VAL	3.7
1	B	139	GLN	3.7
1	A	363	LYS	3.7
1	A	89	GLN	3.7
1	A	192	ARG	3.7
1	B	136	ASP	3.6
1	A	106	SER	3.6
2	E	353	SER	3.5
1	B	83	TYR	3.5
2	E	500	SER	3.5
1	A	297	MET	3.4
2	E	352	TYR	3.3
2	F	358	SER	3.3
2	E	472	LEU	3.3
1	A	196	TYR	3.2
1	B	137	ASN	3.1
1	B	85	LEU	3.1
1	A	81	GLN	3.1
2	F	354	VAL	3.0
1	A	55	THR	3.0
2	E	364	PHE	3.0
2	E	355	LEU	3.0
2	E	345	ILE	3.0
1	B	100	LEU	2.9
1	B	88	ILE	2.9
2	E	325	PHE	2.9
1	B	338	ASN	2.9
1	A	344	CYS	2.9
1	A	302	TRP	2.9
1	A	60	GLN	2.9
1	A	91	LEU	2.8
2	E	421	LEU	2.8
2	E	402	THR	2.8
1	A	298	VAL	2.8
1	B	94	LYS	2.7
2	E	359	THR	2.7
1	A	299	ASP	2.7
1	A	340	GLN	2.7
2	E	349	VAL	2.6
1	A	19	SER	2.6
1	A	301	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	361	CYS	2.6
1	B	92	THR	2.6
1	A	132	VAL	2.6
2	F	472	LEU	2.6
1	B	105	SER	2.6
2	E	357	ASN	2.6
1	B	367	ASP	2.5
1	B	21	THR	2.5
1	A	292	ASP	2.5
1	A	336	PRO	2.5
2	E	382	VAL	2.5
1	B	60	GLN	2.5
1	B	340	GLN	2.5
1	A	85	LEU	2.5
1	B	91	LEU	2.5
1	B	106	SER	2.5
2	F	419	CYS	2.4
1	B	19	SER	2.4
1	B	294	THR	2.4
2	F	360	PHE	2.4
2	E	351	ASP	2.4
1	A	341	LYS	2.4
1	B	213	ASP	2.4
2	E	415	ASP	2.3
2	E	354	VAL	2.3
1	B	553	LYS	2.3
1	A	362	THR	2.3
1	A	360	MET	2.3
1	B	429	GLN	2.3
2	F	350	ALA	2.3
2	F	356	TYR	2.3
1	A	337	GLY	2.3
2	F	445	HIS	2.2
1	A	427	ASP	2.2
1	A	56	GLU	2.2
1	B	28	PHE	2.2
2	E	350	ALA	2.2
1	A	103	ASN	2.2
1	A	319	GLY	2.2
2	E	383	TYR	2.2
1	B	214	GLY	2.2
1	A	93	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	139	GLN	2.1
1	B	111	ASP	2.1
1	B	572	ASN	2.1
1	A	423	LEU	2.1
1	B	366	MET	2.1
1	B	134	ASN	2.1
2	F	365	LYS	2.1
1	B	141	CYS	2.1
1	A	212	VAL	2.1
1	A	213	ASP	2.0
1	A	343	VAL	2.0
1	B	142	LEU	2.0
2	E	419	CYS	2.0
1	B	69	TRP	2.0
2	E	375	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NDG	B	617	15/15	0.21	8.80	79,88,90,90	0
5	ZN	A	901	1/1	0.21	1.54	85,85,85,85	0
6	CL	A	902	1/1	0.22	0.99	66,66,66,66	0
3	NDG	B	618	15/15	0.18	0.14	68,71,76,77	0
3	NDG	A	616	15/15	0.22	-0.30	109,109,110,111	0
3	NDG	B	616	15/15	0.26	-0.58	97,97,98,98	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ZN	B	901	1/1	0.24	-0.96	81,81,81,81	0
6	CL	B	902	1/1	0.15	-1.28	53,53,53,53	0
3	NDG	F	91	15/15	0.25	-	118,119,119,120	0
4	NAG	E	91	15/15	0.34	-	127,128,129,129	0

6.5 Other polymers

There are no such residues in this entry.