



# Full wwPDB X-ray Structure Validation Report

(i)

Mar 1, 2014 – 04:04 AM GMT

PDB ID : 3D0I

Title : Crystal structure of spike protein receptor-binding domain from the 2005-2006 SARS coronavirus civet strain complexed with human-civet chimeric receptor ACE2

Authors : Li, F.

Deposited on : 2008-05-01

Resolution : 2.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---

The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.15 2013

Xtriage (Phenix) : dev-1323

EDS : stable22639

Percentile statistics : 21963

Refmac : 5.8.0049

CCP4 : 6.3.0 (Settle)

Ideal geometry (proteins) : Engh & Huber (2001)


Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)

Validation Pipeline (wwPDB-VP) : stable22683

## 1 Overall quality at a glance (i)


The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1326 (2.90-2.90)
Ramachandran outliers	78287	1290 (2.90-2.90)
Sidechain outliers	78261	1292 (2.90-2.90)
RSRZ outliers	66119	1054 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $>=3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	NDG	B	618	-	X
5	ZN	A	901	-	X

## 2 Entry composition (i)

There are 7 unique types of molecules in this entry. The entry contains 12639 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensin-convertingenzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	597	Total	C	N	O	S	0	0	0
			4864	3110	803	922	29			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	597	Total	C	N	O	S	0	0	0
			4864	3110	803	922	29			

- Molecule 2 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	173	Total	C	N	O	S	0	0	0
			1395	905	229	255	6			


  

2	F	173	Total	C	N	O	S	0	0	0
			1395	905	229	255	6			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	479	ARG	ASN	CONFLICT	UNP P59594
E	480	GLY	ASP	CONFLICT	UNP P59594
E	487	SER	THR	CONFLICT	UNP P59594
F	479	ARG	ASN	CONFLICT	UNP P59594
F	480	GLY	ASP	CONFLICT	UNP P59594
F	487	SER	THR	CONFLICT	UNP P59594

- Molecule 3 is SUGAR (2-(ACETYLAMINO)-2-DEOXY-A-D-GLUCOPYRANOSE) (three-letter code: NDG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 15 8 1 6	0	0
3	B	1	Total C N O 15 8 1 6	0	0
3	B	1	Total C N O 15 8 1 6	0	0
3	B	1	Total C N O 15 8 1 6	0	0
3	B	1	Total C N O 15 8 1 6	0	0

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	E	1	Total C N O 15 8 1 6	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Zn 1 1	0	0
5	A	1	Total Zn 1 1	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Cl 1 1	0	0
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	21	Total O 21 21	0	0
7	F	6	Total O 6 6	0	0

### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Angiotensin-converting enzyme 2

Chain A: 




- Molecule 1: Angiotensin-converting enzyme 2

Chain B: 




- Molecule 2: Spike protein S1

Chain E:





- Molecule 2: Spike protein S1



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.42 Å    119.82 Å    109.77 Å 90.00°    95.50°    90.00°	Depositor
Resolution (Å)	36.47 – 2.90 36.47 – 2.89	Depositor EDS
% Data completeness (in resolution range)	89.9 (36.47-2.90) 89.3 (36.47-2.89)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.71 (at 2.90 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
$R$ , $R_{free}$	0.224 , 0.278 0.244 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	75.7	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 38.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$<  L  > = 0.52$ , $< L^2 > = 0.36$	Xtriage
Outliers	0 of 46219 reflections	Xtriage
$F_o$ , $F_c$ correlation	0.92	EDS
Total number of atoms	12639	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG, NDG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	1/5000 (0.0%)	0.60	0/6796
1	B	0.53	0/5000	0.62	0/6796
2	E	0.55	0/1440	0.61	0/1958
2	F	0.54	0/1440	0.61	0/1958
All	All	0.53	1/12880 (0.0%)	0.61	0/17508

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	300	GLN	CD-OE1	6.23	1.37	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4864	0	0	30	0
1	B	4864	0	0	24	0
2	E	1395	0	0	2	0
2	F	1395	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	15	0	0	1	0
3	B	60	0	0	2	0
4	E	15	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0
6	B	1	0	0	1	0
7	B	21	0	0	1	0
7	F	6	0	0	0	0
All	All	12639	0	0	62	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 5.

All (62) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:269:ASP:OD2	1:A:272:GLY:N	2.29	0.66
1:B:546:ASN:ND2	3:B:619:NDG:N2	2.47	0.62
1:B:541:LYS:NZ	1:B:541:LYS:CB	2.63	0.62
1:B:499:ASP:O	1:B:502:SER:CB	2.49	0.61
1:B:269:ASP:OD2	1:B:272:GLY:N	2.34	0.61
2:F:351:ASP:OD1	2:F:353:SER:OG	2.21	0.58
1:A:457:GLU:OE2	1:A:460:ARG:NH1	2.36	0.58
1:B:431:ASP:OD1	1:B:433:GLU:N	2.37	0.58
1:B:546:ASN:ND2	3:B:619:NDG:C1	2.67	0.57
1:A:397:ASN:OD1	1:A:399:GLY:N	2.39	0.56
1:B:489:GLU:OE1	1:B:489:GLU:N	2.38	0.56
1:A:560:LEU:N	1:A:560:LEU:CD1	2.71	0.54
1:A:100:LEU:O	1:A:102:GLN:N	2.42	0.52
1:B:211:GLY:N	1:B:216:ASP:OD2	2.43	0.52
1:B:177:ARG:NH2	1:B:497:TYR:O	2.44	0.50
1:A:90:ASN:ND2	3:A:616:NDG:O1L	2.46	0.49
1:B:183:TYR:OH	1:B:509:ASP:OD1	2.32	0.48
1:A:312:GLU:OE2	1:A:323:MET:N	2.46	0.48
1:B:169:ARG:NE	6:B:902:CL:CL	2.84	0.48
1:A:73:LEU:O	1:A:77:SER:N	2.46	0.48
1:A:118:THR:O	1:A:122:THR:OG1	2.32	0.47
1:B:401:HIS:CB	7:B:904:HOH:O	2.63	0.47
1:A:493:HIS:CD2	1:A:499:ASP:OD1	2.68	0.47
1:A:183:TYR:OH	1:A:509:ASP:OD1	2.33	0.46
1:B:166:GLU:OE1	1:B:493:HIS:CE1	2.69	0.46
1:A:99:ALA:O	1:A:102:GLN:NE2	2.49	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:436:ILE:O	1:A:437:ASN:C	2.54	0.46
1:A:60:GLN:O	1:A:64:ASN:N	2.49	0.46
1:B:514:ARG:O	1:B:518:ARG:CB	2.64	0.45
1:A:102:GLN:C	1:A:104:GLY:N	2.70	0.44
1:B:580:ASN:OD1	1:B:582:ARG:N	2.51	0.44
1:B:205:GLY:O	1:B:207:TYR:N	2.51	0.44
1:B:295:ASP:O	1:B:299:ASP:N	2.51	0.43
1:A:566:TRP:O	1:A:570:LEU:N	2.51	0.43
2:E:367:TYR:N	2:E:418:GLY:O	2.52	0.43
1:A:505:HIS:CE1	1:A:515:TYR:OH	2.71	0.43
1:B:163:TRP:O	1:B:164:ALA:C	2.57	0.43
1:A:117:ASN:O	1:A:121:ASN:ND2	2.52	0.42
1:A:174:LYS:CG	1:A:496:THR:O	2.67	0.42
2:F:367:TYR:CD1	2:F:367:TYR:N	2.87	0.42
1:B:203:TRP:C	1:B:205:GLY:N	2.73	0.42
1:A:168:TRP:CD1	1:A:502:SER:OG	2.73	0.42
2:F:348:CYS:O	2:F:375:ASN:O	2.37	0.42
1:A:318:VAL:CA	1:A:547:SER:O	2.68	0.42
1:A:489:GLU:N	1:A:489:GLU:OE2	2.52	0.42
2:F:463:ASP:O	2:F:464:GLY:C	2.58	0.41
1:A:494:ASP:OD1	1:A:494:ASP:N	2.53	0.41
1:B:73:LEU:O	1:B:77:SER:N	2.52	0.41
1:A:578:ASN:OD1	1:A:579:MET:N	2.52	0.41
1:A:452:PHE:CD1	1:A:452:PHE:C	2.94	0.41
1:A:476:LYS:NZ	1:A:479:GLU:OE2	2.54	0.41
1:B:589:GLU:O	1:B:590:PRO:C	2.57	0.41
2:F:324:PRO:CD	2:F:348:CYS:SG	3.09	0.41
1:B:203:TRP:O	1:B:205:GLY:N	2.54	0.40
1:B:580:ASN:OD1	1:B:581:VAL:N	2.53	0.40
1:A:203:TRP:C	1:A:205:GLY:N	2.75	0.40
1:A:253:PRO:O	1:A:254:SER:CB	2.69	0.40
1:B:117:ASN:O	1:B:121:ASN:ND2	2.54	0.40
2:F:423:TRP:CD1	2:F:423:TRP:N	2.89	0.40
1:A:76:GLN:O	1:A:80:ALA:N	2.54	0.40
1:A:524:GLN:NE2	1:A:580:ASN:N	2.70	0.40
2:E:364:PHE:CD1	2:E:364:PHE:O	2.74	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	595/597 (100%)	524 (88%)	52 (9%)	19 (3%)	6 25
1	B	595/597 (100%)	528 (89%)	50 (8%)	17 (3%)	7 28
2	E	169/179 (94%)	138 (82%)	21 (12%)	10 (6%)	2 7
2	F	169/179 (94%)	141 (83%)	17 (10%)	11 (6%)	2 5
All	All	1528/1552 (98%)	1331 (87%)	140 (9%)	57 (4%)	5 20

All (57) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	146	PRO
1	A	212	VAL
1	A	289	PRO
1	A	340	GLN
1	A	614	ALA
1	B	146	PRO
1	B	147	GLY
1	B	212	VAL
1	B	290	ASN
1	B	340	GLN
1	B	428	PHE
2	E	370	SER
2	E	402	THR
2	E	415	ASP
2	F	370	SER
2	F	401	GLN
2	F	402	THR
2	F	415	ASP
2	F	416	PHE
2	F	465	LYS
1	A	54	ILE
1	A	147	GLY
1	A	427	ASP
1	A	428	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	432	ASN
1	B	54	ILE
1	B	390	PHE
1	B	427	ASP
1	B	536	GLU
2	E	369	VAL
2	E	401	GLN
2	E	416	PHE
2	F	369	VAL
1	A	206	ASP
1	A	504	PHE
1	B	64	ASN
1	B	213	ASP
2	F	368	GLY
1	A	164	ALA
1	A	213	ASP
1	A	508	ASN
1	B	206	ASP
2	E	365	LYS
2	E	465	LYS
1	A	82	MET
1	A	254	SER
1	A	339	VAL
1	B	92	THR
1	B	108	LEU
1	B	163	TRP
1	B	164	ALA
2	F	482	GLY
1	A	364	VAL
2	E	404	VAL
2	E	482	GLY
2	F	404	VAL
2	F	464	GLY

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	527/527 (100%)	491 (93%)	36 (7%)	22 55
1	B	527/527 (100%)	484 (92%)	43 (8%)	17 44
2	E	150/156 (96%)	137 (91%)	13 (9%)	15 41
2	F	150/156 (96%)	139 (93%)	11 (7%)	20 51
All	All	1354/1366 (99%)	1251 (92%)	103 (8%)	19 48

All (103) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ASN
1	A	89	GLN
1	A	96	GLN
1	A	102	GLN
1	A	114	LYS
1	A	143	LEU
1	A	149	ASN
1	A	172	VAL
1	A	212	VAL
1	A	221	GLN
1	A	250	ASN
1	A	287	GLN
1	A	334	THR
1	A	365	THR
1	A	371	THR
1	A	381	TYR
1	A	385	TYR
1	A	401	HIS
1	A	407	ILE
1	A	410	LEU
1	A	418	LEU
1	A	444	LEU
1	A	449	THR
1	A	460	ARG
1	A	476	LYS
1	A	479	GLU
1	A	508	ASN
1	A	511	SER
1	A	526	GLN
1	A	546	ASN
1	A	549	GLU
1	A	557	MET
1	A	558	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	563	SER
1	A	573	VAL
1	A	600	LYS
1	B	86	GLN
1	B	89	GLN
1	B	96	GLN
1	B	102	GLN
1	B	114	LYS
1	B	129	THR
1	B	139	GLN
1	B	156	LEU
1	B	212	VAL
1	B	221	GLN
1	B	250	ASN
1	B	254	SER
1	B	287	GLN
1	B	334	THR
1	B	357	ARG
1	B	368	ASP
1	B	371	THR
1	B	381	TYR
1	B	385	TYR
1	B	407	ILE
1	B	410	LEU
1	B	418	LEU
1	B	429	GLN
1	B	439	LEU
1	B	444	LEU
1	B	455	MET
1	B	456	LEU
1	B	458	LYS
1	B	471	ASP
1	B	473	TRP
1	B	488	VAL
1	B	491	VAL
1	B	496	THR
1	B	526	GLN
1	B	529	LEU
1	B	541	LYS
1	B	557	MET
1	B	560	LEU
1	B	570	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	593	THR
1	B	595	LEU
1	B	602	SER
1	B	611	SER
2	E	333	LYS
2	E	337	VAL
2	E	359	THR
2	E	364	PHE
2	E	369	VAL
2	E	411	LYS
2	E	412	LEU
2	E	416	PHE
2	E	417	MET
2	E	423	TRP
2	E	439	LYS
2	E	475	TYR
2	E	479	ARG
2	F	333	LYS
2	F	337	VAL
2	F	364	PHE
2	F	369	VAL
2	F	395	ARG
2	F	411	LYS
2	F	416	PHE
2	F	417	MET
2	F	423	TRP
2	F	468	THR
2	F	475	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NDG	A	616	-	15,15,15	0.48	0	21,21,21	0.61	0
3	NDG	B	616	-	15,15,15	0.62	0	21,21,21	0.60	0
3	NDG	B	617	-	15,15,15	0.54	0	21,21,21	1.24	2 (9%)
3	NDG	B	618	-	15,15,15	0.73	0	21,21,21	1.05	1 (4%)
3	NDG	B	619	-	15,15,15	0.63	0	21,21,21	1.08	2 (9%)
4	NAG	E	91	-	15,15,15	0.75	0	21,21,21	1.32	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDG	A	616	-	-	1/6/26/26	0/1/1/1
3	NDG	B	616	-	-	0/6/26/26	0/1/1/1
3	NDG	B	617	-	-	0/6/26/26	0/1/1/1
3	NDG	B	618	-	-	1/6/26/26	0/1/1/1
3	NDG	B	619	-	-	0/6/26/26	0/1/1/1
4	NAG	E	91	-	-	0/6/26/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	E	91	NAG	C1-O5-C5	3.39	119.47	113.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	617	NDG	C1-C2-N2	3.26	114.63	110.85
4	E	91	NAG	O5-C1-C2	2.95	112.60	109.61
3	B	617	NDG	C4-C3-C2	2.79	114.41	110.44
3	B	618	NDG	O-C1-C2	2.75	112.40	109.61
3	B	619	NDG	O-C5-C4	2.23	113.89	109.76
3	B	619	NDG	C1-O-C5	2.17	117.29	113.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	616	NDG	C1-C2-N2-C7
3	B	618	NDG	O7-C7-N2-C2

There are no ring outliers.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	597/597 (100%)	0.41	45 (7%) 14 17	47, 80, 130, 154	0
1	B	597/597 (100%)	0.25	31 (5%) 26 32	38, 74, 130, 156	0
2	E	173/179 (96%)	0.37	11 (6%) 19 23	56, 89, 132, 134	0
2	F	173/179 (96%)	0.36	12 (6%) 17 20	58, 90, 131, 134	0
All	All	1540/1552 (99%)	0.34	99 (6%) 18 23	38, 80, 131, 156	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	290	ASN	7.0
1	A	342	ALA	6.9
1	B	615	ASP	6.7
1	A	341	LYS	6.6
1	A	302	TRP	5.9
1	A	105	SER	5.6
1	A	303	ASP	5.1
1	A	298	VAL	5.0
1	A	106	SER	4.5
2	E	501	PHE	4.5
1	B	89	GLN	4.4
1	A	54	ILE	4.3
1	A	339	VAL	4.3
2	F	352	TYR	4.2
1	B	105	SER	4.0
1	B	85	LEU	4.0
2	F	360	PHE	4.0
1	B	339	VAL	3.9
2	F	345	ILE	3.8
2	E	382	VAL	3.8
1	A	343	VAL	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	340	GLN	3.5
1	A	60	GLN	3.5
2	E	325	PHE	3.5
2	E	345	ILE	3.5
1	A	132	VAL	3.4
1	B	91	LEU	3.4
1	A	62	MET	3.4
1	A	111	ASP	3.4
1	A	299	ASP	3.3
1	B	55	THR	3.3
1	B	341	LYS	3.2
1	A	108	LEU	3.2
1	A	107	VAL	3.2
2	E	421	LEU	3.1
1	B	136	ASP	3.1
2	F	470	PRO	3.1
1	B	54	ILE	3.0
1	A	338	ASN	3.0
1	A	424	LEU	2.9
1	B	84	PRO	2.8
1	B	83	TYR	2.8
1	B	82	MET	2.7
2	E	364	PHE	2.7
1	A	55	THR	2.7
1	A	142	LEU	2.6
1	A	291	ILE	2.6
1	A	363	LYS	2.6
2	E	344	LYS	2.6
1	B	111	ASP	2.6
1	A	328	TRP	2.5
1	A	56	GLU	2.5
1	A	138	PRO	2.5
1	A	320	LEU	2.5
2	E	355	LEU	2.5
1	B	88	ILE	2.5
1	B	21	THR	2.5
1	B	429	GLN	2.5
1	A	19	SER	2.5
1	B	86	GLN	2.5
1	B	137	ASN	2.5
2	F	472	LEU	2.4
1	A	426	PRO	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	114	LYS	2.4
1	B	403	ALA	2.4
1	A	330	ASN	2.3
1	A	59	VAL	2.3
1	A	50	TYR	2.3
2	E	357	ASN	2.3
1	B	22	GLU	2.2
1	B	61	ASN	2.2
1	A	336	PRO	2.2
1	A	86	GLN	2.2
1	B	614	ALA	2.2
2	F	445	HIS	2.2
1	A	344	CYS	2.2
2	F	325	PHE	2.2
2	F	475	TYR	2.2
1	A	331	SER	2.2
2	F	421	LEU	2.2
2	E	359	THR	2.2
1	A	51	ASN	2.1
1	A	307	ILE	2.1
1	B	394	ASN	2.1
2	F	361	PHE	2.1
2	E	472	LEU	2.1
2	F	382	VAL	2.1
1	B	213	ASP	2.1
1	B	28	PHE	2.1
1	A	104	GLY	2.1
1	B	135	PRO	2.1
1	A	119	ILE	2.1
1	A	573	VAL	2.1
1	B	186	LEU	2.0
1	A	83	TYR	2.0
2	F	355	LEU	2.0
1	A	361	CYS	2.0
1	B	214	GLY	2.0
1	B	94	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	ZN	A	901	1/1	0.43	4.40	123,123,123,123	0
3	NDG	B	618	15/15	0.26	2.02	111,113,114,114	0
3	NDG	B	616	15/15	0.27	0.74	123,125,125,126	0
3	NDG	B	619	15/15	0.18	0.14	72,76,81,81	0
5	ZN	B	901	1/1	0.26	0.11	131,131,131,131	0
6	CL	A	902	1/1	0.19	-0.30	84,84,84,84	0
3	NDG	A	616	15/15	0.13	-1.27	109,113,114,114	0
6	CL	B	902	1/1	0.17	-1.68	73,73,73,73	0
3	NDG	B	617	15/15	0.27	-	126,128,129,129	0
4	NAG	E	91	15/15	0.29	-	137,140,141,142	0

## 6.5 Other polymers (i)

There are no such residues in this entry.