



# Full wwPDB X-ray Structure Validation Report

Nov 13, 2014 – 05:31 PM EST

PDB ID : 4D05  
Title : Structure and activity of a minimal-type ATP-dependent DNA ligase from a psychrotolerant bacterium  
Authors : Williamson, A.; Rothweiler, U.; Leiros, H.-K.S.  
Deposited on : 2014-04-24  
Resolution : 1.65 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.16 November 2013  
Xtriage (Phenix) : dev-1439  
EDS : stable24195  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.1.3  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable24195

# 1 Overall quality at a glance

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1404 (1.68-1.64)
Clashscore	79885	1001 (1.66-1.66)
Ramachandran outliers	78287	1581 (1.68-1.64)
Sidechain outliers	78261	1580 (1.68-1.64)
RSRZ outliers	66119	1404 (1.68-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	258	<div><div></div><div></div></div>
1	B	258	<div><div></div><div></div></div>

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	SO4	A	1262	-	X

## 2 Entry composition i


There are 5 unique types of molecules in this entry. The entry contains 9090 atoms, of which 4181 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-DEPENDENT DNA LIGASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	258	Total	C	H	N	O	Se	0	6	0
			4243	1357	2118	361	399	8			
1	B	249	Total	C	H	N	O	Se	101	5	0
			4081	1304	2039	346	383	9			

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula:  $C_{10}H_{14}N_5O_7P$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total 34	C 10	H 12	N 5	O 6	P 1	0	0
3	B	1	Total 34	C 10	H 12	N 5	O 6	P 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	471	Total 471	O 471	0	0
5	B	181	Total 181	O 181	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: ATP-DEPENDENT DNA LIGASE

Chain A: 



#### • Molecule 1: ATP-DEPENDENT DNA LIGASE

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.43 Å   43.98 Å   89.63 Å 90.00°   105.92°   90.00°	Depositor
Resolution (Å)	24.61 – 1.65 43.61 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.6 (24.61-1.65) 94.2 (43.61-1.65)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.10 (at 1.65 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.145   ,   0.192 0.145   ,   0.192	Depositor DCC
$R_{free}$ test set	2298 reflections (3.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.0	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 39.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 80780 reflections (0.001%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9090	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.67	0/2165	0.74	1/2917 (0.0%)
1	B	0.50	0/2077	0.66	2/2794 (0.1%)
All	All	0.59	0/4242	0.71	3/5711 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	155[A]	MSE	CA-CB-CG	-6.90	101.57	113.30
1	B	155[B]	MSE	CA-CB-CG	-6.90	101.57	113.30
1	A	109	ARG	NE-CZ-NH1	5.55	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2125	2118	2111	18	0
1	B	2042	2039	2030	42	2
2	A	25	0	0	2	0
2	B	20	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	22	12	12	1	0
3	B	22	12	12	0	0
4	B	1	0	0	0	0
5	A	471	0	0	11	5
5	B	181	0	0	5	2
All	All	4909	4181	4165	62	7

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (62) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:179[A]:ASP:OD2	5:A:2357:HOH:O	1.75	1.04
2:A:1261:SO4:O4	5:A:2341:HOH:O	1.77	0.99
1:B:109:ARG:NH2	1:B:161:ALA:O	2.10	0.84
1:B:98:ASP:N	1:B:113:MSE:HE1	1.93	0.83
1:A:30:ARG:NH2	1:A:62[B]:ASP:OD2	2.17	0.73
1:B:9:THR:HG23	1:B:168:SER:N	2.06	0.70
1:A:152[B]:GLU:OE1	5:A:2192:HOH:O	2.09	0.69
1:B:219:ASP:OD1	5:B:2155:HOH:O	2.12	0.67
1:B:98:ASP:C	1:B:113:MSE:HE3	2.17	0.64
1:A:167:ARG:NH2	5:A:2341:HOH:O	2.28	0.59
1:B:239:LYS:HD3	1:B:243[B]:ASN:OD1	2.01	0.59
1:B:104:VAL:HG12	1:B:105:SER:H	1.68	0.59
1:B:104:VAL:HG12	1:B:105:SER:N	2.19	0.57
1:B:154:LEU:O	1:B:155[B]:MSE:HG2	2.03	0.57
1:B:106:PHE:HA	1:B:109:ARG:HB2	1.85	0.57
1:B:16:ASP:CG	1:B:18:THR:HG22	2.25	0.56
2:A:1262:SO4:O1	5:A:2164:HOH:O	2.12	0.54
1:A:257:TYR:C	5:A:2434:HOH:O	2.45	0.54
1:B:168:SER:O	1:B:169:ARG:HB2	2.07	0.53
1:B:132:LYS:HD3	1:B:159:GLN:HE21	1.74	0.53
1:B:129:GLU:OE1	5:B:2097:HOH:O	2.19	0.52
1:A:164:GLN:NE2	5:A:2330:HOH:O	2.41	0.52
1:A:179[A]:ASP:CG	5:A:2357:HOH:O	2.36	0.52
1:B:135:ASN:OD1	1:B:137:VAL:CG1	2.59	0.51
1:B:16:ASP:OD2	1:B:18:THR:HG22	2.10	0.51
1:B:154:LEU:C	1:B:155[B]:MSE:HG2	2.31	0.51
1:B:207:GLU:OE1	1:B:243[B]:ASN:ND2	2.35	0.51
1:B:239:LYS:CD	1:B:243[B]:ASN:OD1	2.59	0.50
1:B:144:ASN:O	1:B:148[A]:MSE:HG3	2.12	0.50
1:B:150:LYS:N	1:B:150:LYS:HD3	2.27	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:135:ASN:OD1	1:B:137:VAL:HG12	2.13	0.49
1:A:114:GLN:HG2	5:A:2269:HOH:O	2.12	0.49
1:B:98:ASP:C	1:B:113:MSE:CE	2.81	0.49
1:A:235:ARG:O	1:A:248:ALA:HA	2.14	0.47
1:A:62[B]:ASP:CG	5:A:2099:HOH:O	2.51	0.47
1:A:166:LYS:HG3	1:A:167:ARG:O	2.14	0.47
1:B:239:LYS:NZ	5:B:2165:HOH:O	2.47	0.47
1:B:205:ASN:HB2	5:B:2114:HOH:O	2.15	0.46
1:A:155:MSE:HG3	3:A:1263:AMP:C6	2.50	0.46
1:B:19:GLN:HA	1:B:159:GLN:NE2	2.31	0.46
1:B:15:ILE:CG2	1:B:16:ASP:N	2.78	0.46
1:B:132:LYS:HD3	1:B:159:GLN:NE2	2.31	0.45
1:A:102:SER:HB3	1:A:109:ARG:HD2	1.98	0.45
1:A:195:GLU:HG3	5:A:2379:HOH:O	2.17	0.45
1:B:104:VAL:CG1	1:B:105:SER:H	2.30	0.44
1:B:150:LYS:CD	1:B:150:LYS:N	2.80	0.44
1:B:15:ILE:HD13	1:B:15:ILE:HA	1.89	0.44
1:B:70:GLN:OE1	5:B:2070:HOH:O	2.21	0.44
1:B:27:ASP:HB2	1:B:152:GLU:HG3	2.01	0.43
1:A:179[B]:ASP:OD1	1:A:179[B]:ASP:O	2.36	0.42
1:A:62[B]:ASP:OD1	1:A:100:PRO:HG3	2.19	0.42
1:B:154:LEU:O	1:B:155[A]:MSE:HG2	2.19	0.42
1:A:20:TYR:CE2	1:A:172:MSE:HE2	2.55	0.42
1:B:16:ASP:OD2	1:B:18:THR:CG2	2.68	0.41
1:B:133:ILE:HG23	1:B:134:PRO:HD2	2.02	0.41
1:B:104:VAL:CG1	1:B:105:SER:N	2.83	0.41
1:B:34:ASN:HD21	1:B:37:GLN:CG	2.34	0.41
1:B:9:THR:O	1:B:172:MSE:HA	2.20	0.41
1:B:59:THR:HG22	1:B:60:ALA:N	2.35	0.41
1:B:98:ASP:CA	1:B:113:MSE:HE1	2.50	0.41
1:A:179[B]:ASP:C	1:A:179[B]:ASP:OD1	2.59	0.40
1:B:133:ILE:CG2	1:B:134:PRO:HD2	2.52	0.40

All (7) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:2468:HOH:O	5:B:2080:HOH:O[3.445]	2.02	0.18
5:A:2410:HOH:O	5:B:2067:HOH:O[3.445]	2.08	0.12
5:A:2004:HOH:O	5:A:2057:HOH:O[1.565]	2.11	0.09
5:A:2134:HOH:O	5:A:2393:HOH:O[4.546]	2.18	0.02
5:A:2091:HOH:O	5:A:2318:HOH:O[1.565]	2.19	0.01

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:148[A]:MSE:HE3	1:B:235:ARG:HH22[2.656]	2.11	-0.51
1:B:148[A]:MSE:HE3	1:B:235:ARG:NH2[2.656]	2.16	-0.56

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/258 (102%)	255 (97%)	7 (3%)	0	100	100
1	B	250/258 (97%)	241 (96%)	9 (4%)	0	100	100
All	All	512/516 (99%)	496 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/219 (106%)	231 (100%)	1 (0%)	95	91
1	B	223/219 (102%)	222 (100%)	1 (0%)	95	91
All	All	455/438 (104%)	453 (100%)	2 (0%)	94	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	GLN
1	B	25	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	164	GLN
1	B	159	GLN
1	B	241	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1258	-	4,4,4	0.53	0	6,6,6	0.34	0
2	SO4	A	1259	-	4,4,4	0.41	0	6,6,6	0.27	0
2	SO4	A	1260	-	4,4,4	0.24	0	6,6,6	0.23	0
2	SO4	A	1261	-	4,4,4	0.14	0	6,6,6	0.30	0
2	SO4	A	1262	-	4,4,4	1.07	0	6,6,6	0.66	0
3	AMP	A	1263	1	22,24,25	1.52	4 (18%)	32,35,38	2.54	15 (46%)
2	SO4	B	1256	-	4,4,4	0.15	0	6,6,6	0.14	0
2	SO4	B	1257	-	4,4,4	0.12	0	6,6,6	0.22	0
2	SO4	B	1258	-	4,4,4	0.07	0	6,6,6	0.14	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SO4	B	1259	-	4,4,4	0.16	0	6,6,6	0.33	0
3	AMP	B	1260	-	22,24,25	1.51	3 (13%)	32,35,38	2.82	11 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1258	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1259	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1260	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1261	-	-	0/0/0/0	0/0/0/0
2	SO4	A	1262	-	-	0/0/0/0	0/0/0/0
3	AMP	A	1263	1	-	0/8/25/26	0/3/3/3
2	SO4	B	1256	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1257	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1258	-	-	0/0/0/0	0/0/0/0
2	SO4	B	1259	-	-	0/0/0/0	0/0/0/0
3	AMP	B	1260	-	-	0/8/25/26	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1260	AMP	P-O1P	4.75	1.52	1.46
3	A	1263	AMP	P-O1P	3.47	1.50	1.46
3	A	1263	AMP	C4-N9	-3.35	1.32	1.37
3	A	1263	AMP	C5-C4	2.95	1.47	1.40
3	B	1260	AMP	C5-C4	2.87	1.47	1.40
3	B	1260	AMP	C4-N9	-2.63	1.33	1.37
3	A	1263	AMP	O5'-C5'	-2.38	1.41	1.44

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1260	AMP	C5-C4-N3	-8.24	117.94	125.98
3	B	1260	AMP	N3-C4-N9	7.20	137.75	125.39
3	A	1263	AMP	N3-C2-N1	-6.42	123.24	128.89
3	B	1260	AMP	N3-C2-N1	-6.28	123.37	128.89
3	A	1263	AMP	C5-C4-N3	-5.43	120.68	125.98
3	A	1263	AMP	N3-C4-N9	4.61	133.30	125.39
3	B	1260	AMP	C3'-C2'-C1'	4.26	107.59	100.92

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1263	AMP	C3'-C2'-C1'	4.06	107.28	100.92
3	B	1260	AMP	C8-N9-C1'	-3.81	118.98	126.15
3	A	1263	AMP	O2'-C2'-C3'	-3.76	99.68	111.83
3	A	1263	AMP	C4'-O4'-C1'	3.62	113.70	109.72
3	B	1260	AMP	C8-N9-C4	3.00	109.40	106.96
3	A	1263	AMP	C1'-N9-C4	-2.93	121.57	126.64
3	A	1263	AMP	C2-N1-C6	2.91	123.94	118.76
3	B	1260	AMP	C1'-N9-C4	2.88	131.62	126.64
3	B	1260	AMP	C2-N3-C4	2.81	121.37	113.27
3	A	1263	AMP	O2'-C2'-C1'	2.55	119.54	111.49
3	B	1260	AMP	N6-C6-N1	2.46	124.33	119.11
3	A	1263	AMP	C8-N9-C1'	2.44	130.74	126.15
3	B	1260	AMP	O2'-C2'-C1'	-2.33	104.11	111.49
3	A	1263	AMP	O3'-C3'-C4'	-2.31	104.27	111.07
3	A	1263	AMP	P-O5'-C5'	-2.25	114.11	122.98
3	A	1263	AMP	O4'-C1'-N9	-2.21	103.29	108.10
3	A	1263	AMP	N6-C6-N1	2.20	123.79	119.11
3	A	1263	AMP	C2-N3-C4	2.03	119.11	113.27
3	B	1260	AMP	C5-C4-N9	-2.01	104.30	107.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	258/258 (100%)	-0.37	0 100 100	3, 8, 21, 37	1 (0%)
1	B	249/258 (96%)	0.53	40 (16%) 2 1	5, 28, 50, 55	20 (8%)
All	All	507/516 (98%)	0.07	40 (7%) 13 11	3, 16, 47, 55	21 (4%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	ILE	7.3
1	B	5	ILE	5.9
1	B	17	VAL	4.5
1	B	103	THR	4.4
1	B	18	THR	4.1
1	B	44	ASN	4.0
1	B	162	LEU	3.5
1	B	14	ASP	3.3
1	B	104	VAL	3.2
1	B	168	SER	3.2
1	B	45	ILE	3.0
1	B	13	GLN	3.0
1	B	101	LYS	3.0
1	B	7	LEU	2.9
1	B	60	ALA	2.8
1	B	36	HIS	2.8
1	B	58	THR	2.8
1	B	6	GLN	2.8
1	B	41	LYS	2.6
1	B	241	ASN	2.6
1	B	164	GLN	2.6
1	B	43	GLY	2.6
1	B	165	THR	2.6
1	B	133	ILE	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	138	ALA	2.4
1	B	34	ASN	2.3
1	B	11	TYR	2.3
1	B	32	TYR	2.3
1	B	102	SER	2.3
1	B	159	GLN	2.3
1	B	137	VAL	2.3
1	B	106	PHE	2.3
1	B	134	PRO	2.3
1	B	37	GLN	2.2
1	B	16	ASP	2.2
1	B	158	HIS	2.1
1	B	59	THR	2.1
1	B	161	ALA	2.0
1	B	163	TYR	2.0
1	B	35	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	SO4	A	1262	5/5	0.21	8.04	17,23,33,35	5
2	SO4	B	1257	5/5	0.12	1.10	51,52,54,56	0
2	SO4	A	1260	5/5	0.07	1.09	19,23,23,26	0
2	SO4	A	1261	5/5	0.14	0.86	36,36,40,41	5
2	SO4	B	1258	5/5	0.27	0.33	49,50,53,54	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	AMP	A	1263	22/23	0.09	0.29	4,15,25,28	0
2	SO4	B	1259	5/5	0.11	-0.13	39,40,42,43	5
3	AMP	B	1260	22/23	0.13	-0.16	24,38,52,54	0
4	MG	B	1261	1/1	0.13	-0.26	41,41,41,41	0
2	SO4	A	1258	5/5	0.08	-0.64	13,14,21,25	5
2	SO4	A	1259	5/5	0.06	-0.86	22,22,24,24	0
2	SO4	B	1256	5/5	0.07	-1.19	32,35,36,39	0

## 6.5 Other polymers

There are no such residues in this entry.