



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:33 AM GMT

PDB ID : 3D2R
Title : Crystal structure of pyruvate dehydrogenase kinase isoform 4 in complex with ADP
Authors : Kato, M.; Wynn, R.M.; Chuang, L.C.; Tso, S.-C.; Li, J.; Chuang, D.T.
Deposited on : 2008-05-08
Resolution : 2.03 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	6003 (2.04-2.00)
Clashscore	79885	7467 (2.04-2.00)
Ramachandran outliers	78287	7370 (2.04-2.00)
Sidechain outliers	78261	7368 (2.04-2.00)
RSRZ outliers	66119	6006 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	394	
1	B	394	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
4	GOL	B	502	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6111 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	0	0
			2853	1830	477	531	15			
1	B	362	Total	C	N	O	S	0	0	0
			2903	1861	487	540	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	ALA	-	EXPRESSION TAG	UNP Q16654
A	19	SER	-	EXPRESSION TAG	UNP Q16654
B	18	ALA	-	EXPRESSION TAG	UNP Q16654
B	19	SER	-	EXPRESSION TAG	UNP Q16654

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total 125	O 125	0	0
5	B	162	Total 162	O 162	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 4

Chain A:



- Molecule 1: [Pyruvate dehydrogenase [lipoamide]] kinase isozyme 4

Chain B:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.59Å 69.37Å 82.33Å 90.00° 99.26° 90.00°	Depositor
Resolution (Å)	50.00 – 2.03 49.16 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.03) 99.3 (49.16-2.03)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.24 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, R_{free}	0.183 , 0.230 0.183 , 0.230	Depositor DCC
R_{free} test set	2564 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.315	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 59.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 50489 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6111	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	2/2918 (0.1%)	0.87	2/3952 (0.1%)
1	B	0.97	0/2972	0.90	3/4032 (0.1%)
All	All	0.93	2/5890 (0.0%)	0.89	5/7984 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	206	ASP	CB-CG	-5.93	1.39	1.51
1	A	49	CYS	CB-SG	5.20	1.91	1.82

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	148	ASP	CB-CG-OD1	-8.50	110.65	118.30
1	A	286	ASP	CB-CG-OD2	-5.67	113.20	118.30
1	B	286	ASP	CB-CG-OD2	-5.43	113.41	118.30
1	A	374	LYS	CD-CE-NZ	-5.27	99.59	111.70
1	B	37	MET	CG-SD-CE	5.16	108.46	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2853	0	2822	40	0
1	B	2903	0	2875	51	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	27	0	12	0	0
3	B	27	0	12	0	0
4	A	6	0	8	7	0
4	B	6	0	8	5	0
5	A	125	0	0	0	0
5	B	162	0	0	9	0
All	All	6111	0	5737	86	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 7.

All (86) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:397:ILE:O	1:A:397:ILE:HG13	1.59	1.01
1:A:46:GLU:O	1:A:47:ASN:HB2	1.68	0.94
1:A:175:GLN:HE22	1:A:193:SER:H	1.06	0.93
1:A:315:PRO:O	1:A:316:THR:HB	1.66	0.91
1:A:396:CYS:HB2	5:B:590:HOH:O	1.74	0.86
1:A:342:LYS:HD3	4:A:502:GOL:H32	1.61	0.83
1:B:374:LYS:HD2	5:B:623:HOH:O	1.85	0.77
1:B:268:GLU:HA	5:B:659:HOH:O	1.84	0.76
1:A:175:GLN:NE2	1:A:193:SER:H	1.85	0.74
1:B:342:LYS:HD3	4:B:502:GOL:H31	1.69	0.73
1:A:342:LYS:HD3	4:A:502:GOL:C3	2.23	0.68
1:B:74:ILE:HD11	1:B:383:PHE:CD1	2.29	0.68
1:B:342:LYS:NZ	4:B:502:GOL:H32	2.09	0.67
1:B:144:ALA:O	1:B:146:THR:N	2.29	0.66
1:A:175:GLN:HE22	1:A:193:SER:N	1.89	0.64
1:B:32:PRO:HD2	1:B:377:VAL:HG22	1.80	0.64
1:A:235:PRO:O	1:A:236:ASP:HB2	1.97	0.64
1:A:315:PRO:O	1:A:316:THR:CB	2.42	0.62
1:A:342:LYS:CD	4:A:502:GOL:H32	2.28	0.62
1:B:124:ARG:HD3	1:B:174:ASN:HD21	1.63	0.62
1:B:152:ASN:HD22	1:B:155:LEU:H	1.49	0.60
1:B:44:GLY:HA3	1:B:179:ILE:HD13	1.85	0.58
1:B:342:LYS:HD3	4:B:502:GOL:C3	2.33	0.58
1:B:124:ARG:CD	1:B:174:ASN:HD21	2.17	0.58
1:B:138:ILE:CD1	1:B:155:LEU:HD21	2.33	0.58
1:B:58:ARG:O	1:B:103:HIS:HE1	1.86	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:288:THR:HG21	1:B:353:LEU:HD22	1.86	0.57
1:A:342:LYS:HE2	4:A:502:GOL:H12	1.87	0.56
1:A:288:THR:HG21	1:B:353:LEU:CD2	2.36	0.56
1:B:175:GLN:HE22	1:B:193:SER:H	1.53	0.55
1:B:288:THR:CG2	1:B:361:ILE:HD11	2.37	0.55
1:A:342:LYS:NZ	4:A:502:GOL:H32	2.22	0.54
1:B:148:ASP:O	1:B:152:ASN:HB3	2.08	0.54
1:A:288:THR:CG2	1:B:353:LEU:HD22	2.38	0.54
1:A:136:GLN:O	1:A:140:GLU:HG2	2.08	0.54
1:B:269:ASN:HB2	5:B:653:HOH:O	2.08	0.53
1:A:396:CYS:SG	1:B:377:VAL:CG2	2.97	0.53
1:A:374:LYS:HD3	4:A:502:GOL:H11	1.90	0.53
1:B:44:GLY:CA	1:B:179:ILE:HD13	2.39	0.52
1:A:300:LEU:HD23	5:B:612:HOH:O	2.09	0.52
1:B:41:LEU:HD22	1:B:191:ILE:HD12	1.90	0.52
1:B:124:ARG:HD3	1:B:174:ASN:ND2	2.24	0.52
1:A:361:ILE:HD11	1:B:353:LEU:CD2	2.40	0.52
1:B:223:PRO:HD2	1:B:260:MET:HG2	1.93	0.51
1:B:374:LYS:NZ	5:B:663:HOH:O	2.43	0.50
1:B:148:ASP:O	1:B:152:ASN:CB	2.59	0.50
1:A:46:GLU:O	1:A:47:ASN:CB	2.48	0.50
1:B:374:LYS:CD	5:B:623:HOH:O	2.50	0.49
1:A:74:ILE:O	1:A:74:ILE:HG22	2.13	0.49
1:B:138:ILE:HD12	1:B:155:LEU:HD21	1.95	0.49
1:B:74:ILE:O	1:B:74:ILE:HG23	2.13	0.48
1:A:37:MET:HE2	1:A:371:SER:CB	2.42	0.48
1:B:305:ARG:HG2	1:B:311:TYR:CD1	2.49	0.48
1:A:23:GLU:HB3	1:A:96:LEU:HD21	1.95	0.48
1:B:317:PRO:HD3	1:B:329:GLY:HA3	1.96	0.47
1:B:74:ILE:HD11	1:B:383:PHE:HD1	1.75	0.47
1:B:305:ARG:HB2	1:B:305:ARG:HE	1.56	0.46
1:B:374:LYS:HE3	5:B:623:HOH:O	2.15	0.46
1:B:288:THR:HG21	1:B:361:ILE:HD11	1.99	0.45
1:A:198:CYS:O	1:A:240:HIS:HA	2.16	0.45
1:B:233:LYS:HE2	1:B:283:GLY:O	2.16	0.45
1:B:197:ASN:O	1:B:240:HIS:HD2	1.99	0.45
1:A:51:ARG:CG	1:A:52:THR:N	2.80	0.45
1:B:342:LYS:HZ3	4:B:502:GOL:H32	1.80	0.45
1:B:74:ILE:O	1:B:74:ILE:CG2	2.63	0.45
1:A:124:ARG:HD3	1:A:174:ASN:HD21	1.82	0.45
1:B:269:ASN:CB	5:B:653:HOH:O	2.63	0.44
1:A:298:VAL:CG1	1:A:302:ILE:HG22	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:309:TYR:CE2	1:A:335:PRO:HB2	2.53	0.43
1:B:210:CYS:HA	1:B:213:MET:HE2	1.99	0.43
1:B:175:GLN:NE2	1:B:193:SER:H	2.15	0.43
1:B:152:ASN:HD21	1:B:154:ASN:HB2	1.83	0.43
1:A:397:ILE:CG1	1:A:397:ILE:O	2.40	0.43
1:A:117:VAL:HG11	1:A:181:SER:CB	2.49	0.43
1:B:75:LEU:HD21	1:B:158:PHE:CG	2.55	0.42
1:A:31:SER:HA	1:A:32:PRO:HD3	1.94	0.42
1:A:149:PRO:HA	1:A:152:ASN:HB2	2.02	0.42
1:A:23:GLU:N	1:A:23:GLU:OE1	2.47	0.41
1:B:75:LEU:HB2	1:B:80:VAL:HG22	2.02	0.41
1:A:61:LEU:HB2	1:A:62:PRO:HD3	2.03	0.41
1:A:51:ARG:HG3	1:A:52:THR:N	2.35	0.41
1:B:204:VAL:HA	1:B:252:LEU:HD13	2.02	0.41
1:B:342:LYS:HZ2	4:B:502:GOL:H32	1.81	0.41
1:B:174:ASN:HD22	1:B:174:ASN:HA	1.65	0.40
1:A:342:LYS:CE	4:A:502:GOL:H32	2.50	0.40
1:A:140:GLU:HG2	1:A:140:GLU:H	1.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	342/394 (87%)	335 (98%)	7 (2%)	0	100	100
1	B	354/394 (90%)	344 (97%)	9 (2%)	1 (0%)	50	43
All	All	696/788 (88%)	679 (98%)	16 (2%)	1 (0%)	59	55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	145	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/358 (90%)	312 (96%)	12 (4%)	45	40
1	B	330/358 (92%)	311 (94%)	19 (6%)	28	20
All	All	654/716 (91%)	623 (95%)	31 (5%)	36	29

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	MET
1	A	51	ARG
1	A	87	LEU
1	A	104	GLU
1	A	140	GLU
1	A	143	ASP
1	A	148	ASP
1	A	190	HIS
1	A	206	ASP
1	A	227	LEU
1	A	316	THR
1	A	393	ASP
1	B	37	MET
1	B	43	PHE
1	B	74	ILE
1	B	103	HIS
1	B	110	GLN
1	B	140	GLU
1	B	148	ASP
1	B	152	ASN
1	B	155	LEU
1	B	175	GLN
1	B	183	SER
1	B	184	GLN
1	B	212	ARG
1	B	214	LEU
1	B	227	LEU
1	B	233	LYS
1	B	350	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	381	SER
1	B	385	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	86	GLN
1	A	174	ASN
1	A	175	GLN
1	B	59	GLN
1	B	103	HIS
1	B	110	GLN
1	B	152	ASN
1	B	174	ASN
1	B	175	GLN
1	B	240	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	501	2	29,29,29	1.38	5 (17%)	45,45,45	2.15	9 (20%)
4	GOL	A	502	-	5,5,5	0.97	1 (20%)	5,5,5	0.86	0
3	ADP	B	501	2	29,29,29	1.25	3 (10%)	45,45,45	1.78	8 (17%)
4	GOL	B	502	-	5,5,5	0.39	0	5,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	501	2	-	0/16/32/32	0/1/3/3
4	GOL	A	502	-	-	0/4/4/4	0/0/0/0
3	ADP	B	501	2	-	0/16/32/32	0/1/3/3
4	GOL	B	502	-	-	0/4/4/4	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	501	ADP	C4-N9	-3.92	1.32	1.37
3	A	501	ADP	PB-O3A	3.33	1.66	1.60
3	A	501	ADP	C5-C4	3.09	1.47	1.40
3	B	501	ADP	C5-C4	3.08	1.47	1.40
3	A	501	ADP	C4-N9	-2.53	1.34	1.37
3	A	501	ADP	PB-O2B	-2.25	1.46	1.54
3	B	501	ADP	PB-O3A	2.19	1.64	1.60
3	A	501	ADP	C4-N3	2.13	1.38	1.35
4	A	502	GOL	O2-C2	-2.04	1.37	1.43

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ADP	N3-C2-N1	-9.71	120.59	128.71
3	B	501	ADP	N3-C2-N1	-5.68	123.96	128.71
3	A	501	ADP	N3-C4-N9	5.23	134.87	125.43
3	B	501	ADP	N3-C4-N9	4.99	134.45	125.43
3	A	501	ADP	C8-N9-C4	4.09	110.02	106.90
3	B	501	ADP	C8-N9-C4	3.90	109.87	106.90
3	A	501	ADP	C2-N1-C6	3.10	124.36	118.77
3	B	501	ADP	O2'-C2'-C1'	-2.90	102.44	111.23
3	B	501	ADP	C5-C4-N3	-2.58	120.07	125.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	ADP	O4'-C1'-N9	2.45	110.72	108.44
3	A	501	ADP	C5-C4-N3	-2.37	120.53	125.70
3	A	501	ADP	C2-N3-C4	2.35	120.69	114.01
3	B	501	ADP	C2-N3-C4	2.26	120.43	114.01
3	A	501	ADP	O2A-PA-O1A	2.24	124.71	112.21
3	B	501	ADP	C4-C5-N7	-2.14	107.69	109.52
3	A	501	ADP	C1'-N9-C4	-2.10	123.02	126.64
3	B	501	ADP	O3B-PB-O2B	2.05	115.59	107.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/394 (89%)	0.04	20 (5%) 24 23	29, 37, 51, 67	0
1	B	362/394 (91%)	0.00	11 (3%) 48 48	29, 36, 50, 57	0
All	All	716/788 (90%)	0.02	31 (4%) 34 33	29, 36, 51, 67	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	383	PHE	8.1
1	A	144	ALA	4.5
1	A	143	ASP	4.1
1	A	46	GLU	4.0
1	A	43	PHE	3.7
1	B	384	LYS	3.6
1	B	48	ALA	3.6
1	B	383	PHE	3.4
1	B	43	PHE	3.2
1	B	234	PHE	3.2
1	A	385	HIS	3.1
1	A	384	LYS	3.0
1	A	382	ALA	2.8
1	A	22	ARG	2.8
1	B	20	VAL	2.8
1	B	74	ILE	2.7
1	A	189	SER	2.6
1	A	47	ASN	2.6
1	A	148	ASP	2.5
1	A	182	ASP	2.5
1	B	47	ASN	2.4
1	A	74	ILE	2.4
1	A	397	ILE	2.4
1	A	51	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45	SER	2.3
1	B	147	VAL	2.3
1	A	49	CYS	2.2
1	B	271	PRO	2.2
1	B	272	SER	2.2
1	A	380	LYS	2.2
1	A	233	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	B	502	6/6	0.20	3.01	45,49,52,56	0
4	GOL	A	502	6/6	0.17	1.71	43,47,48,52	0
2	MG	A	500	1/1	0.10	-0.22	31,31,31,31	0
3	ADP	B	501	27/27	0.10	-0.42	24,26,30,32	0
3	ADP	A	501	27/27	0.09	-0.90	23,27,32,36	0
2	MG	B	500	1/1	0.07	-2.27	30,30,30,30	0

6.5 Other polymers ⓘ

There are no such residues in this entry.