



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 02:06 PM GMT

PDB ID : 2D3S  
Title : Crystal Structure of basic winged bean lectin with Tn-antigen  
Authors : Kulkarni, K.A.; Sinha, S.; Katiyar, S.; Surolia, A.; Vijayan, M.; Suguna, K.  
Deposited on : 2005-10-01  
Resolution : 2.35 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3327 (2.40-2.32)
Clashscore	79885	1064 (2.38-2.34)
Ramachandran outliers	78287	1048 (2.38-2.34)
Sidechain outliers	78261	1049 (2.38-2.34)
RSRZ outliers	66119	3330 (2.40-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	242	
1	B	242	
1	C	242	
1	D	242	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
5	NAG	C	601	-	X

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7912 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Basic agglutinin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	0	0	0
			1831	1179	307	345			
1	B	237	Total	C	N	O	0	0	0
			1839	1184	308	347			
1	C	237	Total	C	N	O	0	0	0
			1831	1177	307	347			
1	D	237	Total	C	N	O	0	0	0
			1807	1166	300	341			

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	3	Total	C	N	O	0	0
			38	22	2	14		
2	B	3	Total	C	N	O	0	0
			38	22	2	14		
2	B	3	Total	C	N	O	0	0
			38	22	2	14		
2	D	3	Total	C	N	O	0	0
			38	22	2	14		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			24	14	1	9		

- Molecule 4 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	4	Total	C	N	O	0	0
			49	28	2	19		

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Ca	0	0
			1	1		
6	A	1	Total	Ca	0	0
			1	1		
6	D	1	Total	Ca	0	0
			1	1		
6	C	1	Total	Ca	0	0
			1	1		

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	Mn	0	0
			1	1		
7	A	1	Total	Mn	0	0
			1	1		
7	D	1	Total	Mn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Mn	0	0
			1	1		

- Molecule 8 is O-(2-ACETAMIDO-2-DEOXY-ALPHA-D-GALACTOPYRANOSYL)-L-SERINE (three-letter code: TNR) (formula:  $C_{11}H_{20}N_2O_8$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			21	11	2	8		
8	B	1	Total	C	N	O	0	0
			21	11	2	8		
8	C	1	Total	C	N	O	0	0
			21	11	2	8		
8	D	1	Total	C	N	O	0	0
			21	11	2	8		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	89	Total	O	0	0
			89	89		
9	B	79	Total	O	0	0
			79	79		
9	C	54	Total	O	0	0
			54	54		
9	D	51	Total	O	0	0
			51	51		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Basic agglutinin

Chain A:




#### • Molecule 1: Basic agglutinin

Chain B:




#### • Molecule 1: Basic agglutinin


Chain C:



#### • Molecule 1: Basic agglutinin

Chain D:





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.87Å 91.90Å 73.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.35 28.56 – 2.34	Depositor EDS
% Data completeness (in resolution range)	97.5 (30.00-2.35) 96.8 (28.56-2.34)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.46 (at 2.34Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.198 , 0.244 0.193 , 0.207	Depositor DCC
$R_{free}$ test set	1802 reflections (4.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtriage
Anisotropy	0.547	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 37.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 44523 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7912	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, TNR, CA, MN, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/1884	0.69	1/2580 (0.0%)
1	B	0.40	0/1892	0.70	0/2589
1	C	0.38	0/1884	0.65	0/2579
1	D	0.36	0/1859	0.65	1/2548 (0.0%)
All	All	0.38	0/7519	0.67	2/10296 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	91	PHE	N-CA-C	-5.59	95.90	111.00
1	D	91	PHE	N-CA-C	-5.32	96.64	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1777	39	0
1	B	1839	0	1791	36	0
1	C	1831	0	1770	41	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1807	0	1741	50	0
2	A	38	0	34	0	0
2	B	76	0	68	8	0
2	D	38	0	34	0	0
3	A	24	0	22	5	0
4	C	49	0	43	0	0
5	C	14	0	13	0	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	A	21	0	19	0	0
8	B	21	0	19	0	0
8	C	21	0	19	0	0
8	D	21	0	19	0	0
9	A	89	0	0	3	0
9	B	79	0	0	2	0
9	C	54	0	0	1	0
9	D	51	0	0	1	0
All	All	7912	0	7369	172	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

The worst 5 of 172 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:83:PRO:HG2	1:C:219:ASN:HB2	1.52	0.92
1:D:197:VAL:HG23	1:D:198:LEU:HG	1.53	0.90
1:B:83:PRO:HG2	1:B:219:ASN:HB2	1.55	0.87
3:A:601:NAG:H3	3:A:602:FUC:H5	1.58	0.86
2:B:601:NAG:H4	2:B:603:NAG:HN2	1.40	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/242 (97%)	220 (94%)	11 (5%)	4 (2%)	14	11
1	B	235/242 (97%)	224 (95%)	10 (4%)	1 (0%)	43	52
1	C	235/242 (97%)	219 (93%)	14 (6%)	2 (1%)	25	27
1	D	235/242 (97%)	218 (93%)	14 (6%)	3 (1%)	18	17
All	All	940/968 (97%)	881 (94%)	49 (5%)	10 (1%)	21	21

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	83	PRO
1	C	83	PRO
1	D	83	PRO
1	D	107	PHE
1	B	83	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/211 (96%)	194 (96%)	8 (4%)	42	56
1	B	204/211 (97%)	198 (97%)	6 (3%)	55	71
1	C	202/211 (96%)	195 (96%)	7 (4%)	48	63
1	D	196/211 (93%)	188 (96%)	8 (4%)	41	54
All	All	804/844 (95%)	775 (96%)	29 (4%)	47	61

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	196	GLN
1	C	102	GLU
1	D	196	GLN
1	C	43	TRP
1	C	143	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	203	ASN
1	C	84	HIS
1	D	133	GLN
1	C	38	ASN
1	C	97	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

18 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	A	501	1,2	12,14,15	0.54	0	15,19,21	0.77	0
2	FUC	A	502	2	9,10,11	0.52	0	10,14,16	0.24	0
2	NAG	A	503	2	12,14,15	0.44	0	15,19,21	0.75	0
3	NAG	A	601	1,3	12,14,15	0.66	0	15,19,21	0.90	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FUC	A	602	3	9,10,11	0.37	0	10,14,16	0.35	0
2	NAG	B	501	1,2	12,14,15	0.54	0	15,19,21	0.73	0
2	FUC	B	502	2	9,10,11	0.45	0	10,14,16	0.28	0
2	NAG	B	503	2	12,14,15	0.47	0	15,19,21	0.69	0
2	NAG	B	601	1,2	12,14,15	0.61	0	15,19,21	0.71	0
2	FUC	B	602	2	9,10,11	0.43	0	10,14,16	0.24	0
2	NAG	B	603	2	12,14,15	0.43	0	15,19,21	0.77	0
4	NAG	C	501	1,4	12,14,15	0.53	0	15,19,21	0.73	0
4	FUC	C	502	4	9,10,11	0.45	0	10,14,16	0.25	0
4	NAG	C	503	4	12,14,15	0.63	0	15,19,21	0.93	0
4	BMA	C	504	4	10,11,12	0.35	0	11,15,17	0.59	0
2	NAG	D	601	1,2	12,14,15	0.52	0	15,19,21	0.78	0
2	FUC	D	602	2	9,10,11	0.45	0	10,14,16	0.22	0
2	NAG	D	603	2	12,14,15	0.47	0	15,19,21	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	501	1,2	-	0/6/23/26	0/1/1/1
2	FUC	A	502	2	-	0/0/17/20	0/1/1/1
2	NAG	A	503	2	-	0/6/23/26	0/1/1/1
3	NAG	A	601	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	602	3	-	0/0/17/20	0/1/1/1
2	NAG	B	501	1,2	-	0/6/23/26	0/1/1/1
2	FUC	B	502	2	-	0/0/17/20	0/1/1/1
2	NAG	B	503	2	-	0/6/23/26	0/1/1/1
2	NAG	B	601	1,2	-	0/6/23/26	0/1/1/1
2	FUC	B	602	2	-	0/0/17/20	0/1/1/1
2	NAG	B	603	2	-	0/6/23/26	0/1/1/1
4	NAG	C	501	1,4	-	0/6/23/26	0/1/1/1
4	FUC	C	502	4	-	0/0/17/20	0/1/1/1
4	NAG	C	503	4	-	0/6/23/26	0/1/1/1
4	BMA	C	504	4	-	0/2/19/22	0/1/1/1
2	NAG	D	601	1,2	-	0/6/23/26	0/1/1/1
2	FUC	D	602	2	-	0/0/17/20	0/1/1/1
2	NAG	D	603	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 8 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	TNR	A	401	-	21,21,21	0.69	0	29,29,29	1.01	2 (6%)
8	TNR	B	1401	-	21,21,21	0.71	0	29,29,29	1.04	2 (6%)
8	TNR	C	2401	-	21,21,21	0.79	0	29,29,29	1.02	2 (6%)
5	NAG	C	601	1	12,14,15	0.44	0	15,19,21	0.72	0
8	TNR	D	3401	-	21,21,21	0.79	0	29,29,29	1.03	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	TNR	A	401	-	-	0/15/35/35	0/1/1/1
8	TNR	B	1401	-	-	0/15/35/35	0/1/1/1
8	TNR	C	2401	-	-	0/15/35/35	0/1/1/1
5	NAG	C	601	1	-	0/6/23/26	0/1/1/1
8	TNR	D	3401	-	-	0/15/35/35	0/1/1/1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	1401	TNR	CB-O1-C1	2.94	119.67	113.81
8	D	3401	TNR	CB-O1-C1	2.84	119.46	113.81

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	401	TNR	CB-O1-C1	2.81	119.41	113.81
8	C	2401	TNR	CB-O1-C1	2.68	119.15	113.81
8	B	1401	TNR	OXT-C-O	-2.21	119.08	124.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	237/242 (97%)	-0.07	7 (2%) 48 52	24, 36, 57, 71	0
1	B	237/242 (97%)	-0.07	3 (1%) 74 77	21, 34, 63, 75	0
1	C	237/242 (97%)	0.10	13 (5%) 24 27	29, 42, 65, 86	0
1	D	237/242 (97%)	0.32	17 (7%) 15 17	28, 47, 74, 85	0
All	All	948/968 (97%)	0.07	40 (4%) 35 38	21, 40, 68, 86	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	237	GLY	7.1
1	D	37	VAL	5.5
1	B	82	ARG	4.9
1	B	113	LEU	4.9
1	C	237	GLY	4.9

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	BMA	C	504	11/12	0.25	3.73	83,85,87,87	0
2	FUC	B	502	10/11	0.25	0.90	77,79,80,80	0
3	NAG	A	601	14/15	0.18	0.83	60,64,68,76	0
2	NAG	B	601	14/15	0.18	0.19	49,60,67,73	0
2	FUC	A	502	10/11	0.15	0.06	66,68,68,69	0
4	NAG	C	503	14/15	0.18	-0.08	64,72,76,79	0
4	FUC	C	502	10/11	0.25	-0.13	70,71,72,75	0
2	NAG	B	503	14/15	0.28	-0.17	75,77,79,79	0
4	NAG	C	501	14/15	0.17	-0.18	58,60,66,67	0
2	NAG	B	501	14/15	0.16	-0.18	60,66,71,73	0
2	NAG	A	501	14/15	0.11	-0.45	59,61,64,71	0
2	NAG	A	503	14/15	0.26	-0.49	76,79,82,83	0
2	FUC	B	602	10/11	0.20	-0.75	58,62,63,63	0
2	NAG	D	601	14/15	0.13	-1.16	75,79,83,84	0
3	FUC	A	602	10/11	0.36	-	80,82,83,84	0
2	NAG	B	603	14/15	0.32	-	78,80,84,85	0
2	NAG	D	603	14/15	0.38	-	88,90,91,91	0
2	FUC	D	602	10/11	0.37	-	86,88,89,89	0

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	C	601	14/15	0.26	2.84	63,65,68,70	0
6	CA	D	3303	1/1	0.17	0.81	38,38,38,38	0
6	CA	B	1303	1/1	0.15	0.81	34,34,34,34	0
6	CA	C	2303	1/1	0.14	-0.09	43,43,43,43	0
6	CA	A	303	1/1	0.13	-0.23	34,34,34,34	0
8	TNR	C	2401	21/21	0.12	-0.28	45,50,59,60	0
8	TNR	A	401	21/21	0.12	-0.31	40,43,56,58	0
8	TNR	D	3401	21/21	0.13	-0.45	44,51,65,67	0
8	TNR	B	1401	21/21	0.11	-0.46	33,37,53,55	0
7	MN	B	1300	1/1	0.09	-0.74	35,35,35,35	0
7	MN	A	300	1/1	0.06	-1.81	43,43,43,43	0
7	MN	C	2300	1/1	0.05	-1.97	49,49,49,49	0
7	MN	D	3300	1/1	0.05	-2.12	49,49,49,49	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.