



# Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 01:22 AM GMT

PDB ID : 3D4Z  
Title : GOLGI MANNOSIDASE II complex with gluco-imidazole  
Authors : Kuntz, D.A.; Tarling, C.A.; Withers, S.G.; Rose, D.R.  
Deposited on : 2008-05-15  
Resolution : 1.39 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1246 (1.42-1.38)
Ramachandran outliers	78287	1206 (1.42-1.38)
Sidechain outliers	78261	1205 (1.42-1.38)
RSRZ outliers	66119	1097 (1.42-1.38)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	1045	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9422 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-mannosidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1016	Total	C	N	O	S	0	9	0
			8264	5256	1445	1522	41			


There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	ARG	-	EXPRESSION TAG	UNP Q24451
A	2	SER	-	EXPRESSION TAG	UNP Q24451
A	3	SER	-	EXPRESSION TAG	UNP Q24451
A	4	HIS	-	EXPRESSION TAG	UNP Q24451
A	5	HIS	-	EXPRESSION TAG	UNP Q24451
A	6	HIS	-	EXPRESSION TAG	UNP Q24451
A	7	HIS	-	EXPRESSION TAG	UNP Q24451
A	8	HIS	-	EXPRESSION TAG	UNP Q24451
A	9	HIS	-	EXPRESSION TAG	UNP Q24451
A	10	GLY	-	EXPRESSION TAG	UNP Q24451
A	11	GLU	-	EXPRESSION TAG	UNP Q24451
A	12	PHE	-	EXPRESSION TAG	UNP Q24451
A	907	LYS	GLU	SEE REMARK 999	UNP Q24451

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is (4R)-2-METHYLPENTANE-2,4-DIOL (three-letter code: MRD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is GLUCOIMIDAZOLE (three-letter code: GIM) (formula:  $C_8H_{13}N_2O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	2	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1135	Total 1135	O 1135	0	6

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-mannosidase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.65Å 109.30Å 137.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.39 29.63 – 1.39	Depositor EDS
% Data completeness (in resolution range)	97.6 (30.00-1.39) 94.7 (29.63-1.39)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.06 (at 1.39Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.144 , 0.181 0.139 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	11.2	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 59.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 203042 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9422	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, ZN, GIM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.57	0/8500	1.09	15/11541 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	793	ARG	NE-CZ-NH1	-5.98	117.31	120.30
1	A	422	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	A	617	ARG	CA-CB-CG	5.94	126.47	113.40
1	A	617	ARG	NE-CZ-NH2	5.83	123.21	120.30
1	A	616	TYR	CB-CG-CD1	-5.78	117.53	121.00
1	A	869	ARG	NE-CZ-NH1	-5.76	117.42	120.30
1	A	427	ARG	NE-CZ-NH2	-5.49	117.55	120.30
1	A	820	ASP	CB-CG-OD1	5.38	123.14	118.30
1	A	804	PHE	CB-CG-CD1	5.37	124.56	120.80
1	A	102	TYR	CB-CG-CD1	5.28	124.17	121.00
1	A	818	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	A	884	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	430	ARG	NE-CZ-NH1	5.16	122.88	120.30
1	A	370	PHE	CB-CG-CD2	-5.09	117.24	120.80
1	A	381	GLU	OE1-CD-OE2	-5.03	117.26	123.30



There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	954	ARG	Sidechain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8264	0	0	26	0
2	A	1	0	0	0	0
3	A	8	0	0	0	0
4	A	14	0	0	1	0
5	A	1135	0	0	17	0
All	All	9422	0	0	26	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (26) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:54:PHE:N	5:A:2162:HOH:O	2.34	0.60
1:A:954:ARG:NH1	5:A:1364:HOH:O	2.33	0.59
1:A:992:GLU:OE1	1:A:992:GLU:N	2.40	0.54
1:A:51:ARG:NH2	5:A:1067:HOH:O	2.40	0.54
1:A:583:LEU:CD2	1:A:583:LEU:O	2.58	0.52
1:A:916:SER:OG	1:A:955:GLU:OE1	2.32	0.48
1:A:539:SER:O	1:A:539:SER:OG	2.30	0.47
1:A:154:LYS:NZ	5:A:2146:HOH:O	2.48	0.46
1:A:621:LYS:NZ	5:A:1580:HOH:O	2.49	0.46
1:A:986:GLN:NE2	5:A:1130:HOH:O	2.47	0.46
1:A:610:GLN:NE2	5:A:1449:HOH:O	2.49	0.46
1:A:714:LYS:NZ	5:A:1249:HOH:O	2.50	0.45
1:A:945:GLN:NE2	5:A:1375:HOH:O	2.49	0.45
1:A:146:LYS:NZ	5:A:1563:HOH:O	2.49	0.45
1:A:235:LYS:NZ	5:A:1665:HOH:O	2.49	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:842:THR:OG1	1:A:898:LYS:NZ	2.50	0.44
1:A:996:LYS:NZ	5:A:2016:HOH:O	2.50	0.44
1:A:300:LYS:NZ	5:A:2014:HOH:O	2.50	0.44
1:A:615:LYS:NZ	5:A:1446:HOH:O	2.50	0.44
1:A:700:THR:N	1:A:703:SER:OG	2.50	0.43
1:A:988:GLY:N	5:A:1139:HOH:O	2.52	0.43
1:A:341:ASP:OD2	4:A:1048:GIM:N1	2.52	0.42
1:A:80:LYS:NZ	5:A:1631:HOH:O	2.52	0.42
1:A:97:GLN:OE1	1:A:105:HIS:ND1	2.52	0.42
1:A:538:ASP:O	1:A:538:ASP:OD1	2.37	0.41
1:A:711:LYS:NZ	5:A:1756:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1023/1045 (98%)	1000 (98%)	21 (2%)	2 (0%)	56 24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	95	TRP
1	A	204	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	911/929 (98%)	898 (99%)	13 (1%)	78 50

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	SER
1	A	57	ILE
1	A	78	HIS
1	A	144	ASN
1	A	275	CYS
1	A	375	GLN
1	A	447	SER
1	A	539	SER
1	A	663	GLN
1	A	691	GLN
1	A	828	TYR
1	A	906	SER
1	A	907	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	MRD	A	1047	-	7,7,7	0.23	0	10,10,10	0.65	0
4	GIM	A	1048	2	15,15,15	1.26	2 (13%)	19,22,22	112.07	8 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MRD	A	1047	-	-	0/5/5/5	0/0/0/0
4	GIM	A	1048	2	-	0/2/22/22	0/0/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1048	GIM	C8-N10	-2.77	1.36	1.39
4	A	1048	GIM	C5-N10	-2.00	1.47	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1048	GIM	C2-C1-N10	-488.37	118.36	128.81
4	A	1048	GIM	C8-N10-C1	-6.30	105.42	109.47
4	A	1048	GIM	C2-C1-N1	4.94	132.17	121.27
4	A	1048	GIM	C6-C5-C4	-3.37	106.35	112.49
4	A	1048	GIM	C4-C5-N10	3.10	114.88	109.44
4	A	1048	GIM	C1-C2-C3	-3.01	107.50	112.76
4	A	1048	GIM	C4-C3-C2	-2.18	107.95	110.64
4	A	1048	GIM	O4-C4-C5	-2.00	106.15	109.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1016/1045 (97%)	0.48	65 (6%) 19 18	7, 14, 31, 69	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	536	VAL	8.2
1	A	603	LEU	8.0
1	A	539	SER	7.8
1	A	993	HIS	7.0
1	A	534	SER	6.5
1	A	701	GLN	5.7
1	A	602	THR	5.6
1	A	638	ASP	5.3
1	A	720	HIS	5.3
1	A	78	HIS	5.2
1	A	703	SER	5.1
1	A	990	PRO	5.1
1	A	989	THR	4.9
1	A	702	ASP	4.9
1	A	639	SER	4.8
1	A	682	ASN	4.6
1	A	537	GLU	4.1
1	A	57	ILE	4.1
1	A	583	LEU	4.1
1	A	535	GLY	4.1
1	A	700	THR	4.0
1	A	77	ALA	3.9
1	A	655	ASN	3.9
1	A	73	LEU	3.6
1	A	600	HIS	3.6
1	A	1045	SER	3.5
1	A	139	HIS	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	72	PRO	3.2
1	A	538	ASP	3.2
1	A	297	CYS	3.1
1	A	683	GLY	3.1
1	A	395	ALA	3.1
1	A	1024	ASP	3.0
1	A	521	PHE	2.9
1	A	147	LEU	2.9
1	A	994	THR	2.8
1	A	389	ALA	2.8
1	A	392	ALA	2.7
1	A	992	GLU	2.7
1	A	614	THR	2.7
1	A	141	LEU	2.7
1	A	601	ASP	2.7
1	A	908	LEU	2.6
1	A	31	CYS	2.5
1	A	291	GLY	2.5
1	A	606	THR	2.5
1	A	1022	HIS	2.4
1	A	335	LEU	2.4
1	A	662	GLY	2.4
1	A	721	GLY	2.4
1	A	322	TRP	2.4
1	A	547	GLU	2.3
1	A	304	ARG	2.3
1	A	290	MET	2.3
1	A	1026	MET	2.2
1	A	1027	VAL	2.2
1	A	940	ILE	2.2
1	A	368	ALA	2.2
1	A	71	ASP	2.1
1	A	1040	VAL	2.1
1	A	256	ASP	2.1
1	A	677	SER	2.1
1	A	697	ILE	2.1
1	A	308	ASP	2.0
1	A	592	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates

There are no carbohydrates in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MRD	A	1047	8/8	0.14	1.39	14,18,27,27	0
4	GIM	A	1048	14/14	0.08	0.86	7,9,10,10	0
2	ZN	A	1046	1/1	0.04	-1.81	9,9,9,9	0

### 6.5 Other polymers

There are no such residues in this entry.