



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 01:09 AM GMT

PDB ID : 1D5T
Title : GUANINE NUCLEOTIDE DISSOCIATION INHIBITOR, ALPHA-ISOFORM
Authors : Peng, L.; Zeng, K.; Heine, A.; Moyer, B.; Greasley, S.E.; Kuhn, P.; Balch, W.E.; Wilson, I.A.
Deposited on : 1999-10-11
Resolution : 1.04 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1106 (1.14-0.94)
Clashscore	79885	1011 (1.12-0.96)
Ramachandran outliers	78287	1151 (1.14-0.94)
Sidechain outliers	78261	1150 (1.14-0.94)
RSRZ outliers	66119	1106 (1.14-0.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	433	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3827 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GUANINE NUCLEOTIDE DISSOCIATION INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	3430	2182	564	658	26	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	HIS	-	CONFLICT	UNP P21856
A	-1	HIS	-	CONFLICT	UNP P21856

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	387	Total 387	O 387	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GUANINE NUCLEOTIDE DISSOCIATION INHIBITOR

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.29Å 42.74Å 61.78Å 90.00° 104.55° 90.00°	Depositor
Resolution (Å)	10.00 – 1.04 34.77 – 1.04	Depositor EDS
% Data completeness (in resolution range)	91.0 (10.00-1.04) 92.1 (34.77-1.04)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 1.04Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.173 , 0.209 0.181 , 0.212	Depositor DCC
R_{free} test set	9973 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	11.0	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 43.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 207147 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3827	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	4/3513 (0.1%)	1.50	53/4752 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	275	ALA	C-O	37.82	1.95	1.23
1	A	208	ARG	CD-NE	-6.09	1.36	1.46
1	A	275	ALA	C-N	-5.38	1.21	1.34
1	A	40	GLY	N-CA	-5.09	1.38	1.46

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	LEU	C-N-CA	22.36	177.61	121.70
1	A	275	ALA	CA-C-N	16.96	154.52	117.20
1	A	208	ARG	CD-NE-CZ	16.07	146.10	123.60
1	A	275	ALA	O-C-N	-15.29	98.23	122.70
1	A	138	ARG	NE-CZ-NH1	14.27	127.43	120.30
1	A	23	MET	CG-SD-CE	13.84	122.34	100.20
1	A	138	ARG	CD-NE-CZ	13.28	142.19	123.60
1	A	391	ASP	CB-CG-OD1	11.28	128.45	118.30
1	A	366	VAL	CA-CB-CG2	-10.57	95.04	110.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39	TYR	C-N-CA	10.27	143.87	122.30
1	A	55	ARG	NE-CZ-NH1	9.30	124.95	120.30
1	A	141	ARG	NE-CZ-NH2	9.16	124.88	120.30
1	A	145	VAL	CA-CB-CG1	8.94	124.31	110.90
1	A	292	ARG	CD-NE-CZ	8.19	135.07	123.60
1	A	275	ALA	C-N-CA	8.08	141.90	121.70
1	A	360	THR	C-N-CA	8.02	141.75	121.70
1	A	141	ARG	NE-CZ-NH1	-7.82	116.39	120.30
1	A	275	ALA	CA-C-O	-7.80	103.72	120.10
1	A	208	ARG	NE-CZ-NH2	7.63	124.11	120.30
1	A	183	ASP	CB-CG-OD1	-7.54	111.51	118.30
1	A	313	ASP	CB-CG-OD2	7.24	124.81	118.30
1	A	290	ARG	CD-NE-CZ	7.13	133.59	123.60
1	A	195	ASP	O-C-N	-7.07	111.39	122.70
1	A	158	PHE	CB-CG-CD1	6.96	125.67	120.80
1	A	39	TYR	O-C-N	-6.92	111.44	123.20
1	A	249	TYR	CG-CD2-CE2	6.70	126.66	121.30
1	A	240	ARG	CD-NE-CZ	6.35	132.50	123.60
1	A	423	ARG	NE-CZ-NH2	6.27	123.43	120.30
1	A	50	GLU	OE1-CD-OE2	-6.26	115.79	123.30
1	A	93	TYR	CB-CG-CD1	-6.25	117.25	121.00
1	A	423	ARG	NE-CZ-NH1	-6.15	117.22	120.30
1	A	245	TYR	CB-CG-CD1	6.14	124.68	121.00
1	A	26	ASN	CB-CG-OD1	6.07	133.73	121.60
1	A	240	ARG	NE-CZ-NH2	5.98	123.29	120.30
1	A	35	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	A	124	THR	O-C-N	-5.90	113.26	122.70
1	A	377	ASP	CB-CG-OD1	-5.87	113.01	118.30
1	A	56	PHE	CB-CG-CD2	5.86	124.90	120.80
1	A	175	PHE	CB-CG-CD1	5.77	124.84	120.80
1	A	173	ARG	CD-NE-CZ	5.59	131.43	123.60
1	A	226	TYR	CB-CG-CD2	-5.51	117.69	121.00
1	A	70	ARG	NE-CZ-NH1	-5.47	117.56	120.30
1	A	35	ARG	CD-NE-CZ	5.40	131.16	123.60
1	A	26	ASN	OD1-CG-ND2	-5.35	109.60	121.90
1	A	158	PHE	CB-CG-CD2	-5.23	117.14	120.80
1	A	164	GLN	CA-CB-CG	5.16	124.74	113.40
1	A	117	TYR	CB-CG-CD1	5.08	124.05	121.00
1	A	257	ASP	CB-CG-OD1	5.08	122.88	118.30
1	A	184	PHE	CG-CD2-CE2	5.07	126.37	120.80
1	A	429	PHE	CB-CG-CD2	-5.06	117.25	120.80
1	A	68	ARG	CD-NE-CZ	5.05	130.67	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	TYR	CG-CD2-CE2	5.03	125.32	121.30
1	A	124	THR	C-N-CA	5.03	134.27	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	216	LEU	Peptide
1	A	275	ALA	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3430	0	0	9	0
2	A	10	0	0	0	0
3	A	387	0	0	5	0
All	All	3827	0	0	9	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 3.

All (9) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:275:ALA:O	1:A:275:ALA:C	1.95	1.05
1:A:275:ALA:O	1:A:276:ARG:N	2.44	0.47
1:A:55:ARG:NH2	3:A:1262:HOH:O	2.48	0.46
1:A:253:LYS:NZ	3:A:1166:HOH:O	2.49	0.46
1:A:290:ARG:NH1	1:A:391:ASP:OD1	2.50	0.45
1:A:240:ARG:NE	3:A:1377:HOH:O	2.49	0.45
1:A:290:ARG:NH2	3:A:1308:HOH:O	2.50	0.43
1:A:207:ASN:ND2	3:A:1244:HOH:O	2.50	0.43
1:A:123:GLU:OE1	1:A:141:ARG:NE	2.51	0.43

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	434/433 (100%)	426 (98%)	5 (1%)	3 (1%)	30 5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	216	LEU
1	A	217	ALA
1	A	220	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	381/380 (100%)	369 (97%)	12 (3%)	52 12

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-2	HIS
1	A	33[A]	MET
1	A	33[B]	MET
1	A	64	GLU
1	A	87	LEU
1	A	112	LYS
1	A	141	ARG
1	A	145	VAL
1	A	283	ASP
1	A	312	ASN
1	A	315	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	337	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	501	-	4,4,4	1.18	0	6,6,6	0.27	0
2	SO4	A	502	-	4,4,4	0.51	0	6,6,6	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	501	-	-	0/0/0/0	0/0/0/0
2	SO4	A	502	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	433/433 (100%)	0.89	37 (8%)	11 12	8, 14, 32, 60	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	219	TYR	13.6
1	A	217	ALA	10.6
1	A	218	ARG	8.8
1	A	220	GLY	7.9
1	A	216	LEU	5.7
1	A	247	GLY	5.5
1	A	221	LYS	4.1
1	A	58	LEU	3.6
1	A	261	GLU	3.3
1	A	426	GLY	3.2
1	A	431	PHE	3.1
1	A	-2	HIS	3.0
1	A	289	ASP	2.9
1	A	127	LEU	2.9
1	A	179	GLN	2.9
1	A	231	LEU	2.5
1	A	246	GLY	2.5
1	A	360	THR	2.5
1	A	298	ILE	2.4
1	A	63	PRO	2.4
1	A	18	ILE	2.3
1	A	10	LEU	2.3
1	A	404	TYR	2.2
1	A	308	ILE	2.2
1	A	145	VAL	2.2
1	A	383	ILE	2.2
1	A	429	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	241	LEU	2.1
1	A	240	ARG	2.1
1	A	421	TYR	2.1
1	A	266	VAL	2.1
1	A	77	ILE	2.1
1	A	259	ILE	2.1
1	A	425	ALA	2.0
1	A	135	PHE	2.0
1	A	262	ASN	2.0
1	A	417	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	A	502	5/5	0.09	-0.33	13,13,19,20	0
2	SO4	A	501	5/5	0.08	-0.53	15,17,23,25	0

6.5 Other polymers ⓘ

There are no such residues in this entry.