



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 05:16 AM GMT

PDB ID : 3D6G
Title : Fc fragment of IgG1 (Herceptin) with protein-A mimetic peptide dendrimer ligand.
Authors : Bujacz, A.D.; Redzynia, I.; Bujacz, G.D.; Dinon, F.; Pengo, P.; Fassina, G.
Deposited on : 2008-05-19
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	210	
1	B	210	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	X12	B	446	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4002 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ig gamma-1 chain C region.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	4	0
			1702	1084	288	324	6			
1	B	209	Total	C	N	O	S	0	2	0
			1682	1070	285	321	6			

- Molecule 2 is a polymer of unknown type called SUGAR (8-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	8	Total	C	N	O	0	0
			99	56	4	39		
2	B	8	Total	C	N	O	0	0
			99	56	4	39		

- Molecule 3 is 2-[[[(2S)-2,6-BIS[[[(2S)-2,6-BIS[[[(2R)-2-[[[(2R,3R)-2-[[[(2R)-2-AMINO-5-CARBA MIMIDAMIDO-PENTANOYL]AMINO]-3-HYDROXY-BUTANOYL]AMINO]-3-(4-HYDROXYPHENYL)PROPANOYL]AMINO]HEXANOYL]AMINO]HEXANOYL]AMINO]ETHANOICACID (three-letter code: X12) (formula: C₉₆H₁₅₃N₃₁O₂₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	0	0
			152	96	31	25		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	145	Total	O	0	0
			145	145		
4	B	123	Total	O	0	0
			123	123		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ig gamma-1 chain C region

Chain A: 



- Molecule 1: Ig gamma-1 chain C region

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.78Å 79.16Å 139.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.30 30.11 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (40.00-2.30) 99.3 (30.11-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.188 , 0.257 0.202 , 0.263	Depositor DCC
R_{free} test set	806 reflections (3.33%)	DCC
Wilson B-factor (Å ²)	53.1	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 98.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 25040 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4002	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, X12, BMA, NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/1762	0.77	0/2400
1	B	0.76	0/1734	0.74	0/2361
All	All	0.76	0/3496	0.75	0/4761

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1702	0	1674	12	0
1	B	1682	0	1654	43	0
2	A	99	0	85	2	0
2	B	99	0	85	2	0
3	B	152	0	146	13	0
4	A	145	0	0	0	0
4	B	123	0	0	4	0
All	All	4002	0	3644	67	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including

hydrogens) of the entry. The overall clashscore for this entry is 9.

All (67) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
3:B:446:X12:HNBS	3:B:446:X12:HBI	1.29	0.96
1:B:276:ASN:HA	4:B:532:HOH:O	1.65	0.95
1:B:282:VAL:HG12	1:B:283:GLU:N	1.91	0.83
3:B:446:X12:HBI	3:B:446:X12:NBS	1.93	0.83
1:B:429:HIS:HD2	1:B:431:ALA:H	1.32	0.77
1:B:274:LYS:HG2	1:B:275:PHE:N	2.01	0.76
1:B:278:TYR:HD1	1:B:282:VAL:O	1.73	0.72
1:B:282:VAL:HG12	1:B:283:GLU:O	1.90	0.72
1:B:274:LYS:HG2	1:B:275:PHE:H	1.55	0.71
1:B:278:TYR:CD1	1:B:282:VAL:O	2.44	0.70
1:B:242:LEU:HD13	1:B:336:ILE:HB	1.72	0.70
1:A:297:ASN:HD22	2:A:1:NAG:H83	1.58	0.69
1:B:429:HIS:CD2	1:B:431:ALA:H	2.12	0.65
1:B:268:HIS:CD2	1:B:300:TYR:CE1	2.86	0.64
1:B:322:LYS:HG2	1:B:333:GLU:HG2	1.79	0.64
1:B:264:VAL:HG13	1:B:265:ASP:H	1.63	0.64
1:B:268:HIS:CD2	1:B:300:TYR:HE1	2.18	0.61
1:B:278:TYR:HB3	1:B:282:VAL:O	1.99	0.61
1:B:264:VAL:HG13	1:B:265:ASP:N	2.17	0.60
1:B:358:LEU:HD23	1:B:363:VAL:HG11	1.84	0.60
1:B:282:VAL:CG1	1:B:283:GLU:N	2.61	0.59
1:A:374:PRO:O	1:A:429:HIS:HE1	1.86	0.59
1:B:250:THR:O	3:B:446:X12:NDU	2.36	0.58
1:B:325:ASN:HB3	1:B:328:LEU:HG	1.88	0.56
1:B:278:TYR:CB	1:B:282:VAL:O	2.56	0.54
1:B:282:VAL:HG12	1:B:283:GLU:H	1.70	0.53
1:A:272[A]:GLU:HA	1:A:272[A]:GLU:OE1	2.08	0.53
1:B:264:VAL:CG1	1:B:265:ASP:N	2.71	0.53
1:A:429:HIS:CD2	1:A:431:ALA:H	2.27	0.52
3:B:446:X12:HCQ	3:B:446:X12:HB5	1.92	0.52
1:A:295:GLN:HE21	2:A:1:NAG:H62	1.75	0.51
1:A:429:HIS:HD2	1:A:431:ALA:H	1.57	0.51
3:B:446:X12:HCJ	3:B:446:X12:HC6	1.93	0.50
1:B:274:LYS:CG	1:B:275:PHE:N	2.74	0.50
1:A:259:VAL:HG23	1:A:308:VAL:HG11	1.94	0.50
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.48	0.49
1:B:276:ASN:HB2	1:B:322:LYS:HB2	1.95	0.48
1:B:325:ASN:HD21	1:B:327:ALA:HB3	1.79	0.48
1:A:311[B]:GLN:H	1:A:311[B]:GLN:NE2	2.11	0.48
3:B:446:X12:HAW	3:B:446:X12:NBC	2.29	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:274:LYS:CG	1:B:275:PHE:H	2.26	0.46
3:B:446:X12:HAT	3:B:446:X12:HAQA	1.48	0.45
3:B:446:X12:HNAG	3:B:446:X12:CC1	2.28	0.45
1:B:435:HIS:HE1	3:B:446:X12:NDT	2.14	0.45
1:B:424:SER:HB3	1:B:438:GLN:HE21	1.82	0.45
1:B:266:VAL:HB	1:B:300:TYR:HB2	2.00	0.44
1:B:325:ASN:ND2	1:B:327:ALA:H	2.15	0.44
1:A:358:LEU:HD23	1:A:363:VAL:HG11	1.98	0.44
1:B:282:VAL:CG1	1:B:283:GLU:H	2.29	0.43
1:B:325:ASN:ND2	1:B:327:ALA:HB3	2.34	0.43
1:B:365:LEU:HB3	1:B:441:LEU:HD23	2.01	0.42
2:B:3:BMA:O4	4:B:512:HOH:O	2.21	0.42
1:B:276:ASN:HB2	1:B:322:LYS:CB	2.48	0.42
1:B:335:THR:HG22	1:B:336:ILE:N	2.34	0.42
1:B:416:ARG:HD3	4:B:509:HOH:O	2.18	0.42
1:B:238:PRO:HB3	1:B:264:VAL:O	2.20	0.41
1:B:271:PRO:HG2	4:B:560:HOH:O	2.20	0.41
1:B:374:PRO:O	1:B:429:HIS:HE1	2.02	0.41
1:B:435:HIS:HE1	3:B:446:X12:CDS	2.33	0.41
1:B:386:GLN:HA	1:B:387:PRO:HD3	1.96	0.41
3:B:446:X12:HCU	3:B:446:X12:HDH	2.01	0.41
1:B:290:LYS:HA	1:B:291:PRO:HD3	1.79	0.41
3:B:446:X12:HBP	3:B:446:X12:HNBA	1.86	0.40
2:B:4:MAN:H4	2:B:5:NAG:H83	2.03	0.40
1:A:325:ASN:OD1	1:A:327:ALA:HB3	2.21	0.40
3:B:446:X12:CAW	3:B:446:X12:NBC	2.84	0.40
1:A:365:LEU:HB3	1:A:441:LEU:HD23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/210 (101%)	210 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	209/210 (100%)	202 (97%)	6 (3%)	1 (0%)	38	45
All	All	421/420 (100%)	412 (98%)	8 (2%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	238	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/195 (102%)	194 (98%)	5 (2%)	60	77
1	B	196/195 (100%)	187 (95%)	9 (5%)	37	48
All	All	395/390 (101%)	381 (96%)	14 (4%)	50	63

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	390	ASN
1	A	413	ASP
1	A	419	GLN
1	A	442	SER
1	A	443	LEU
1	B	242	LEU
1	B	292	ARG
1	B	293	GLU
1	B	301[A]	ARG
1	B	301[B]	ARG
1	B	322	LYS
1	B	326	LYS
1	B	354	SER
1	B	418	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	295	GLN
1	A	361	ASN
1	A	386	GLN
1	A	419	GLN
1	A	429	HIS
1	A	438	GLN
1	B	268	HIS
1	B	325	ASN
1	B	389	ASN
1	B	390	ASN
1	B	418	GLN
1	B	419	GLN
1	B	429	HIS
1	B	435	HIS
1	B	438	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

16 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1	1,2	12,14,15	1.08	1 (8%)	15,19,21	2.11	5 (33%)
2	NAG	A	2	2	12,14,15	0.87	0	15,19,21	1.76	5 (33%)
2	BMA	A	3	2	10,11,12	0.86	0	11,15,17	1.76	3 (27%)
2	MAN	A	4	2	10,11,12	0.86	0	11,15,17	1.86	3 (27%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	5	2	12,14,15	1.11	1 (8%)	15,19,21	1.25	1 (6%)
2	MAN	A	6	2	10,11,12	0.98	0	11,15,17	1.28	1 (9%)
2	NAG	A	7	2	12,14,15	0.83	1 (8%)	15,19,21	1.60	3 (20%)
2	FUC	A	8	2	9,10,11	0.67	0	10,14,16	1.05	0
2	NAG	B	1	1,2	12,14,15	0.91	0	15,19,21	1.53	2 (13%)
2	NAG	B	2	2	12,14,15	1.05	0	15,19,21	1.30	2 (13%)
2	BMA	B	3	2	10,11,12	0.53	0	11,15,17	1.53	3 (27%)
2	MAN	B	4	2	10,11,12	0.78	0	11,15,17	1.40	1 (9%)
2	NAG	B	5	2	12,14,15	1.08	2 (16%)	15,19,21	0.92	0
2	MAN	B	6	2	10,11,12	0.53	0	11,15,17	0.79	1 (9%)
2	NAG	B	7	2	12,14,15	0.99	1 (8%)	15,19,21	0.93	1 (6%)
2	FUC	B	8	2	9,10,11	0.76	0	10,14,16	1.11	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2	2	-	0/6/23/26	0/1/1/1
2	BMA	A	3	2	-	0/2/19/22	0/1/1/1
2	MAN	A	4	2	-	0/2/19/22	0/1/1/1
2	NAG	A	5	2	-	0/6/23/26	0/1/1/1
2	MAN	A	6	2	-	0/2/19/22	0/1/1/1
2	NAG	A	7	2	-	0/6/23/26	0/1/1/1
2	FUC	A	8	2	-	0/0/17/20	0/1/1/1
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	NAG	B	5	2	-	0/6/23/26	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	NAG	B	7	2	-	0/6/23/26	0/1/1/1
2	FUC	B	8	2	-	0/0/17/20	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5	NAG	C8-C7	2.42	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	7	NAG	C2-N2	2.38	1.49	1.46
2	A	7	NAG	C8-C7	2.21	1.55	1.50
2	A	1	NAG	O5-C5	2.11	1.49	1.45
2	B	5	NAG	C8-C7	2.03	1.54	1.50
2	B	5	NAG	C4-C5	2.01	1.57	1.53

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	NAG	O6-C6-C5	-4.41	96.19	111.36
2	A	2	NAG	C3-C4-C5	-4.30	102.53	110.20
2	B	4	MAN	O2-C2-C3	-4.01	101.53	110.18
2	B	1	NAG	O5-C5-C4	3.98	115.70	110.65
2	A	1	NAG	C3-C2-N2	-3.77	106.02	111.76
2	A	5	NAG	O5-C5-C6	-3.75	103.05	106.98
2	A	7	NAG	O5-C5-C4	-3.72	105.93	110.65
2	A	3	BMA	O6-C6-C5	-3.63	98.85	111.36
2	A	6	MAN	O5-C5-C6	3.51	110.66	106.98
2	A	4	MAN	O2-C2-C3	-3.46	102.71	110.18
2	A	1	NAG	O5-C5-C4	3.43	115.01	110.65
2	A	4	MAN	C4-C3-C2	-3.38	105.97	110.50
2	A	7	NAG	O5-C5-C6	-3.28	103.54	106.98
2	A	1	NAG	C6-C5-C4	-3.08	105.56	113.00
2	A	3	BMA	O3-C3-C4	-2.98	103.67	110.35
2	B	2	NAG	O5-C5-C4	2.83	114.25	110.65
2	B	3	BMA	C4-C3-C2	2.62	114.02	110.50
2	A	7	NAG	C6-C5-C4	2.58	119.24	113.00
2	B	1	NAG	C6-C5-C4	-2.57	106.80	113.00
2	A	1	NAG	O5-C5-C6	2.52	109.63	106.98
2	B	3	BMA	O6-C6-C5	-2.43	103.00	111.36
2	A	2	NAG	O4-C4-C5	-2.42	102.90	109.28
2	A	4	MAN	O5-C5-C4	2.41	113.72	110.65
2	B	3	BMA	C6-C5-C4	-2.31	107.42	113.00
2	A	2	NAG	C8-C7-N2	-2.31	111.59	116.11
2	B	8	FUC	C4-C3-C2	-2.27	107.46	110.50
2	A	2	NAG	O7-C7-N2	2.24	126.58	121.90
2	B	2	NAG	O4-C4-C5	-2.23	103.40	109.28
2	A	3	BMA	O5-C5-C6	2.20	109.29	106.98
2	A	2	NAG	C3-C2-N2	-2.19	108.43	111.76
2	B	6	MAN	O5-C5-C6	-2.12	104.76	106.98
2	B	7	NAG	O5-C5-C4	-2.03	108.07	110.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	X12	B	446	-	155,155,155	0.93	3 (1%)	205,205,205	1.19	13 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	X12	B	446	-	-	0/193/193/193	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	446	X12	CBG-NBH	4.80	1.57	1.46
3	B	446	X12	CB2-NBK	4.76	1.53	1.32
3	B	446	X12	CB2-NBH	2.25	1.38	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	446	X12	CBH-NBG-CBX	6.04	134.99	121.63
3	B	446	X12	CBG-NBH-CB2	-4.99	114.75	124.14
3	B	446	X12	CBD-CB1-NAV	3.84	121.41	116.31
3	B	446	X12	NBJ-CB2-NBH	-3.56	112.11	119.69
3	B	446	X12	CDG-NDF-CDV	3.53	130.94	121.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	446	X12	CAW-NAV-CB1	3.34	130.46	121.81
3	B	446	X12	CAP-NAO-CBE	3.22	128.76	121.63
3	B	446	X12	CBT-CBX-NBG	3.08	123.27	116.48
3	B	446	X12	CAT-NAU-C	2.96	128.66	122.57
3	B	446	X12	CCU-NCT-CDK	2.87	127.97	121.63
3	B	446	X12	CB5-NBW-CCN	2.77	127.76	121.63
3	B	446	X12	CDN-CDV-NDF	2.33	119.40	116.31
3	B	446	X12	CBI-CBH-NBG	2.03	115.06	110.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/210 (100%)	-0.20	3 (1%) 72 80	65, 71, 79, 86	0
1	B	209/210 (99%)	0.32	17 (8%) 12 18	65, 72, 78, 84	0
All	All	419/420 (99%)	0.06	20 (4%) 31 39	65, 72, 79, 86	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	445	PRO	6.1
1	B	296	TYR	5.6
1	B	300	TYR	4.9
1	B	330	ALA	4.3
1	B	366	THR	3.4
1	A	444	SER	3.4
1	B	281	GLY	3.2
1	B	239	SER	3.1
1	B	368	LEU	2.8
1	B	295	GLN	2.7
1	B	236	GLY	2.7
1	B	367	CYS	2.6
1	B	326	LYS	2.5
1	B	297	ASN	2.4
1	B	237	GLY	2.3
1	A	366	THR	2.3
1	B	351	LEU	2.2
1	B	327	ALA	2.1
1	B	408	SER	2.1
1	B	240	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	BMA	B	3	11/12	0.23	3.21	62,65,66,68	0
2	NAG	B	1	14/15	0.40	2.85	59,65,66,68	0
2	NAG	B	2	14/15	0.33	2.59	62,68,71,71	0
2	FUC	A	8	10/11	0.16	0.83	71,75,75,76	0
2	FUC	B	8	10/11	0.39	0.39	69,70,71,71	0
2	NAG	B	7	14/15	0.17	-0.14	64,66,68,70	0
2	NAG	A	2	14/15	0.11	-0.29	67,69,70,73	0
2	NAG	A	1	14/15	0.08	-0.37	66,69,74,79	0
2	NAG	A	7	14/15	0.10	-0.98	65,69,73,74	0
2	MAN	B	6	11/12	0.14	-1.58	65,67,69,70	0
2	BMA	A	3	11/12	0.07	-3.17	64,68,71,71	0
2	MAN	A	6	11/12	0.09	-3.60	66,69,70,72	0
2	MAN	A	4	11/12	0.10	-	64,68,72,77	0
2	MAN	B	4	11/12	0.19	-	61,64,67,72	0
2	NAG	A	5	14/15	0.20	-	82,86,89,90	0
2	NAG	B	5	14/15	0.28	-	76,78,79,79	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	X12	B	446	152/152	0.76	11.09	51,79,83,83	152

6.5 Other polymers ⓘ

There are no such residues in this entry.