



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 04:17 PM GMT

PDB ID : 1DDO
Title : REDUCED D-AMINO ACID OXIDASE FROM PIG KIDNEY IN COMPLEX
WITH IMINO-TRP
Authors : Todone, F.; Mattevi, A.
Deposited on : 1997-01-16
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)
RSRZ outliers	66119	1008 (3.18-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	347	
1	B	347	
1	C	347	
1	D	347	
1	E	347	
1	F	347	
1	G	347	
1	H	347	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
3	ITR	A	349	-	X

2 Entry composition


There are 5 unique types of molecules in this entry. The entry contains 22441 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-AMINO ACID OXIDASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	161	0	0
			2720	1749	473	489	9			
1	B	339	Total	C	N	O	S	190	0	0
			2720	1749	473	489	9			
1	C	339	Total	C	N	O	S	173	0	0
			2720	1749	473	489	9			
1	D	339	Total	C	N	O	S	131	0	0
			2720	1749	473	489	9			
1	E	339	Total	C	N	O	S	164	0	0
			2720	1749	473	489	9			
1	F	339	Total	C	N	O	S	154	0	0
			2720	1749	473	489	9			
1	G	339	Total	C	N	O	S	210	0	0
			2720	1749	473	489	9			
1	H	339	Total	C	N	O	S	166	0	0
			2720	1749	473	489	9			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	E	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	F	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	G	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	H	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is IMINO-TRYPTOPHAN (three-letter code: ITR) (formula: C₁₁H₁₀N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			15	11	2	2		
3	B	1	Total	C	N	O	0	0
			15	11	2	2		
3	C	1	Total	C	N	O	0	0
			15	11	2	2		
3	D	1	Total	C	N	O	0	0
			15	11	2	2		
3	E	1	Total	C	N	O	0	0
			15	11	2	2		
3	F	1	Total	C	N	O	0	0
			15	11	2	2		
3	G	1	Total	C	N	O	0	0
			15	11	2	2		
3	H	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 4 is D-TRYPTOPHAN (three-letter code: DTR) (formula: C₁₁H₁₂N₂O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	E	1	Total	C	N	O	0	0
			15	11	2	2		
4	F	1	Total	C	N	O	0	0
			15	11	2	2		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	17	Total	O	0	0
			17	17		
5	B	7	Total	O	0	0
			7	7		
5	C	12	Total	O	0	0
			12	12		
5	D	21	Total	O	0	0
			21	21		
5	E	16	Total	O	0	0
			16	16		
5	F	21	Total	O	0	0
			21	21		
5	G	3	Total	O	0	0
			3	3		
5	H	10	Total	O	0	0
			10	10		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: D-AMINO ACID OXIDASE

Chain A: 



• Molecule 1: D-AMINO ACID OXIDASE

Chain B: 



• Molecule 1: D-AMINO ACID OXIDASE




Age Group	Percentage
18-24	2%
25-34	45%
35-44	35%
45-54	10%
55-64	2%
65-74	1%
75-84	1%
85+	1%






• Molecule 1: D-AMINO ACID OXIDASE

Chain F:




• Molecule 1: D-AMINO ACID OXIDASE


Chain G:



• Molecule 1: D-AMINO ACID OXIDASE

Chain H:





4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	325.20Å 137.10Å 196.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.10 78.06 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (20.00-3.10) 98.0 (78.06-3.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.18 (at 3.13Å)	Xtriage
Refinement program	TNT V. 5-E	Depositor
R, R_{free}	0.236 , 0.250 0.247 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	53.4	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 81.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 78287 reflections	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	22441	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DTR, ITR, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.94	1/2796 (0.0%)	1.37	28/3808 (0.7%)
1	B	0.91	0/2796	1.34	27/3808 (0.7%)
1	C	0.90	0/2796	1.32	24/3808 (0.6%)
1	D	0.97	1/2796 (0.0%)	1.41	36/3808 (0.9%)
1	E	0.93	3/2796 (0.1%)	1.37	28/3808 (0.7%)
1	F	0.93	1/2796 (0.0%)	1.37	24/3808 (0.6%)
1	G	0.89	0/2796	1.34	21/3808 (0.6%)
1	H	0.92	2/2796 (0.1%)	1.34	24/3808 (0.6%)
All	All	0.92	8/22368 (0.0%)	1.36	212/30464 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	1	0
1	E	0	2
1	F	0	1
All	All	1	4

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	60	SER	CB-OG	8.68	1.53	1.42
1	E	23	TYR	CE1-CZ	-8.57	1.27	1.38
1	D	177	VAL	CB-CG1	-7.10	1.38	1.52
1	H	177	VAL	CB-CG1	-5.69	1.41	1.52
1	E	58	GLU	CG-CD	5.65	1.60	1.51

The worst 5 of 212 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	23	TYR	CG-CD1-CE1	-13.65	110.38	121.30
1	G	61	ASN	C-N-CD	-11.53	95.24	120.60
1	D	27	LEU	CA-CB-CG	10.98	140.55	115.30
1	C	194	LEU	CA-CB-CG	-8.93	94.75	115.30
1	D	194	LEU	CA-CB-CG	-8.93	94.75	115.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	23	TYR	CA

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	54	PRO	Mainchain
1	E	23	TYR	Sidechain
1	E	58	GLU	Sidechain
1	F	23	TYR	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2675	181	1
1	B	2720	0	2675	197	0
1	C	2720	0	2675	176	3
1	D	2720	0	2675	223	1
1	E	2720	0	2675	187	0
1	F	2720	0	2675	186	0
1	G	2720	0	2675	204	0
1	H	2720	0	2675	189	3
2	A	53	0	31	5	0
2	B	53	0	31	3	0
2	C	53	0	31	4	0
2	D	53	0	31	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	53	0	31	2	0
2	F	53	0	31	4	0
2	G	53	0	31	5	0
2	H	53	0	31	5	0
3	A	15	0	8	0	0
3	B	15	0	8	0	0
3	C	15	0	8	0	0
3	D	15	0	8	0	0
3	E	15	0	8	0	0
3	F	15	0	8	1	0
3	G	15	0	8	0	0
3	H	15	0	8	0	0
4	E	15	0	11	7	0
4	F	15	0	11	3	0
5	A	17	0	0	0	0
5	B	7	0	0	3	0
5	C	12	0	0	2	0
5	D	21	0	0	5	0
5	E	16	0	0	0	0
5	F	21	0	0	3	0
5	G	3	0	0	1	0
5	H	10	0	0	0	0
All	All	22441	0	21734	1493	4

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 36.

The worst 5 of 1493 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:223:ILE:HG13	1:D:224:TYR:CE1	1.52	1.42
1:D:223:ILE:HG13	1:D:224:TYR:CD1	1.69	1.27
1:A:253:ILE:HG12	1:F:42:PHE:CD1	1.73	1.23
1:E:22:ARG:HG3	1:E:22:ARG:HH11	1.09	1.14
1:F:223:ILE:HG12	1:F:224:TYR:CD1	1.85	1.11

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:123:ASP:OD1	1:H:162:ARG:NH2[4_565]	2.01	0.19
1:A:332:LYS:NZ	1:D:126:PRO:CG[7_545]	2.11	0.09
1:C:123:ASP:OD2	1:H:35:TYR:OH[4_565]	2.15	0.05

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:162:ARG:NH2	1:H:123:ASP:OD1[3.655]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/347 (97%)	310 (92%)	25 (7%)	2 (1%)	33	78
1	B	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	25	71
1	C	337/347 (97%)	312 (93%)	23 (7%)	2 (1%)	33	78
1	D	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	25	71
1	E	337/347 (97%)	309 (92%)	25 (7%)	3 (1%)	25	71
1	F	337/347 (97%)	311 (92%)	24 (7%)	2 (1%)	33	78
1	G	337/347 (97%)	306 (91%)	27 (8%)	4 (1%)	19	62
1	H	337/347 (97%)	310 (92%)	23 (7%)	4 (1%)	19	62
All	All	2696/2776 (97%)	2476 (92%)	197 (7%)	23 (1%)	25	71

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	28	GLN
1	E	28	GLN
1	A	28	GLN
1	B	24	HIS
1	G	222	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/298 (97%)	263 (91%)	27 (9%)	13	44
1	B	290/298 (97%)	265 (91%)	25 (9%)	15	51
1	C	290/298 (97%)	266 (92%)	24 (8%)	16	53
1	D	290/298 (97%)	265 (91%)	25 (9%)	15	51
1	E	290/298 (97%)	263 (91%)	27 (9%)	13	44
1	F	290/298 (97%)	263 (91%)	27 (9%)	13	44
1	G	290/298 (97%)	263 (91%)	27 (9%)	13	44
1	H	290/298 (97%)	268 (92%)	22 (8%)	19	58
All	All	2320/2384 (97%)	2116 (91%)	204 (9%)	14	49

5 of 204 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	216	THR
1	E	184	VAL
1	H	129	ARG
1	D	238	LEU
1	E	22	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 49 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	86	ASN
1	D	288	GLN
1	H	96	ASN
1	D	201	GLN
1	D	307	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.82	14 (16%)
3	ITR	A	349	-	16,16,16	1.00	1 (6%)	21,22,22	1.11	2 (9%)
2	FAD	B	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.82	14 (16%)
3	ITR	B	349	-	16,16,16	1.00	1 (6%)	21,22,22	1.11	2 (9%)
2	FAD	C	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.82	14 (16%)
3	ITR	C	349	-	16,16,16	1.00	1 (6%)	21,22,22	1.11	2 (9%)
2	FAD	D	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.83	14 (16%)
3	ITR	D	349	-	16,16,16	1.01	1 (6%)	21,22,22	1.11	2 (9%)
2	FAD	E	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.83	14 (16%)
3	ITR	E	349	-	16,16,16	1.00	1 (6%)	21,22,22	1.11	2 (9%)
4	DTR	E	350	-	16,16,16	1.23	1 (6%)	22,22,22	1.12	1 (4%)
2	FAD	F	348	-	58,58,58	0.91	1 (1%)	85,89,89	1.82	14 (16%)
3	ITR	F	349	-	16,16,16	1.01	1 (6%)	21,22,22	1.10	2 (9%)
4	DTR	F	350	-	16,16,16	1.48	1 (6%)	22,22,22	0.96	1 (4%)
2	FAD	G	348	-	58,58,58	0.90	1 (1%)	85,89,89	1.83	14 (16%)
3	ITR	G	349	-	16,16,16	1.01	1 (6%)	21,22,22	1.11	2 (9%)
2	FAD	H	348	-	58,58,58	0.91	1 (1%)	85,89,89	1.83	14 (16%)
3	ITR	H	349	-	16,16,16	1.00	1 (6%)	21,22,22	1.12	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	348	-	-	0/34/50/50	0/1/6/6
3	ITR	A	349	-	-	0/5/8/8	0/0/2/2
2	FAD	B	348	-	-	0/34/50/50	0/1/6/6
3	ITR	B	349	-	-	0/5/8/8	0/0/2/2
2	FAD	C	348	-	-	0/34/50/50	0/1/6/6
3	ITR	C	349	-	-	0/5/8/8	0/0/2/2
2	FAD	D	348	-	-	0/34/50/50	0/1/6/6
3	ITR	D	349	-	-	0/5/8/8	0/0/2/2
2	FAD	E	348	-	-	0/34/50/50	0/1/6/6
3	ITR	E	349	-	-	0/5/8/8	0/0/2/2
4	DTR	E	350	-	-	0/8/8/8	0/0/2/2
2	FAD	F	348	-	-	0/34/50/50	0/1/6/6
3	ITR	F	349	-	-	0/5/8/8	0/0/2/2
4	DTR	F	350	-	-	0/8/8/8	0/0/2/2
2	FAD	G	348	-	-	0/34/50/50	0/1/6/6
3	ITR	G	349	-	-	0/5/8/8	0/0/2/2
2	FAD	H	348	-	-	0/34/50/50	0/1/6/6
3	ITR	H	349	-	-	0/5/8/8	0/0/2/2

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	350	DTR	CB-CG	-4.91	1.44	1.51
4	E	350	DTR	CB-CG	-3.88	1.46	1.51
2	F	348	FAD	C6-C5X	-2.25	1.39	1.41
2	E	348	FAD	C6-C5X	-2.21	1.39	1.41
2	D	348	FAD	C6-C5X	-2.20	1.39	1.41

The worst 5 of 130 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	348	FAD	C2-N1-C10	6.91	121.94	114.98
2	G	348	FAD	C2-N1-C10	6.90	121.93	114.98
2	E	348	FAD	C2-N1-C10	6.88	121.91	114.98
2	D	348	FAD	C2-N1-C10	6.86	121.89	114.98
2	A	348	FAD	C2-N1-C10	6.85	121.88	114.98

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/347 (95%)	-0.02	0 100 100	18, 35, 71, 100	23 (6%)
1	B	328/347 (94%)	0.19	9 (2%) 52 8	18, 36, 77, 100	24 (7%)
1	C	331/347 (95%)	0.03	3 (0%) 81 25	18, 36, 82, 100	26 (7%)
1	D	332/347 (95%)	0.06	4 (1%) 75 20	12, 36, 69, 100	19 (5%)
1	E	331/347 (95%)	0.13	2 (0%) 86 36	18, 36, 78, 100	26 (7%)
1	F	329/347 (94%)	-0.03	1 (0%) 91 53	18, 36, 70, 100	19 (5%)
1	G	327/347 (94%)	0.51	13 (3%) 36 5	18, 36, 83, 100	26 (7%)
1	H	329/347 (94%)	0.05	2 (0%) 86 36	18, 36, 73, 100	22 (6%)
All	All	2637/2776 (94%)	0.12	34 (1%) 72 19	12, 36, 76, 100	185 (7%)

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	301	SER	6.4
1	D	299	GLY	4.0
1	G	336	GLU	3.3
1	G	297	ARG	3.2
1	D	301	SER	3.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	ITR	A	349	15/15	0.29	2.03	45,52,58,72	0
3	ITR	H	349	15/15	0.33	1.82	45,52,58,72	0
3	ITR	C	349	15/15	0.33	1.64	45,52,58,72	0
3	ITR	E	349	15/15	0.32	1.47	45,52,58,72	0
4	DTR	F	350	15/15	0.35	0.94	1,49,100,100	0
2	FAD	C	348	53/53	0.26	0.73	20,27,42,43	0
2	FAD	D	348	53/53	0.27	0.69	20,27,42,43	0
3	ITR	F	349	15/15	0.31	0.68	45,52,58,72	0
3	ITR	D	349	15/15	0.32	0.68	45,52,58,72	0
2	FAD	G	348	53/53	0.36	0.64	20,27,42,43	0
2	FAD	B	348	53/53	0.26	0.55	20,27,42,43	0
3	ITR	G	349	15/15	0.43	0.46	45,52,58,72	0
2	FAD	A	348	53/53	0.23	0.45	20,27,42,43	0
2	FAD	H	348	53/53	0.24	0.39	20,27,42,43	0
4	DTR	E	350	15/15	0.36	0.38	9,42,100,100	0
3	ITR	B	349	15/15	0.26	0.24	45,52,58,72	0
2	FAD	E	348	53/53	0.23	0.20	20,27,42,43	0
2	FAD	F	348	53/53	0.23	0.19	20,27,42,43	0

6.5 Other polymers ⓘ

There are no such residues in this entry.