



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 01:34 PM GMT

PDB ID : 1DFK
Title : NUCLEOTIDE-FREE SCALLOP MYOSIN S1-NEAR RIGOR STATE
Authors : Houdusse, A.; Szent-Gyorgyi, A.G.; Cohen, C.
Deposited on : 1999-11-19
Resolution : 4.20 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 4.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1259 (4.84-3.50)
Ramachandran outliers	78287	1192 (4.84-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	830	
2	Y	139	
3	Z	152	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5032 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	0	0	0
			3597	2140	728	729			

- Molecule 2 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	Y	139	Total	C	N	O	0	0	0
			688	409	139	140			

- Molecule 3 is a protein called MYOSIN HEAD.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Z	152	Total	C	N	O	0	0	0
			746	441	152	153			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Z	1	Total	Ca	0	0
			1	1		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: MYOSIN HEAD



• Molecule 2: MYOSIN HEAD



• Molecule 3: MYOSIN HEAD



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.50Å 52.10Å 164.60Å 90.00° 100.30° 90.00°	Depositor
Resolution (Å)	20.00 – 4.20	Depositor
% Data completeness (in resolution range)	95.0 (20.00-4.20)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.415 , 0.410	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5032	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.97	9/3587 (0.3%)	1.33	26/4978 (0.5%)
2	Y	0.42	0/687	0.80	2/954 (0.2%)
3	Z	0.58	1/745 (0.1%)	0.76	0/1031
All	All	0.86	10/5019 (0.2%)	1.20	28/6963 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	8

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	268	GLU	C-N	-31.26	0.62	1.34
1	A	705	LYS	C-N	23.07	1.74	1.33
1	A	706	GLY	C-N	19.97	1.79	1.34
1	A	683	ASP	C-N	-18.55	0.91	1.34
3	Z	7	GLU	C-N	11.26	1.59	1.34

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	515	ASP	O-C-N	-39.30	59.82	122.70
1	A	268	GLU	O-C-N	-35.48	65.94	122.70
1	A	268	GLU	C-N-CA	-24.70	59.95	121.70
1	A	268	GLU	CA-C-N	-24.51	63.29	117.20
1	A	515	ASP	CA-C-N	16.06	152.53	117.20

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	268	GLU	Mainchain
1	A	445	THR	Mainchain
1	A	515	ASP	Mainchain,Peptide
1	A	600	ASP	Mainchain
1	A	601	PRO	Mainchain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3597	0	1592	58	0
2	Y	688	0	307	17	0
3	Z	746	0	340	11	0
4	Z	1	0	0	0	0
All	All	5032	0	2239	85	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:705:LYS:C	1:A:706:GLY:N	1.74	1.40
1:A:706:GLY:C	1:A:707:PHE:N	1.80	1.35
1:A:584:ALA:HB1	1:A:693:CYS:HA	1.37	1.07
1:A:770:LEU:O	1:A:774:ARG:CB	2.10	0.99
1:A:507:TRP:O	1:A:508:GLU:C	2.07	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	708/830 (85%)	576 (81%)	95 (13%)	37 (5%)	3	41
2	Y	137/139 (99%)	96 (70%)	34 (25%)	7 (5%)	3	42
3	Z	150/152 (99%)	106 (71%)	33 (22%)	11 (7%)	2	31
All	All	995/1121 (89%)	778 (78%)	162 (16%)	55 (6%)	3	40

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	GLU
1	A	269	LYS
1	A	542	ALA
1	A	601	PRO
1	A	623	LYS

5.3.2 Protein sidechains ⓘ

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.