



# wwPDB X-ray Structure Validation Summary Report

(i)

Feb 28, 2014 – 12:38 AM GMT

PDB ID : 1DML  
Title : CRYSTAL STRUCTURE OF HERPES SIMPLEX UL42 BOUND TO THE C-TERMINUS OF HSV POL  
Authors : Zuccola, H.J.; Filman, D.J.; Coen, D.M.; Hogle, J.M.  
Deposited on : 1999-12-14  
Resolution : 2.70 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance (i)


The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1557 (2.70-2.70)
Clashscore	79885	1939 (2.70-2.70)
Ramachandran outliers	78287	1905 (2.70-2.70)
Sidechain outliers	78261	1905 (2.70-2.70)
RSRZ outliers	66119	1559 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 9414 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE PROCESSIVITY FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
		2059	1309	362	381	7				
1	C	272	Total	C	N	O	S	0	0	0
		2087	1325	368	387	7				
1	E	275	Total	C	N	O	S	0	0	0
		2108	1336	370	395	7				
1	G	275	Total	C	N	O	S	0	0	0
		2108	1336	370	395	7				

- Molecule 2 is a protein called DNA POLYMERASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	36	Total	C	N	O	S	0	0	0
		263	159	53	50	1				
2	D	36	Total	C	N	O	S	0	0	0
		263	159	53	50	1				
2	F	36	Total	C	N	O	S	0	0	0
		263	159	53	50	1				
2	H	36	Total	C	N	O	S	0	0	0
		263	159	53	50	1				

### 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR

## Chain A



- Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR

### Chain C



- Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR


### Chain E





- Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR

Chain G: 



- Molecule 2: DNA POLYMERASE

Chain B:



- Molecule 2: DNA POLYMERASE

Chain D:




- Molecule 2: DNA POLYMERASE

Chain F:



- Molecule 2: DNA POLYMERASE

Chain H:



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.32 Å    100.07 Å    129.53 Å 90.00°    100.62°    90.00°	Depositor
Resolution (Å)	12.00 – 2.70 29.87 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.6 (12.00-2.70) 99.9 (29.87-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	5.21 (at 2.51 Å)	Xtriage
Refinement program	CNS 0.5	Depositor
$R$ , $R_{free}$	0.230 , 0.281 0.190 , 0.224	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 17.7	EDS
Estimated twinning fraction	0.000 for h,-k,-h-1	Xtriage
L-test for twinning	$<  L  > = 0.49$ , $< L^2 > = 0.32$	Xtriage
Outliers	2 of 47173 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9414	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.37 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7625e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.41	0/2097	0.71	2/2851 (0.1%)
1	C	0.41	0/2125	0.70	2/2889 (0.1%)
1	E	0.39	0/2146	0.65	0/2918
1	G	0.38	0/2146	0.64	0/2918
2	B	0.41	0/265	0.56	0/354
2	D	0.40	0/265	0.56	0/354
2	F	0.38	0/265	0.52	0/354
2	H	0.39	0/265	0.55	0/354
All	All	0.40	0/9574	0.66	4/12992 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	155	GLU	N-CA-C	-6.85	92.49	111.00
1	C	155	GLU	N-CA-C	-6.85	92.50	111.00
1	A	136	LEU	CA-CB-CG	5.30	127.49	115.30
1	C	136	LEU	CA-CB-CG	5.27	127.43	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2059	0	2104	100	0
1	C	2087	0	2130	96	0
1	E	2108	0	2150	104	0
1	G	2108	0	2150	104	0
2	B	263	0	256	18	0
2	D	263	0	256	20	0
2	F	263	0	256	25	0
2	H	263	0	256	22	0
All	All	9414	0	9558	444	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 23.

The worst 5 of 444 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:G:121:ARG:HH11	1:G:143:THR:HG22	1.18	1.06
1:A:267:VAL:HG11	1:A:315:LEU:HD13	1.34	1.05
1:G:171:GLN:HA	2:H:1233:THR:HG21	1.39	1.04
1:C:149:ALA:HB3	1:C:152:LEU:HD22	1.41	0.99
1:E:286:GLY:HA2	1:E:304:THR:HG23	1.44	0.99

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/319 (82%)	237 (90%)	18 (7%)	8 (3%)	7 15
1	C	268/319 (84%)	240 (90%)	16 (6%)	12 (4%)	4 7
1	E	271/319 (85%)	243 (90%)	21 (8%)	7 (3%)	8 20
1	G	271/319 (85%)	238 (88%)	28 (10%)	5 (2%)	13 31
2	B	34/36 (94%)	33 (97%)	0	1 (3%)	7 16
2	D	34/36 (94%)	33 (97%)	0	1 (3%)	7 16

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	F	34/36 (94%)	31 (91%)	2 (6%)	1 (3%)	7 16
2	H	34/36 (94%)	30 (88%)	3 (9%)	1 (3%)	7 16
All	All	1209/1420 (85%)	1085 (90%)	88 (7%)	36 (3%)	7 15

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	LEU
1	A	223	ALA
1	A	270	ASP
1	A	272	CYS
2	B	1216	ALA

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	228/266 (86%)	214 (94%)	14 (6%)	26 54
1	C	229/266 (86%)	216 (94%)	13 (6%)	29 58
1	E	235/266 (88%)	224 (95%)	11 (5%)	36 69
1	G	235/266 (88%)	222 (94%)	13 (6%)	30 60
2	B	22/22 (100%)	20 (91%)	2 (9%)	14 30
2	D	22/22 (100%)	20 (91%)	2 (9%)	14 30
2	F	22/22 (100%)	19 (86%)	3 (14%)	5 13
2	H	22/22 (100%)	19 (86%)	3 (14%)	5 13
All	All	1015/1152 (88%)	954 (94%)	61 (6%)	27 56

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	204	LEU
1	E	71	THR
1	G	287	THR
1	C	263	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	D	1234	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	70	ASN
1	E	207	ASN
1	G	171	GLN
1	E	36	GLN
1	G	207	ASN

### 5.3.3 RNA (i)

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	267/319 (83%)	-0.12	12 (4%) 32 36	5, 19, 64, 85	0
1	C	272/319 (85%)	-0.02	13 (4%) 29 33	5, 23, 67, 87	0
1	E	275/319 (86%)	0.01	10 (3%) 41 46	8, 29, 69, 90	0
1	G	275/319 (86%)	0.20	15 (5%) 24 26	8, 33, 76, 91	0
2	B	36/36 (100%)	-0.25	0 100 100	9, 23, 45, 69	0
2	D	36/36 (100%)	-0.12	1 (2%) 50 56	11, 23, 48, 71	0
2	F	36/36 (100%)	0.26	1 (2%) 50 56	11, 46, 68, 82	0
2	H	36/36 (100%)	0.43	3 (8%) 11 12	15, 48, 71, 86	0
All	All	1233/1420 (86%)	0.02	55 (4%) 32 36	5, 27, 69, 91	0

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	144	THR	7.5
1	G	146	ASP	7.0
1	C	152	LEU	6.0
1	E	146	ASP	5.5
2	H	1200	ASP	5.1

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

## 6.4 Ligands (i)

There are no ligands in this entry.

## 6.5 Other polymers (i)

There are no such residues in this entry.