



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 08:34 AM GMT

PDB ID : 3DP9
Title : Crystal structure of Vibrio cholerae 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase (MTAN) complexed with butylthio-DADMe-ImmucillinA
Authors : Ho, M.; Gutierrez, J.A.; Crowder, T.; Rinaldo-Matthis, A.; Almo, S.C.; Schramm, V.L.
Deposited on : 2008-07-07
Resolution : 2.30 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2929 (2.30-2.30)
Clashscore	79885	3679 (2.30-2.30)
Ramachandran outliers	78287	3642 (2.30-2.30)
Sidechain outliers	78261	3641 (2.30-2.30)
RSRZ outliers	66119	2930 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	231	
1	C	231	

2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 3523 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MTA/SAH nucleosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	0	0
			1701	1077	284	330	10			
1	C	230	Total	C	N	O	S	0	0	0
			1709	1082	285	331	11			

- Molecule 2 is (3R,4S)-1-[(4-AMINO-5H-PYRROLO[3,2-D]PYRIMIDIN-7-YL)METHYL]-4-[(BUTYLSULFANYL)METHYL]PYRROLIDIN-3-OL (three-letter code: BIG) (formula: C₁₆H₂₅N₅OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			23	16	5	1	1		
2	C	1	Total	C	N	O	S	0	0
			23	16	5	1	1		

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total I 1 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	38	Total O 38 38	0	0
4	C	28	Total O 28 28	0	0

3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MTA/SAH nucleosidase

Chain A:



- Molecule 1: MTA/SAH nucleosidase

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.74Å 54.33Å 69.79Å 90.00° 115.35° 90.00°	Depositor
Resolution (Å)	31.53 – 2.30 31.53 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.7 (31.53-2.30) 90.7 (31.53-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.264 0.198 , 0.261	Depositor DCC
R_{free} test set	952 reflections (5.36%)	DCC
Wilson B-factor (Å ²)	42.8	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.0	EDS
Estimated twinning fraction	0.023 for l,-k,h	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 18707 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3523	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BIG, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1726	0.61	0/2341
1	C	0.54	1/1734 (0.1%)	0.61	1/2351 (0.0%)
All	All	0.53	1/3460 (0.0%)	0.61	1/4692 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	137	PRO	N-CD	9.00	1.60	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	57	LEU	CA-CB-CG	5.28	127.44	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1701	0	1707	59	0
1	C	1709	0	1719	46	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	23	0	25	1	0
2	C	23	0	25	1	0
3	C	1	0	0	3	0
4	A	38	0	0	1	0
4	C	28	0	0	0	0
All	All	3523	0	3476	99	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

All (99) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:148:THR:HG21	1:A:177:SER:H	1.05	1.15
1:A:115:GLN:HE22	1:A:185:GLN:NE2	1.52	1.05
1:C:143:ARG:HG2	1:C:143:ARG:HH11	1.17	1.04
1:A:143:ARG:HG2	1:A:143:ARG:HH11	1.24	0.99
1:C:158:ARG:HG3	1:C:158:ARG:HH11	1.33	0.94
1:A:148:THR:HG21	1:A:177:SER:N	1.83	0.93
1:A:115:GLN:NE2	1:A:185:GLN:HE21	1.68	0.89
1:C:158:ARG:CG	1:C:158:ARG:HH11	1.84	0.89
1:A:115:GLN:NE2	1:A:185:GLN:NE2	2.21	0.88
1:A:86:VAL:HG22	1:A:214:LEU:HD23	1.57	0.86
1:A:38:ILE:CG2	1:A:227:VAL:HG11	2.06	0.85
1:A:166:PHE:O	1:A:169:VAL:HG23	1.77	0.85
1:A:159:GLN:OE1	1:A:163:ARG:NH1	2.12	0.83
1:A:115:GLN:HE22	1:A:185:GLN:HE21	0.85	0.82
1:C:166:PHE:O	1:C:169:VAL:HG23	1.82	0.79
1:A:148:THR:CG2	1:A:177:SER:H	1.92	0.78
1:C:143:ARG:HG2	1:C:143:ARG:NH1	1.92	0.78
1:C:143:ARG:HH11	1:C:143:ARG:CG	1.96	0.78
1:A:148:THR:HG23	1:A:177:SER:HB3	1.66	0.78
1:A:143:ARG:HG2	1:A:143:ARG:NH1	1.99	0.76
1:C:73:ASN:HD21	1:C:194:ARG:HH21	1.31	0.76
1:C:158:ARG:HG3	1:C:158:ARG:NH1	2.03	0.73
1:C:178:ALA:O	1:C:182:THR:HG23	1.89	0.72
1:A:38:ILE:HG23	1:A:227:VAL:HG11	1.71	0.72
1:A:57:LEU:HD12	1:C:182:THR:HB	1.73	0.69
1:A:86:VAL:HG21	1:A:211:PHE:HB3	1.78	0.65
1:C:124:LYS:O	1:C:128:VAL:HG23	1.97	0.65
1:C:38:ILE:HG23	1:C:227:VAL:HG11	1.79	0.65
1:C:225:LYS:O	1:C:228:GLU:HB3	1.97	0.64
1:A:38:ILE:HG23	1:A:38:ILE:O	1.95	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:19:LEU:HB3	1:A:38:ILE:HD11	1.80	0.63
1:A:229:LEU:O	4:A:311:HOH:O	2.15	0.62
1:A:136:LEU:HB3	1:A:139:THR:CG2	2.30	0.62
1:A:38:ILE:HG21	1:A:227:VAL:HG11	1.82	0.61
1:A:135:GLN:C	1:A:136:LEU:HD13	2.22	0.60
1:A:2:LYS:N	1:A:69:ASP:OD2	2.35	0.59
1:A:112:MET:CE	1:C:50:ILE:O	2.51	0.58
1:C:135:GLN:HG2	1:C:225:LYS:HE2	1.85	0.58
1:A:86:VAL:HG21	1:A:211:PHE:CG	2.39	0.58
1:C:166:PHE:O	1:C:169:VAL:CG2	2.51	0.58
1:A:173:GLU:HA	2:A:301:BIG:N3	2.19	0.57
1:C:213:PRO:O	1:C:217:LYS:HG3	2.05	0.57
1:A:225:LYS:O	1:A:229:LEU:HD13	2.05	0.56
1:A:147:CYS:SG	1:A:169:VAL:HG22	2.45	0.56
1:A:112:MET:HE1	1:C:50:ILE:O	2.06	0.55
1:A:148:THR:HG23	1:A:177:SER:CB	2.35	0.54
1:C:92:SER:HB3	1:C:146:ILE:HD12	1.90	0.54
1:C:173:GLU:HA	2:C:301:BIG:N3	2.23	0.54
1:A:213:PRO:O	1:A:217:LYS:HG3	2.07	0.53
1:C:139:THR:CG2	3:C:300:IOD:I	3.28	0.52
1:C:178:ALA:O	1:C:182:THR:CG2	2.57	0.52
1:A:20:ILE:HD11	1:A:23:VAL:HG23	1.90	0.52
1:A:38:ILE:CG2	1:A:38:ILE:O	2.57	0.52
1:C:38:ILE:CG2	1:C:227:VAL:HG11	2.41	0.51
1:C:137:PRO:O	3:C:300:IOD:I	2.98	0.51
1:A:92:SER:HB3	1:A:146:ILE:HD12	1.93	0.51
1:C:158:ARG:CG	1:C:158:ARG:NH1	2.55	0.50
1:A:73:ASN:HB3	1:A:192:VAL:HG22	1.93	0.50
1:C:139:THR:HG22	3:C:300:IOD:I	2.82	0.50
1:C:86:VAL:HG11	1:C:211:PHE:CG	2.46	0.50
1:A:89:VAL:HG21	1:A:222:MET:HG3	1.93	0.49
1:A:86:VAL:HG21	1:A:211:PHE:CB	2.40	0.49
1:C:98:HIS:HB3	1:C:147:CYS:HB3	1.94	0.49
1:C:38:ILE:HD11	1:C:224:LEU:HD13	1.95	0.48
1:A:20:ILE:HD12	1:A:36:GLY:HA3	1.96	0.47
1:C:38:ILE:HD11	1:C:224:LEU:CD1	2.45	0.47
1:C:225:LYS:O	1:C:229:LEU:HD12	2.15	0.47
1:C:181:GLN:HE21	1:C:185:GLN:NE2	2.12	0.47
1:A:124:LYS:O	1:A:128:VAL:HG23	2.14	0.47
1:A:98:HIS:HB3	1:A:147:CYS:HB3	1.96	0.47
1:A:57:LEU:HD23	1:A:57:LEU:C	2.36	0.47
1:A:86:VAL:CG2	1:A:211:PHE:HB3	2.44	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:162:ILE:HG22	1:C:169:VAL:HG21	1.97	0.46
1:A:47:GLN:O	1:A:47:GLN:HG3	2.14	0.46
1:C:73:ASN:ND2	1:C:194:ARG:HH21	2.08	0.45
1:A:112:MET:HE2	1:C:50:ILE:O	2.16	0.45
1:C:24:GLN:HB3	1:C:24:GLN:HE21	1.65	0.45
1:A:136:LEU:N	1:A:136:LEU:HD13	2.32	0.45
1:A:229:LEU:N	1:A:229:LEU:HD12	2.32	0.45
1:C:13:VAL:HB	1:C:45:LEU:HD21	1.98	0.45
1:C:158:ARG:HG2	1:C:158:ARG:HH11	1.77	0.45
1:A:86:VAL:CG2	1:A:214:LEU:HD23	2.40	0.45
1:C:225:LYS:O	1:C:229:LEU:CD1	2.65	0.44
1:A:198:ASP:HB2	1:A:204:SER:HB3	2.00	0.44
1:A:206:LEU:HA	1:A:209:GLU:HB3	2.00	0.44
1:A:80:PHE:CZ	1:A:159:GLN:HG3	2.53	0.43
1:C:159:GLN:OE1	1:C:163:ARG:NH1	2.34	0.43
1:A:100:ALA:HA	1:C:150:ASP:O	2.19	0.43
1:A:90:VAL:CG2	1:A:170:VAL:HG23	2.49	0.43
1:A:92:SER:HB3	1:A:192:VAL:HB	2.01	0.42
1:C:212:LEU:N	1:C:213:PRO:CD	2.82	0.42
1:C:137:PRO:O	1:C:138:ASN:HB2	2.20	0.42
1:C:73:ASN:HB3	1:C:192:VAL:HG22	2.00	0.42
1:A:10:GLN:HA	1:A:47:GLN:HE22	1.85	0.41
1:C:1:MET:O	1:C:42:ASP:HB2	2.21	0.41
1:A:202:LYS:HB2	1:A:202:LYS:HE3	1.98	0.41
1:A:185:GLN:OE1	1:C:57:LEU:HD13	2.22	0.40
1:A:159:GLN:NE2	1:A:200:ALA:O	2.48	0.40
1:A:166:PHE:HB2	1:A:169:VAL:CG2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/231 (98%)	211 (93%)	16 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	228/231 (99%)	217 (95%)	11 (5%)	0	100	100
All	All	455/462 (98%)	428 (94%)	27 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/184 (99%)	164 (90%)	18 (10%)	11	13
1	C	183/184 (100%)	170 (93%)	13 (7%)	21	26
All	All	365/368 (99%)	334 (92%)	31 (8%)	15	18

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ASP
1	A	20	ILE
1	A	27	ASN
1	A	38	ILE
1	A	47	GLN
1	A	57	LEU
1	A	130	GLU
1	A	135	GLN
1	A	136	LEU
1	A	139	THR
1	A	143	ARG
1	A	148	THR
1	A	172	VAL
1	A	173	GLU
1	A	203	GLU
1	A	204	SER
1	A	206	LEU
1	A	230	LEU
1	C	24	GLN
1	C	32	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	57	LEU
1	C	86	VAL
1	C	90	VAL
1	C	123	GLU
1	C	135	GLN
1	C	139	THR
1	C	143	ARG
1	C	158	ARG
1	C	173	GLU
1	C	182	THR
1	C	199	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	47	GLN
1	A	65	GLN
1	A	115	GLN
1	A	165	HIS
1	C	24	GLN
1	C	28	GLN
1	C	47	GLN
1	C	65	GLN
1	C	73	ASN
1	C	115	GLN
1	C	164	GLN
1	C	185	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BIG	A	301	-	25,25,25	0.81	0	32,34,34	2.19	9 (28%)
2	BIG	C	301	-	25,25,25	0.93	1 (4%)	32,34,34	2.16	9 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BIG	A	301	-	-	0/9/22/22	0/1/3/3
2	BIG	C	301	-	-	0/9/22/22	0/1/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	BIG	C5'-S5'	-2.22	1.76	1.81

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	BIG	N3-C2-N1	-6.88	122.96	128.71
2	C	301	BIG	N3-C2-N1	-6.65	123.15	128.71
2	C	301	BIG	C2-N3-C4	4.38	118.66	115.13
2	A	301	BIG	C9-C10-N1'	4.10	119.34	113.93
2	C	301	BIG	C9-C10-N1'	3.92	119.11	113.93
2	A	301	BIG	C5'-C4'-C1'	-3.92	107.08	113.51
2	A	301	BIG	C2-N3-C4	3.83	118.22	115.13
2	A	301	BIG	C10-N1'-C2'	-3.28	108.11	113.76
2	C	301	BIG	C5-C4-N3	-3.28	121.85	125.25
2	A	301	BIG	C2-N1-C6	2.80	123.83	118.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	BIG	C4'-C5'-S5'	-2.77	109.59	114.71
2	C	301	BIG	C2-N1-C6	2.48	123.24	118.77
2	A	301	BIG	C2'-C3'-C4'	2.44	105.13	103.47
2	C	301	BIG	C9-C8-N7	2.38	111.97	107.94
2	C	301	BIG	C2'-N1'-C1'	2.36	108.63	104.58
2	A	301	BIG	C5-C4-N3	-2.28	122.88	125.25
2	C	301	BIG	C5'-C4'-C1'	-2.22	109.88	113.51
2	A	301	BIG	C2'-N1'-C1'	2.03	108.05	104.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	229/231 (99%)	0.47	16 (6%) 16 23	33, 45, 62, 68	0
1	C	230/231 (99%)	0.45	24 (10%) 7 11	32, 45, 58, 60	0
All	All	459/462 (99%)	0.46	40 (8%) 10 16	32, 45, 60, 68	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	22	ASP	4.3
1	C	134	ALA	4.3
1	C	53	VAL	4.0
1	A	178	ALA	3.9
1	C	209	GLU	3.1
1	A	56	ALA	3.0
1	C	52	LYS	2.9
1	C	206	LEU	2.8
1	C	24	GLN	2.8
1	C	37	GLN	2.7
1	A	53	VAL	2.7
1	C	39	GLN	2.7
1	A	38	ILE	2.7
1	A	177	SER	2.6
1	C	23	VAL	2.6
1	A	52	LYS	2.5
1	C	178	ALA	2.5
1	A	55	ALA	2.5
1	A	39	GLN	2.5
1	C	51	GLY	2.5
1	C	8	ALA	2.4
1	C	7	GLY	2.4
1	C	49	GLY	2.4
1	C	75	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	42	ASP	2.3
1	C	41	VAL	2.3
1	A	174	MET	2.3
1	A	22	ASP	2.3
1	C	54	SER	2.2
1	A	179	ILE	2.2
1	A	50	ILE	2.2
1	C	149	GLY	2.2
1	A	54	SER	2.2
1	C	148	THR	2.1
1	A	149	GLY	2.1
1	C	202	LYS	2.1
1	C	175	GLU	2.1
1	C	56	ALA	2.1
1	C	124	LYS	2.0
1	A	148	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	BIG	A	301	23/23	0.12	-0.45	23,28,36,38	0
2	BIG	C	301	23/23	0.12	-0.61	32,33,41,43	0
3	IOD	C	300	1/1	0.03	-8.85	67,67,67,67	0

6.5 Other polymers ⓘ

There are no such residues in this entry.