



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 04:32 PM GMT

PDB ID : 4DPL  
Title : Structure of malonyl-coenzyme A reductase from crenarchaeota in complex with NadP  
Authors : Demmer, U.; Warkentin, E; Srivastava, A.; Kockelkorn, D.; Fuchs, G.; Ermler, U.  
Deposited on : 2012-02-13  
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	359	
1	B	359	
1	C	359	
1	D	359	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	NAP	B	401	-	X
3	UNL	A	402	-	X
3	UNL	B	402	-	X
3	UNL	C	402	-	X
3	UNL	D	402	-	X

## 2 Entry composition i


There are 4 unique types of molecules in this entry. The entry contains 12082 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malonyl-CoA/succinyl-CoA reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	354	Total	C	N	O	S	0	1	0
			2746	1761	466	510	9			
1	B	354	Total	C	N	O	S	0	2	0
			2749	1763	466	511	9			
1	C	353	Total	C	N	O	S	0	1	0
			2738	1757	463	510	8			
1	D	354	Total	C	N	O	S	0	1	0
			2746	1761	466	510	9			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDEPHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is UNKNOWN LIGAND (three-letter code: UNL) (formula: ).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	233	Total	O	0	0
			233	233		
4	B	225	Total	O	0	0
			225	225		
4	C	245	Total	O	0	0
			245	245		
4	D	184	Total	O	0	0
			184	184		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Malonyl-CoA/succinyl-CoA reductase

Chain A:




- Molecule 1: Malonyl-CoA/succinyl-CoA reductase

Chain B:




- Molecule 1: Malonyl-CoA/succinyl-CoA reductase

Chain C:



- Molecule 1: Malonyl-CoA/succinyl-CoA reductase

Chain D:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	94.58Å 128.80Å 140.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.90 19.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-1.90) 99.5 (19.95-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.56 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.6.0095	Depositor
R, $R_{free}$	0.163 , 0.194 0.163 , 0.194	Depositor DCC
$R_{free}$ test set	6757 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.3	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 38.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 134386 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12082	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, UNL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.01	1/2810 (0.0%)	0.90	5/3816 (0.1%)
1	B	0.99	1/2816 (0.0%)	0.86	2/3824 (0.1%)
1	C	0.98	0/2802	0.89	5/3806 (0.1%)
1	D	0.93	3/2810 (0.1%)	0.85	5/3816 (0.1%)
All	All	0.98	5/11238 (0.0%)	0.87	17/15262 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	138	ARG	CZ-NH1	8.54	1.44	1.33
1	D	135	ASP	C-O	7.65	1.37	1.23
1	A	251	VAL	CB-CG2	5.78	1.65	1.52
1	D	314	VAL	CB-CG1	5.69	1.64	1.52
1	B	302	PHE	CE1-CZ	5.20	1.47	1.37

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	241	ARG	NE-CZ-NH1	11.22	125.91	120.30
1	C	241	ARG	NE-CZ-NH2	-9.37	115.61	120.30
1	A	241	ARG	NE-CZ-NH2	-8.90	115.85	120.30
1	A	241	ARG	NE-CZ-NH1	8.87	124.73	120.30
1	D	318	ARG	NE-CZ-NH1	7.31	123.95	120.30
1	B	53	ARG	NE-CZ-NH1	7.02	123.81	120.30
1	A	325	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	D	138	ARG	NE-CZ-NH2	6.81	123.70	120.30
1	C	18	LEU	CA-CB-CG	-6.38	100.62	115.30
1	B	18	LEU	CA-CB-CG	-6.29	100.83	115.30
1	A	18	LEU	CA-CB-CG	-5.77	102.03	115.30
1	C	207	ASP	CB-CG-OD1	5.50	123.25	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	18	LEU	CA-CB-CG	-5.39	102.90	115.30
1	C	303	ASP	CB-CG-OD2	5.34	123.11	118.30
1	D	172	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	304	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	D	241	ARG	NE-CZ-NH1	5.01	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2746	0	0	5	0
1	B	2749	0	0	7	0
1	C	2738	0	0	9	0
1	D	2746	0	0	5	0
2	A	48	0	0	0	0
2	B	48	0	0	1	0
2	C	48	0	0	1	0
2	D	48	0	0	0	0
3	A	6	0	0	1	0
3	B	6	0	0	1	0
3	C	6	0	0	3	0
3	D	6	0	0	0	0
4	A	233	0	0	1	0
4	B	225	0	0	3	0
4	C	245	0	0	4	0
4	D	184	0	0	2	0
All	All	12082	0	0	27	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 2.

All (27) close contacts within the same asymmetric unit are listed below.



Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:266:LYS:NZ	1:B:293:ASN:ND2	2.31	0.78
1:B:212:LYS:NZ	4:B:723:HOH:O	2.26	0.69
1:C:258:GLU:OE1	4:C:610:HOH:O	2.10	0.69
1:C:34:LYS:NZ	4:C:532:HOH:O	2.28	0.67
1:C:241:ARG:NH1	3:C:402:UNL:O41	2.30	0.64
1:A:266:LYS:NZ	1:A:293:ASN:ND2	2.47	0.63
1:C:266:LYS:NZ	1:C:293:ASN:ND2	2.50	0.60
1:C:153:CYS:SG	3:C:402:UNL:C1	2.91	0.56
1:C:43:SER:OG	2:C:401:NAP:O1X	2.28	0.52
1:A:273:ARG:NH2	1:A:278:ASP:OD1	2.43	0.52
1:D:266:LYS:NZ	1:D:293:ASN:ND2	2.58	0.51
1:D:275:GLU:N	1:D:276:PRO:CD	2.75	0.50
1:C:153:CYS:SG	3:C:402:UNL:O	2.71	0.49
2:B:401:NAP:C5N	3:B:402:UNL:O	2.60	0.49
1:B:266:LYS:NZ	4:B:693:HOH:O	2.47	0.48
1:A:262:ALA:N	1:A:321:GLN:NE2	2.64	0.46
1:D:89:GLN:NE2	4:D:643:HOH:O	2.49	0.45
1:B:63:GLU:OE1	1:B:63:GLU:N	2.50	0.45
1:C:89:GLN:NE2	4:C:615:HOH:O	2.50	0.43
1:A:241:ARG:NH1	3:A:402:UNL:O41	2.52	0.43
1:D:178:THR:OG1	1:D:180:GLN:NE2	2.53	0.42
1:D:275:GLU:N	4:D:634:HOH:O	2.52	0.42
1:C:72:THR:N	4:C:656:HOH:O	2.53	0.41
1:A:321:GLN:NE2	4:A:679:HOH:O	2.53	0.41
1:B:30:HIS:NE2	1:B:349:GLU:OE2	2.54	0.41
1:B:320:LYS:NZ	4:B:511:HOH:O	2.54	0.41
1:B:178:THR:OG1	1:B:180:GLN:NE2	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	353/359 (98%)	344 (98%)	7 (2%)	2 (1%)	33 19
1	B	354/359 (99%)	343 (97%)	9 (2%)	2 (1%)	33 19

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	352/359 (98%)	341 (97%)	8 (2%)	3 (1%)	25	10
1	D	353/359 (98%)	340 (96%)	11 (3%)	2 (1%)	33	19
All	All	1412/1436 (98%)	1368 (97%)	35 (2%)	9 (1%)	33	19

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	86	PRO
1	B	86	PRO
1	C	86	PRO
1	D	86	PRO
1	C	340	ALA
1	D	337	VAL
1	B	337	VAL
1	A	337	VAL
1	C	337	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/302 (99%)	277 (93%)	21 (7%)	21	10
1	B	299/302 (99%)	278 (93%)	21 (7%)	21	10
1	C	297/302 (98%)	282 (95%)	15 (5%)	33	19
1	D	298/302 (99%)	280 (94%)	18 (6%)	27	13
All	All	1192/1208 (99%)	1117 (94%)	75 (6%)	26	12

All (75) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ARG
1	A	34	LYS
1	A	41	LYS
1	A	55	GLN
1	A	62	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	64	ILE
1	A	75	LYS
1	A	76	LEU
1	A	79	ASP
1	A	87	LEU
1	A	89	GLN
1	A	101	LYS
1	A	112	ASP
1	A	136	GLU
1	A	140	ARG
1	A	143	TRP
1	A	222	LYS
1	A	257	LYS
1	A	279	LEU
1	A	329	LEU
1	A	359	LYS
1	B	34	LYS
1	B	43	SER
1	B	46	LYS
1	B	55	GLN
1	B	62	LYS
1	B	68	GLU
1	B	76	LEU
1	B	79	ASP
1	B	87	LEU
1	B	89	GLN
1	B	101	LYS
1	B	129	HIS
1	B	140	ARG
1	B	143	TRP
1	B	144	LYS
1	B	167	LYS
1	B	222	LYS
1	B	229	LYS
1	B	257	LYS
1	B	358	GLU
1	B	359	LYS
1	C	55	GLN
1	C	59[A]	GLN
1	C	59[B]	GLN
1	C	76	LEU
1	C	79	ASP

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	87	LEU
1	C	89	GLN
1	C	101	LYS
1	C	136	GLU
1	C	143	TRP
1	C	222	LYS
1	C	264	LYS
1	C	279	LEU
1	C	280	LYS
1	C	329	LEU
1	D	41	LYS
1	D	55	GLN
1	D	62	LYS
1	D	74	PRO
1	D	75	LYS
1	D	76	LEU
1	D	79	ASP
1	D	87	LEU
1	D	98	GLN
1	D	101	LYS
1	D	112	ASP
1	D	129	HIS
1	D	140	ARG
1	D	143	TRP
1	D	144	LYS
1	D	222	LYS
1	D	279	LEU
1	D	280	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are unknown - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAP	A	401	-	52,52,52	1.55	8 (15%)	80,80,80	1.87	13 (16%)
2	NAP	B	401	-	52,52,52	1.63	9 (17%)	80,80,80	1.73	13 (16%)
2	NAP	C	401	-	52,52,52	1.67	9 (17%)	80,80,80	1.73	11 (13%)
2	NAP	D	401	-	52,52,52	1.40	7 (13%)	80,80,80	1.69	11 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	401	-	-	0/35/67/67	0/3/5/5
2	NAP	B	401	-	-	0/35/67/67	0/3/5/5
2	NAP	C	401	-	-	0/35/67/67	0/3/5/5
2	NAP	D	401	-	-	0/35/67/67	0/3/5/5

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NAP	C3N-C7N	-6.21	1.39	1.50
2	C	401	NAP	C3N-C7N	-5.72	1.40	1.50
2	B	401	NAP	C3N-C7N	-5.43	1.41	1.50
2	D	401	NAP	C3N-C7N	-4.90	1.42	1.50
2	B	401	NAP	C2N-N1N	4.38	1.40	1.35
2	C	401	NAP	C2N-N1N	4.27	1.40	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	401	NAP	C2N-N1N	3.80	1.40	1.35
2	C	401	NAP	PN-O3	3.66	1.67	1.60
2	C	401	NAP	PA-O3	3.64	1.66	1.59
2	A	401	NAP	C2N-N1N	3.34	1.39	1.35
2	B	401	NAP	PN-O2N	3.21	1.56	1.48
2	C	401	NAP	C4A-N9A	-3.16	1.33	1.37
2	B	401	NAP	PN-O3	3.08	1.66	1.60
2	C	401	NAP	PN-O2N	3.08	1.55	1.48
2	A	401	NAP	PA-O3	3.07	1.65	1.59
2	A	401	NAP	O4B-C1B	3.05	1.46	1.41
2	B	401	NAP	PA-O3	3.03	1.65	1.59
2	A	401	NAP	O4D-C1D	2.99	1.45	1.41
2	D	401	NAP	PN-O3	2.82	1.66	1.60
2	C	401	NAP	O4B-C1B	2.82	1.45	1.41
2	C	401	NAP	O4D-C1D	2.78	1.45	1.41
2	D	401	NAP	PN-O2N	2.70	1.55	1.48
2	A	401	NAP	PN-O3	2.60	1.65	1.60
2	B	401	NAP	O4B-C1B	2.56	1.45	1.41
2	B	401	NAP	PN-O1N	2.45	1.54	1.48
2	B	401	NAP	O4D-C1D	2.45	1.45	1.41
2	A	401	NAP	PN-O2N	2.38	1.54	1.48
2	C	401	NAP	PN-O1N	2.37	1.54	1.48
2	D	401	NAP	C4A-N9A	-2.21	1.34	1.37
2	A	401	NAP	PN-O1N	2.18	1.54	1.48
2	D	401	NAP	PA-O3	2.15	1.63	1.59
2	D	401	NAP	O4B-C1B	2.11	1.44	1.41
2	B	401	NAP	C2D-C1D	2.07	1.56	1.53

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	NAP	N3A-C2A-N1A	-10.51	119.92	128.71
2	A	401	NAP	N3A-C2A-N1A	-9.52	120.75	128.71
2	D	401	NAP	N3A-C2A-N1A	-9.18	121.03	128.71
2	B	401	NAP	N3A-C2A-N1A	-8.41	121.68	128.71
2	A	401	NAP	O4D-C1D-N1N	7.14	115.25	107.95
2	B	401	NAP	N3A-C4A-N9A	4.26	133.13	125.43
2	B	401	NAP	O4D-C1D-N1N	4.26	112.31	107.95
2	A	401	NAP	N3A-C4A-N9A	4.26	133.12	125.43
2	D	401	NAP	N3A-C4A-N9A	4.20	133.02	125.43
2	D	401	NAP	O4B-C1B-N9A	-3.70	105.00	108.44
2	C	401	NAP	N3A-C4A-N9A	3.69	132.09	125.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	NAP	O4D-C1D-N1N	3.50	111.53	107.95
2	D	401	NAP	O4D-C1D-N1N	3.47	111.50	107.95
2	B	401	NAP	O4B-C1B-N9A	-3.30	105.37	108.44
2	B	401	NAP	O7N-C7N-N7N	3.07	127.02	122.59
2	D	401	NAP	C8A-N9A-C4A	3.04	109.22	106.90
2	C	401	NAP	C8A-N9A-C4A	2.98	109.17	106.90
2	A	401	NAP	O2X-P2B-O1X	2.95	120.10	110.44
2	D	401	NAP	C4A-C5A-N7A	-2.86	107.07	109.52
2	B	401	NAP	C5A-C4A-N3A	-2.80	119.61	125.70
2	D	401	NAP	C5A-C4A-N3A	-2.75	119.71	125.70
2	A	401	NAP	O3X-P2B-O2X	2.70	118.12	107.61
2	A	401	NAP	C5A-C4A-N3A	-2.69	119.85	125.70
2	A	401	NAP	C3N-C7N-N7N	-2.62	114.79	117.77
2	A	401	NAP	C2D-C1D-N1N	-2.61	109.44	113.86
2	C	401	NAP	O3X-P2B-O1X	2.60	118.94	110.44
2	B	401	NAP	C3N-C7N-N7N	-2.56	114.86	117.77
2	B	401	NAP	C2B-C3B-C4B	2.54	107.96	101.94
2	B	401	NAP	O2N-PN-O3	2.43	115.34	108.79
2	B	401	NAP	C4A-C5A-N7A	-2.36	107.50	109.52
2	A	401	NAP	C2A-N3A-C4A	2.32	120.61	114.01
2	D	401	NAP	C2A-N3A-C4A	2.31	120.58	114.01
2	D	401	NAP	O3X-P2B-O1X	2.29	117.92	110.44
2	D	401	NAP	N7A-C8A-N9A	-2.26	107.97	114.36
2	A	401	NAP	O7N-C7N-N7N	2.24	125.83	122.59
2	C	401	NAP	C3N-C7N-N7N	-2.21	115.25	117.77
2	C	401	NAP	C5A-C4A-N3A	-2.16	121.00	125.70
2	C	401	NAP	N7A-C8A-N9A	-2.12	108.36	114.36
2	C	401	NAP	O2N-PN-O3	2.11	114.48	108.79
2	C	401	NAP	C2A-N3A-C4A	2.11	120.01	114.01
2	D	401	NAP	O2N-PN-O3	2.10	114.46	108.79
2	B	401	NAP	C2A-N3A-C4A	2.10	119.99	114.01
2	B	401	NAP	O2X-P2B-O2B	-2.10	101.04	107.09
2	C	401	NAP	C2A-N1A-C6A	2.08	122.52	118.77
2	A	401	NAP	C2B-C3B-C4B	2.07	106.85	101.94
2	B	401	NAP	O2D-C2D-C1D	2.06	117.47	111.23
2	A	401	NAP	C8A-N9A-C4A	2.02	108.44	106.90
2	A	401	NAP	O2X-P2B-O2B	-2.01	101.31	107.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	354/359 (98%)	-0.39	2 (0%) 86 88	16, 28, 59, 84	0
1	B	354/359 (98%)	-0.35	3 (0%) 83 85	15, 30, 58, 83	0
1	C	353/359 (98%)	-0.30	7 (1%) 62 63	16, 29, 58, 98	0
1	D	354/359 (98%)	-0.19	10 (2%) 50 51	17, 35, 67, 89	0
All	All	1415/1436 (98%)	-0.31	22 (1%) 68 70	15, 30, 62, 98	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	78	ASP	5.8
1	D	6	ARG	4.7
1	A	6	ARG	4.5
1	C	76	LEU	4.4
1	C	75	LYS	3.8
1	C	79	ASP	3.7
1	D	358	GLU	3.4
1	C	359	LYS	3.3
1	D	139	LYS	2.9
1	B	144	LYS	2.9
1	D	75	LYS	2.8
1	D	359	LYS	2.7
1	D	140	ARG	2.6
1	B	78	ASP	2.5
1	D	144	LYS	2.5
1	B	75	LYS	2.5
1	C	106	VAL	2.3
1	C	7	THR	2.2
1	D	142	GLU	2.1
1	D	147	ILE	2.1
1	A	139	LYS	2.0
1	D	79	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	UNL	D	402	6/-	0.15	7.42	33,40,48,50	0
3	UNL	C	402	6/-	0.13	4.95	33,37,43,59	0
3	UNL	B	402	6/-	0.14	3.48	36,40,41,43	0
3	UNL	A	402	6/-	0.11	3.15	31,35,41,44	0
2	NAP	B	401	48/48	0.26	3.01	30,62,84,103	0
2	NAP	D	401	48/48	0.18	1.93	35,59,73,90	0
2	NAP	A	401	48/48	0.16	1.87	31,52,63,69	0
2	NAP	C	401	48/48	0.14	1.25	29,45,56,67	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.