



# wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 05:21 PM GMT

PDB ID : 1DS5  
Title : DIMERIC CRYSTAL STRUCTURE OF THE ALPHA SUBUNIT IN COMPLEX WITH TWO BETA PEPTIDES MIMICKING THE ARCHITECTURE OF THE TETRAMERIC PROTEIN KINASE CK2 HOLOENZYME.  
Authors : Battistutta, R.; Sarno, S.; De Moliner, E.; Marin, O.; Zanotti, G.; Pinna, L.A.  
Deposited on : 2000-01-07  
Resolution : 3.16 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 21963  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 3.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1681 (3.22-3.10)
Ramachandran outliers	78287	1639 (3.22-3.10)
Sidechain outliers	78261	1638 (3.22-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	332	
1	B	332	
1	C	332	
1	D	332	
2	E	23	
2	F	23	
2	G	23	
2	H	23	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11932 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CASEIN KINASE, ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	328	Total	C	N	O	S	0	0	0
			2735	1762	471	490	12			
1	B	328	Total	C	N	O	S	0	0	0
			2735	1762	471	490	12			
1	C	328	Total	C	N	O	S	0	0	0
			2735	1762	471	490	12			
1	D	328	Total	C	N	O	S	0	0	0
			2735	1762	471	490	12			

- Molecule 2 is a protein called CASEIN KINASE, BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	16	Total	C	N	O	S	0	0	0
			135	90	22	22	1			
2	F	18	Total	C	N	O	S	0	0	0
			154	102	27	24	1			
2	G	16	Total	C	N	O	S	0	0	0
			135	90	22	22	1			
2	H	16	Total	C	N	O	S	0	0	0
			135	90	22	22	1			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mg	0	0
			2	2		
3	D	2	Total	Mg	0	0
			2	2		

- Molecule 4 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	C	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	D	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
4	B	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	77	Total	O	0	0
			77	77		
5	B	74	Total	O	0	0
			74	74		
5	C	89	Total	O	0	0
			89	89		
5	D	73	Total	O	0	0
			73	73		
5	E	6	Total	O	0	0
			6	6		
5	F	4	Total	O	0	0
			4	4		
5	G	6	Total	O	0	0
			6	6		
5	H	8	Total	O	0	0
			8	8		

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: CASEIN KINASE, ALPHA CHAIN



#### • Molecule 1: CASEIN KINASE, ALPHA CHAIN





● Molecule 2: CASEIN KINASE, BETA CHAIN



● Molecule 2: CASEIN KINASE, BETA CHAIN



● Molecule 2: CASEIN KINASE, BETA CHAIN



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.68Å 119.85Å 145.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.00 – 3.16	Depositor
% Data completeness (in resolution range)	95.3 (45.00-3.16)	Depositor
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, $R_{free}$	0.214 , 0.289	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11932	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: AMP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.40	18/2803 (0.6%)	1.23	13/3788 (0.3%)
1	B	1.30	7/2803 (0.2%)	1.21	11/3788 (0.3%)
1	C	1.34	10/2803 (0.4%)	1.24	18/3788 (0.5%)
1	D	1.29	10/2803 (0.4%)	1.20	11/3788 (0.3%)
2	E	1.78	2/139 (1.4%)	1.57	3/185 (1.6%)
2	F	1.74	4/158 (2.5%)	1.85	5/210 (2.4%)
2	G	1.55	1/139 (0.7%)	1.63	2/185 (1.1%)
2	H	1.84	3/139 (2.2%)	1.65	2/185 (1.1%)
All	All	1.35	55/11787 (0.5%)	1.25	65/15917 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	2
1	C	0	4
1	D	0	3
2	F	0	1
2	G	0	1
2	H	0	1
All	All	0	16

The worst 5 of 55 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	41	VAL	CA-CB	-8.84	1.36	1.54
1	D	35	GLU	CB-CG	8.82	1.69	1.52
1	D	35	GLU	CG-CD	8.30	1.64	1.51

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	167	GLU	CG-CD	8.13	1.64	1.51
1	D	147	CYS	CB-SG	-8.10	1.68	1.82

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	198	GLN	N-CA-C	8.81	134.78	111.00
1	B	124	LEU	CA-CB-CG	-8.31	96.19	115.30
1	A	134	ARG	NE-CZ-NH1	-7.92	116.34	120.30
2	F	198	GLN	N-CA-C	7.59	131.50	111.00
1	C	108	THR	N-CA-C	-6.92	92.31	111.00

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	12	TYR	Sidechain
1	A	125	TYR	Sidechain
1	A	182	TYR	Sidechain
1	A	307	TYR	Sidechain
1	B	182	TYR	Sidechain

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2735	0	2721	280	0
1	B	2735	0	2721	342	0
1	C	2735	0	2721	301	0
1	D	2735	0	2721	309	0
2	E	135	0	132	49	0
2	F	154	0	156	58	0
2	G	135	0	132	57	0
2	H	135	0	132	48	0
3	B	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	2	0	0	0	0
4	A	23	0	11	6	0
4	B	23	0	11	5	0
4	C	23	0	11	3	0
4	D	23	0	11	3	0
5	A	77	0	0	1	0
5	B	74	0	0	2	0
5	C	89	0	0	2	0
5	D	73	0	0	0	0
5	E	6	0	0	0	0
5	F	4	0	0	1	0
5	G	6	0	0	3	0
5	H	8	0	0	4	0
All	All	11932	0	11480	1344	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 58.

The worst 5 of 1344 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:H:194:PRO:CB	2:H:194:PRO:CG	1.76	1.46
1:D:119:THR:HB	2:G:202:GLN:NE2	1.42	1.30
1:A:41:VAL:HG12	2:F:197:TYR:HB3	1.24	1.19
2:F:200:GLN:HB2	2:F:202:GLN:NE2	1.61	1.13
1:A:100:ILE:HD12	1:A:101:VAL:N	1.64	1.12

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/332 (98%)	271 (83%)	41 (13%)	14 (4%)	4	30
1	B	326/332 (98%)	270 (83%)	40 (12%)	16 (5%)	3	26

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	326/332 (98%)	268 (82%)	44 (14%)	14 (4%)	4	30
1	D	326/332 (98%)	268 (82%)	44 (14%)	14 (4%)	4	30
2	E	14/23 (61%)	8 (57%)	2 (14%)	4 (29%)	0	0
2	F	16/23 (70%)	12 (75%)	1 (6%)	3 (19%)	0	0
2	G	14/23 (61%)	10 (71%)	2 (14%)	2 (14%)	0	2
2	H	14/23 (61%)	10 (71%)	2 (14%)	2 (14%)	0	2
All	All	1362/1420 (96%)	1117 (82%)	176 (13%)	69 (5%)	3	25

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	ARG
1	A	195	ARG
1	B	37	ASP
1	B	155	ARG
1	B	195	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/300 (99%)	249 (84%)	48 (16%)	3	15
1	B	297/300 (99%)	255 (86%)	42 (14%)	5	22
1	C	297/300 (99%)	257 (86%)	40 (14%)	6	25
1	D	297/300 (99%)	254 (86%)	43 (14%)	5	21
2	E	13/20 (65%)	3 (23%)	10 (77%)	0	0
2	F	15/20 (75%)	8 (53%)	7 (47%)	0	0
2	G	13/20 (65%)	8 (62%)	5 (38%)	0	0
2	H	13/20 (65%)	5 (38%)	8 (62%)	0	0
All	All	1242/1280 (97%)	1039 (84%)	203 (16%)	3	15

5 of 203 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	30	THR
1	C	195	ARG
2	F	199	LEU
1	C	43	ARG
1	C	118	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 58 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	150	GLN
1	C	291	HIS
2	F	193	HIS
1	C	160	HIS
1	C	189	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	AMP	A	501	-	25,25,25	1.83	7 (28%)	38,38,38	3.11	9 (23%)
4	AMP	B	801	-	25,25,25	1.55	7 (28%)	38,38,38	3.37	13 (34%)
4	AMP	C	601	-	25,25,25	1.59	5 (20%)	38,38,38	2.53	11 (28%)
4	AMP	D	701	-	25,25,25	1.76	8 (32%)	38,38,38	2.72	10 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	AMP	A	501	-	1/1/5/5	0/10/26/26	0/1/3/3
4	AMP	B	801	-	1/1/5/5	0/10/26/26	0/1/3/3
4	AMP	C	601	-	1/1/5/5	0/10/26/26	0/1/3/3
4	AMP	D	701	-	1/1/5/5	0/10/26/26	0/1/3/3

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	AMP	C4-N3	5.16	1.43	1.35
4	D	701	AMP	O5'-C5'	3.60	1.59	1.44
4	A	501	AMP	C2'-C1'	3.42	1.58	1.53
4	D	701	AMP	C3'-C4'	3.19	1.61	1.53
4	C	601	AMP	P-O3P	3.12	1.66	1.54

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	801	AMP	O4'-C1'-N9	16.60	123.88	108.44
4	A	501	AMP	O4'-C1'-N9	15.05	122.44	108.44
4	D	701	AMP	O4'-C1'-N9	12.75	120.30	108.44
4	C	601	AMP	O4'-C1'-N9	10.33	118.05	108.44
4	B	801	AMP	C4'-O4'-C1'	-6.28	102.93	109.75

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	501	AMP	C1'
4	C	601	AMP	C1'
4	B	801	AMP	C1'
4	D	701	AMP	C1'

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.