



Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 05:03 AM GMT

PDB ID : 4DS4
Title : Ternary complex of Bacillus DNA Polymerase I Large Fragment, DNA duplex,
and rCTP in presence of Mn²⁺
Authors : Wang, W.; Beese, L.S.
Deposited on : 2012-02-17
Resolution : 1.68 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3587 (1.70-1.66)
Clashscore	79885	4225 (1.70-1.66)
Ramachandran outliers	78287	4144 (1.70-1.66)
Sidechain outliers	78261	4143 (1.70-1.66)
RSRZ outliers	66119	3587 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	592	
1	D	592	
2	B	9	
2	E	9	
3	C	13	
3	F	13	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 20911 atoms, of which 9809 are hydrogens and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	580	Total	C	H	N	O	S	0	2	0
			9391	2963	4731	810	870	17			
1	D	561	Total	C	H	N	O	S	0	20	0
			9228	2914	4655	791	853	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ALA	ASP	ENGINEERED MUTATION	UNP Q5KWC1
A	823	HIS	ARG	ENGINEERED MUTATION	UNP Q5KWC1
D	598	ALA	ASP	ENGINEERED MUTATION	UNP Q5KWC1
D	823	HIS	ARG	ENGINEERED MUTATION	UNP Q5KWC1

- Molecule 2 is a DNA chain called DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	9	Total	C	H	N	O	P	0	0	0
			274	85	100	29	52	8			
2	E	9	Total	C	H	N	O	P	0	0	0
			274	85	100	29	52	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*CP*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	11	Total	C	H	N	O	P	0	0	0
			356	109	123	47	66	11			
3	F	9	Total	C	H	N	O	P	0	0	0
			291	89	100	40	53	9			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	354	Total	O	0	0
			354	354		
5	D	570	Total	O	0	0
			570	570		
5	B	28	Total	O	0	0
			28	28		
5	C	47	Total	O	0	0
			47	47		
5	E	29	Total	O	0	0
			29	29		
5	F	54	Total	O	0	0
			54	54		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase



- Molecule 1: DNA polymerase



- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3')



- Molecule 2: DNA (5'-D(*CP*CP*TP*GP*AP*CP*TP*CP*(DOC))-3')



- Molecule 3: DNA (5'-D(*CP*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain C: 



- Molecule 3: DNA (5'-D(*CP*AP*TP*GP*GP*GP*AP*GP*TP*CP*AP*GP*G)-3')

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.79Å 108.77Å 150.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.25 – 1.68 64.25 – 1.68	Depositor EDS
% Data completeness (in resolution range)	87.7 (64.25-1.68) 87.7 (64.25-1.68)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.68Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1026)	Depositor
R, R_{free}	0.174 , 0.199 0.174 , 0.199	Depositor DCC
R_{free} test set	6893 reflections (4.49%)	DCC
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.140	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1730 of 153533 reflections (1.127%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	20911	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DOC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/4750	0.57	0/6418
1	D	0.63	0/4737	0.70	4/6401 (0.1%)
2	B	0.91	0/173	1.94	8/264 (3.0%)
2	E	1.09	0/173	1.82	2/264 (0.8%)
3	C	1.04	0/262	1.59	6/404 (1.5%)
3	F	1.30	0/215	1.95	5/331 (1.5%)
All	All	0.60	0/10310	0.81	25/14082 (0.2%)

There are no bond length outliers.

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	23	DT	O4'-C1'-N1	-9.98	101.02	108.00
2	B	25	DA	O4'-C4'-C3'	-9.32	100.41	106.00
3	F	6	DA	OP1-P-OP2	-7.18	108.83	119.60
2	B	25	DA	O5'-P-OP1	6.36	118.33	110.70
3	F	12	DG	O4'-C1'-N9	6.30	112.41	108.00
1	D	660	ARG	NE-CZ-NH1	6.25	123.42	120.30
3	F	10	DA	O4'-C1'-N9	-6.24	103.63	108.00
3	F	8	DT	O4'-C1'-N1	-6.18	103.67	108.00
3	F	5	DG	C2-N3-C4	-5.97	108.92	111.90
3	C	11	DG	O4'-C1'-N9	-5.94	103.84	108.00
3	C	5	DG	C5-C6-N1	-5.76	108.62	111.50
1	D	859	ARG	NE-CZ-NH1	5.75	123.18	120.30
3	C	8	DT	O4'-C1'-N1	-5.74	103.98	108.00
2	B	23	DT	O4'-C1'-N1	-5.67	104.03	108.00
1	D	409	ASP	CB-CG-OD2	5.66	123.40	118.30
2	B	26	DC	O4'-C1'-N1	5.60	111.92	108.00
2	B	25	DA	O5'-P-OP2	-5.43	100.81	105.70
3	C	5	DG	N1-C6-O6	5.38	123.12	119.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	25	DA	C5'-C4'-C3'	5.37	123.77	114.10
2	B	26	DC	O4'-C1'-C2'	5.37	110.19	105.90
1	D	859	ARG	NE-CZ-NH2	-5.16	117.72	120.30
2	E	26	DC	N1-C2-O2	-5.15	115.81	118.90
3	C	5	DG	N3-C4-N9	-5.02	122.99	126.00
3	C	6	DA	C5-N7-C8	-5.01	101.39	103.90
2	B	26	DC	C6-N1-C2	5.00	122.30	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4660	4731	0	24	0
1	D	4573	4655	0	46	0
2	B	174	100	0	5	0
2	E	174	100	0	1	0
3	C	233	123	0	0	0
3	F	191	100	0	1	0
4	D	15	0	0	0	0
5	A	354	0	0	7	2
5	B	28	0	0	0	0
5	C	47	0	0	0	0
5	D	570	0	0	20	2
5	E	29	0	0	0	0
5	F	54	0	0	1	0
All	All	11102	9809	0	70	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 4.

All (70) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:845:CYS:SG	5:D:1375:HOH:O	2.10	1.07
1:A:388:CYS:SG	5:A:1043:HOH:O	2.23	0.95
1:D:416:MET:HE3	5:D:1543:HOH:O	1.66	0.94
1:A:408:ASP:OD1	5:A:1012:HOH:O	1.89	0.91
1:D:845:CYS:SG	5:D:1481:HOH:O	2.29	0.90
1:A:582:LYS:HD2	2:B:26:DC:H1'	1.60	0.83
1:D:846:ARG:HD2	5:D:1535:HOH:O	1.80	0.80
1:D:758:LYS:CE	5:D:1562:HOH:O	2.33	0.77
1:D:788:GLU:OE2	5:D:1562:HOH:O	2.01	0.77
1:D:305:ASP:OD1	5:D:1435:HOH:O	2.03	0.75
1:A:423:ARG:NE	5:A:1239:HOH:O	2.20	0.70
1:D:459:ARG:NH2	5:D:1342:HOH:O	2.25	0.69
1:A:464:GLU:OE1	5:A:1159:HOH:O	2.11	0.69
1:D:664:HIS:HD1	1:D:859:ARG:H	1.40	0.68
1:D:751:GLU:OE2	5:D:1267:HOH:O	2.11	0.67
3:F:11:DG:OP1	5:F:144:HOH:O	2.13	0.67
1:D:733:ALA:O	5:D:1522:HOH:O	2.12	0.65
1:A:812:ASN:O	5:A:1073:HOH:O	2.15	0.64
1:D:658:GLU:OE1	5:D:1559:HOH:O	2.15	0.63
1:D:758:LYS:NZ	5:D:1562:HOH:O	2.23	0.63
1:D:758:LYS:HE2	5:D:1562:HOH:O	1.98	0.63
1:D:520:GLU:OE1	5:D:1490:HOH:O	2.16	0.62
1:D:561:LEU:HB3	1:D:571:VAL:HG13	1.83	0.60
1:D:561:LEU:O	1:D:571:VAL:HG11	2.01	0.60
1:A:467:ARG:NH2	5:A:1107:HOH:O	2.35	0.59
1:D:708:VAL:HG13	1:D:712:ILE:CD1	2.34	0.57
1:D:515:GLU:HG2	1:D:519:TYR:CE2	2.43	0.54
1:D:708:VAL:HG13	1:D:712:ILE:HD12	1.90	0.52
1:D:520:GLU:CD	5:D:1490:HOH:O	2.48	0.50
1:A:469:GLU:HG3	1:D:466:ARG:NE	2.26	0.50
1:A:534:LEU:HD11	1:A:574:ILE:HD13	1.95	0.48
1:A:459:ARG:HB2	1:A:460:PRO:HD3	1.94	0.48
1:D:721:LEU:HB3	1:D:732:ALA:HB1	1.96	0.48
1:A:848:VAL:HB	1:A:849:PRO:HD3	1.96	0.47
1:D:415:LYS:NZ	5:D:1318:HOH:O	2.48	0.47
1:A:325:GLU:OE1	5:A:1248:HOH:O	2.20	0.47
1:A:440:GLU:N	1:A:441:PRO:HD2	2.29	0.47
1:D:734:GLU:HB3	1:D:738:ARG:NH1	2.28	0.47
1:A:469:GLU:CD	1:D:466:ARG:HE	2.17	0.47
1:D:459:ARG:CZ	5:D:1342:HOH:O	2.62	0.47
1:D:510:GLN:O	1:D:514:VAL:HG23	2.15	0.46
1:D:814:ARG:HG3	1:D:851[B]:VAL:HG11	1.98	0.46
1:D:645:SER:O	1:D:646:ASP:HB2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:22:DC:H2'	2:B:23:DT:H71	1.97	0.45
1:D:525:GLU:O	5:D:1365:HOH:O	2.21	0.45
1:D:550:THR:CG2	5:D:1376:HOH:O	2.65	0.45
1:A:469:GLU:CD	1:D:466:ARG:NE	2.70	0.44
1:D:814:ARG:NH2	1:D:847[A]:LEU:HD13	2.33	0.44
1:D:304:ALA:CB	1:D:311:MET:HE1	2.48	0.44
1:D:756:GLU:OE2	1:D:760:LYS:HE2	2.18	0.44
1:D:500:LEU:HD21	1:D:591:LEU:HD23	1.99	0.44
1:D:754:VAL:CG1	1:D:758:LYS:HE3	2.48	0.44
1:D:629:ARG:HH22	2:E:29:DOC:H5	1.83	0.44
1:A:687:MET:HE3	1:A:694:GLU:N	2.33	0.43
1:D:722:ALA:HB2	1:D:729:ARG:HA	2.01	0.43
1:D:708:VAL:HG13	1:D:712:ILE:HG13	2.01	0.43
1:A:551:LYS:NZ	2:B:23:DT:OP1	2.52	0.43
1:A:446:HIS:NE2	1:A:450:LYS:HD2	2.34	0.42
1:D:712:ILE:HA	1:D:716:ILE:HG22	2.02	0.42
1:D:814:ARG:CZ	1:D:847[A]:LEU:HD13	2.48	0.42
2:B:21:DC:H2'	2:B:22:DC:C6	2.54	0.42
1:D:764:THR:HA	1:D:769:ARG:O	2.20	0.42
1:A:578:ARG:NH1	2:B:25:DA:H5''	2.35	0.42
1:A:522:ALA:O	1:A:541:LYS:HE2	2.20	0.41
1:A:349:GLU:CD	1:A:349:GLU:H	2.23	0.41
1:D:681:ILE:HD12	1:D:681:ILE:HA	1.91	0.41
1:D:435:ARG:NH2	5:D:1428:HOH:O	2.53	0.41
1:A:582:LYS:HG3	1:A:582:LYS:O	2.20	0.41
1:A:446:HIS:CE1	1:A:450:LYS:HD2	2.56	0.41
1:A:766:LEU:HD23	1:A:766:LEU:HA	1.96	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:A:977:HOH:O	5:D:1176:HOH:O[2_745]	1.72	0.48
5:A:1010:HOH:O	5:D:1088:HOH:O[2_745]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone

conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	580/592 (98%)	567 (98%)	13 (2%)	0	100	100
1	D	577/592 (98%)	566 (98%)	11 (2%)	0	100	100
All	All	1157/1184 (98%)	1133 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	497/507 (98%)	494 (99%)	3 (1%)	92	86
1	D	497/507 (98%)	494 (99%)	3 (1%)	92	86
All	All	994/1014 (98%)	988 (99%)	6 (1%)	92	86

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	477	LEU
1	A	582	LYS
1	A	838	LYS
1	D	303	LEU
1	D	513	THR
1	D	782	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	468	ASN
1	A	470	GLN
1	A	510	GLN
1	A	529	ASN
1	A	573	ASN
1	A	691	GLN
1	A	724	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	755	GLN
1	D	755	GLN
1	D	793	ASN
1	D	854	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DOC	B	29	3,2	17,19,20	0.76	0	20,26,29	1.13	1 (5%)
2	DOC	E	29	3,2	17,19,20	0.92	1 (5%)	20,26,29	1.28	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DOC	B	29	3,2	-	0/5/18/19	0/2/2/2
2	DOC	E	29	3,2	-	0/5/18/19	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	29	DOC	C6-C5	-2.28	1.32	1.38

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	29	DOC	C2-N3-C4	3.16	120.15	115.57
2	E	29	DOC	O4'-C4'-C5'	-2.85	105.69	109.67
2	E	29	DOC	C2-N3-C4	2.15	118.69	115.57
2	E	29	DOC	C6-C5-C4	2.11	118.35	117.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	D	901	-	4,4,4	0.25	0	6,6,6	0.35	0
4	SO4	D	902	-	4,4,4	0.18	0	6,6,6	0.21	0
4	SO4	D	903	-	4,4,4	0.22	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	D	901	-	-	0/0/0/0	0/0/0/0
4	SO4	D	902	-	-	0/0/0/0	0/0/0/0
4	SO4	D	903	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	580/592 (97%)	0.49	56 (9%) 8 9	21, 41, 66, 91	0
1	D	561/592 (94%)	0.31	30 (5%) 25 28	11, 25, 47, 63	0
2	B	9/9 (100%)	-0.47	0 100 100	26, 32, 47, 60	0
2	E	9/9 (100%)	-0.20	0 100 100	20, 27, 44, 57	0
3	C	11/13 (84%)	-0.30	0 100 100	21, 28, 47, 66	0
3	F	9/13 (69%)	-0.27	0 100 100	16, 21, 34, 41	0
All	All	1179/1228 (96%)	0.38	86 (7%) 15 17	11, 33, 59, 91	0

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	712	ILE	10.1
1	D	685	THR	5.7
1	D	727	ILE	5.6
1	D	735	PHE	5.6
1	A	297	ALA	5.5
1	A	298	LYS	5.3
1	A	518	ILE	5.3
1	D	713	VAL	5.3
1	A	689	ILE	5.3
1	A	679	LEU	5.0
1	A	736	ILE	4.8
1	A	819	ARG	4.8
1	D	731	GLU	4.7
1	A	571	VAL	4.6
1	A	537	ILE	4.5
1	A	514	VAL	4.5
1	D	307	VAL	4.1
1	A	697	VAL	4.1
1	A	699	PRO	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	736	ILE	4.0
1	D	708	VAL	4.0
1	A	708	VAL	3.8
1	D	710	PHE	3.7
1	D	296	LEU	3.7
1	D	509	GLU	3.7
1	A	681	ILE	3.6
1	A	816	LYS	3.6
1	D	724	ASN	3.5
1	A	570	ILE	3.5
1	A	856	VAL	3.4
1	A	521	LEU	3.4
1	D	725	LEU	3.3
1	D	734	GLU	3.2
1	A	630	LEU	3.2
1	A	703	ARG	3.1
1	A	846	ARG	3.1
1	D	707	ALA	3.1
1	D	683	THR	3.1
1	A	727	ILE	3.1
1	A	528	ILE	3.0
1	A	833	ILE	3.0
1	D	681	ILE	2.8
1	A	588	ILE	2.8
1	D	716	ILE	2.8
1	A	685	THR	2.8
1	D	732	ALA	2.8
1	A	700	ASN	2.8
1	A	574	ILE	2.8
1	A	695	ASP	2.8
1	A	692	VAL	2.8
1	A	442	VAL	2.7
1	A	842	GLU	2.7
1	A	337	VAL	2.7
1	A	649	ILE	2.7
1	A	602	VAL	2.7
1	A	864	VAL	2.7
1	A	303	LEU	2.6
1	A	860	VAL	2.5
1	A	710	PHE	2.5
1	A	495	VAL	2.4
1	A	746	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	684	LYS	2.4
1	A	307	VAL	2.4
1	A	353	ALA	2.4
1	A	636	ILE	2.4
1	D	726	ASN	2.4
1	A	783	VAL	2.4
1	D	297	ALA	2.3
1	D	514	VAL	2.3
1	A	698	THR	2.3
1	D	595	VAL	2.3
1	A	569	GLU	2.2
1	A	308	THR	2.2
1	D	570	ILE	2.2
1	A	714	TYR	2.1
1	A	848	VAL	2.1
1	A	676	ARG	2.1
1	D	493[A]	VAL	2.1
1	A	691	GLN	2.1
1	A	513	THR	2.1
1	D	730	LYS	2.1
1	A	712	ILE	2.0
1	D	537	ILE	2.0
1	A	866	TYR	2.0
1	A	507	LEU	2.0
1	D	721	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	DOC	E	29	18/19	0.10	-0.25	17,21,29,35	0
2	DOC	B	29	18/19	0.07	-0.72	28,33,43,43	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	D	901	5/5	0.08	0.29	33,36,48,50	0
4	SO4	D	902	5/5	0.09	-0.39	36,44,49,59	0
4	SO4	D	903	5/5	0.12	-0.61	75,82,85,85	0

6.5 Other polymers

There are no such residues in this entry.