



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 04:36 AM GMT

PDB ID : 2DVY
Title : Crystal structure of restriction endonucleases PabI
Authors : Miyazono, K.; Watanabe, M.; Kamo, M.; Sawasaki, T.; Nagata, K.; Endo, Y.;
Tanokura, M.; Kobayashi, I.
Deposited on : 2006-08-01
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)
RSRZ outliers	66119	1217 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	226	
1	B	226	
1	C	226	
1	D	226	
1	E	226	
1	F	226	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 10623 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Restriction endonuclease PabI.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1817	1167	310	335	5			
1	B	215	Total	C	N	O	S	0	0	0
			1747	1123	295	325	4			
1	C	218	Total	C	N	O	S	0	0	0
			1776	1142	300	329	5			
1	D	214	Total	C	N	O	S	0	0	0
			1747	1120	299	324	4			
1	E	223	Total	C	N	O	S	0	0	0
			1817	1167	310	335	5			
1	F	211	Total	C	N	O	S	0	0	0
			1719	1106	289	320	4			

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Restriction endonuclease PabI

Chain A:




• Molecule 1: Restriction endonuclease PabI

Chain B:



• Molecule 1: Restriction endonuclease PabI

Chain C:



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	84.57Å 113.96Å 89.17Å 90.00° 116.23° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00 19.94 – 2.99	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-3.00) 94.6 (19.94-2.99)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.21 (at 2.98Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.247 , 0.318 0.246 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	60.9	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 40.6	EDS
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 30339 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10623	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/1853	0.73	1/2501 (0.0%)
1	B	0.48	0/1781	0.68	0/2404
1	C	0.56	0/1811	0.81	1/2444 (0.0%)
1	D	0.48	1/1782 (0.1%)	0.67	0/2404
1	E	0.45	0/1853	0.67	0/2501
1	F	0.43	0/1752	0.67	0/2364
All	All	0.49	1/10832 (0.0%)	0.71	2/14618 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	58	ARG	CZ-NH2	5.62	1.40	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	160	TYR	N-CA-C	5.84	126.76	111.00
1	A	120	THR	N-CA-C	-5.52	96.10	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1817	0	1859	144	0
1	B	1747	0	1776	128	0
1	C	1776	0	1810	180	0
1	D	1747	0	1775	175	0
1	E	1817	0	1859	186	0
1	F	1719	0	1750	234	0
All	All	10623	0	10829	960	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 45.

The worst 5 of 960 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:124:LEU:HB2	1:D:129:ILE:HD11	1.16	1.16
1:F:13:PHE:HB2	1:F:84:LEU:HD21	1.33	1.09
1:A:61:TYR:CD2	1:A:167:CYS:HB3	1.94	1.01
1:E:132:ASP:HB2	1:F:134:TYR:HB2	1.38	1.00
1:F:50:SER:HA	1:F:181:VAL:HB	1.42	0.99

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/226 (98%)	178 (80%)	33 (15%)	10 (4%)	4	22
1	B	211/226 (93%)	165 (78%)	37 (18%)	9 (4%)	4	23
1	C	214/226 (95%)	173 (81%)	30 (14%)	11 (5%)	3	18
1	D	212/226 (94%)	150 (71%)	46 (22%)	16 (8%)	2	8
1	E	221/226 (98%)	171 (77%)	35 (16%)	15 (7%)	2	10
1	F	207/226 (92%)	141 (68%)	50 (24%)	16 (8%)	1	7
All	All	1286/1356 (95%)	978 (76%)	231 (18%)	77 (6%)	2	14

5 of 77 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	55	GLU
1	A	56	ASN
1	A	184	ARG
1	B	57	LEU
1	C	49	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/204 (99%)	182 (90%)	20 (10%)	11	40
1	B	194/204 (95%)	173 (89%)	21 (11%)	9	35
1	C	198/204 (97%)	171 (86%)	27 (14%)	5	24
1	D	193/204 (95%)	173 (90%)	20 (10%)	10	37
1	E	202/204 (99%)	180 (89%)	22 (11%)	9	35
1	F	191/204 (94%)	171 (90%)	20 (10%)	10	37
All	All	1180/1224 (96%)	1050 (89%)	130 (11%)	9	34

5 of 130 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	140	GLU
1	D	24	ILE
1	F	116	ARG
1	C	145	GLU
1	C	196	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	185	ASN
1	D	121	ASN
1	F	161	GLN
1	D	15	ASN
1	D	128	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	A	223/226 (98%)	-0.24	0	100	100	31, 52, 76, 91	0
1	B	215/226 (95%)	-0.18	0	100	100	29, 59, 87, 94	0
1	C	218/226 (96%)	-0.28	0	100	100	22, 48, 78, 95	0
1	D	214/226 (94%)	-0.07	0	100	100	25, 80, 113, 119	0
1	E	223/226 (98%)	-0.11	0	100	100	39, 70, 101, 111	0
1	F	211/226 (93%)	0.03	1 (0%)	88	36	45, 80, 118, 134	0
All	All	1304/1356 (96%)	-0.14	1 (0%)	93	63	22, 65, 104, 134	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	55	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.