



wwPDB X-ray Structure Validation Summary Report

Mar 1, 2014 – 01:19 AM GMT

PDB ID : 2E2K
Title : Helicobacter pylori formamidase AmiF contains a fine-tuned cysteine-glutamate-lysine catalytic triad
Authors : Wang, W.C.; Hung, C.L.
Deposited on : 2006-11-14
Resolution : 2.50 Å (reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	334	
1	B	334	
1	C	334	
1	D	334	
1	E	334	
1	F	334	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15744 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formamidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2482	1597	416	458	11			
1	B	314	Total	C	N	O	S	0	0	0
			2485	1598	417	459	11			
1	C	314	Total	C	N	O	S	0	0	0
			2489	1600	417	461	11			
1	D	311	Total	C	N	O	S	0	0	0
			2458	1581	412	454	11			
1	E	315	Total	C	N	O	S	0	0	0
			2493	1602	419	461	11			
1	F	314	Total	C	N	O	S	0	0	0
			2487	1601	415	460	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	166	SER	CYS	ENGINEERED	UNP O25836
B	166	SER	CYS	ENGINEERED	UNP O25836
C	166	SER	CYS	ENGINEERED	UNP O25836
D	166	SER	CYS	ENGINEERED	UNP O25836
E	166	SER	CYS	ENGINEERED	UNP O25836
F	166	SER	CYS	ENGINEERED	UNP O25836

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	148	Total	O	0	0
			148	148		
2	B	151	Total	O	0	0
			151	151		
2	C	133	Total	O	0	0
			133	133		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	D	159	Total 159	O 159	0	0
2	E	138	Total 138	O 138	0	0
2	F	121	Total 121	O 121	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Formamidase



• Molecule 1: Formamidase




• Molecule 1: Formamidase




- Molecule 1: Formamidase

Chain D:




- Molecule 1: Formamidase

Chain E:



- Molecule 1: Formamidase

Chain F:



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.72Å 130.53Å 144.59Å 90.00° 99.44° 90.00°	Depositor
Resolution (Å)	30.00 – 2.50 26.90 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.50) 99.8 (26.90-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	13.24 (at 2.50Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.256 , 0.294 0.256 , 0.264	Depositor DCC
R_{free} test set	3760 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	13.2	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 7.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 74588 reflections	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	15744	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2550	0.46	0/3459
1	B	0.27	0/2553	0.47	0/3465
1	C	0.26	0/2558	0.47	0/3474
1	D	0.27	0/2523	0.46	0/3421
1	E	0.27	0/2562	0.46	0/3480
1	F	0.26	0/2556	0.47	0/3471
All	All	0.27	0/15302	0.46	0/20770

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	249	ASN	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the

Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2482	0	2407	111	0
1	B	2485	0	2417	84	0
1	C	2489	0	2409	101	0
1	D	2458	0	2382	83	0
1	E	2493	0	2423	92	0
1	F	2487	0	2411	75	0
2	A	148	0	0	6	0
2	B	151	0	0	1	0
2	C	133	0	0	6	0
2	D	159	0	0	4	0
2	E	138	0	0	3	0
2	F	121	0	0	0	0
All	All	15744	0	14449	455	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

The worst 5 of 455 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:F:192:TYR:HB3	1:F:193:SER:CA	1.69	1.23
1:E:248:ARG:HD3	1:E:248:ARG:H	1.06	1.20
1:A:192:TYR:CB	1:A:193:SER:HA	1.73	1.18
1:D:192:TYR:HB3	1:D:193:SER:CA	1.76	1.14
1:C:192:TYR:HB3	1:C:193:SER:CA	1.78	1.13

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	303/334 (91%)	282 (93%)	19 (6%)	2 (1%)	30 50
1	B	306/334 (92%)	282 (92%)	18 (6%)	6 (2%)	11 17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	306/334 (92%)	287 (94%)	12 (4%)	7 (2%)	10	14
1	D	298/334 (89%)	278 (93%)	16 (5%)	4 (1%)	18	29
1	E	309/334 (92%)	281 (91%)	22 (7%)	6 (2%)	12	19
1	F	306/334 (92%)	278 (91%)	22 (7%)	6 (2%)	11	17
All	All	1828/2004 (91%)	1688 (92%)	109 (6%)	31 (2%)	14	22

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	194	THR
1	C	194	THR
1	E	194	THR
1	E	226	VAL
1	F	250	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/277 (94%)	247 (95%)	14 (5%)	31	53
1	B	262/277 (95%)	252 (96%)	10 (4%)	44	71
1	C	262/277 (95%)	254 (97%)	8 (3%)	52	79
1	D	259/277 (94%)	252 (97%)	7 (3%)	57	83
1	E	263/277 (95%)	254 (97%)	9 (3%)	49	75
1	F	262/277 (95%)	251 (96%)	11 (4%)	40	66
All	All	1569/1662 (94%)	1510 (96%)	59 (4%)	44	71

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	224	ASP
1	D	223	TYR
1	F	232	GLU
1	C	248	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	297	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	281	HIS
1	D	197	ASN
1	F	167	HIS
1	C	292	HIS
1	D	35	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/334 (93%)	0.44	24 (7%) 13 13	2, 8, 20, 28	0
1	B	314/334 (94%)	0.39	24 (7%) 14 13	3, 8, 18, 25	0
1	C	314/334 (94%)	0.33	24 (7%) 14 13	2, 7, 15, 24	0
1	D	311/334 (93%)	0.38	23 (7%) 14 14	2, 7, 17, 28	0
1	E	315/334 (94%)	0.57	25 (7%) 13 12	3, 10, 21, 29	0
1	F	314/334 (94%)	0.56	29 (9%) 9 8	4, 10, 21, 26	0
All	All	1881/2004 (93%)	0.44	149 (7%) 13 12	2, 8, 20, 29	0

The worst 5 of 149 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	222	GLY	11.8
1	A	222	GLY	10.9
1	E	228	TYR	10.1
1	B	222	GLY	10.1
1	F	225	ASN	10.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.