



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 04:10 AM GMT

PDB ID : 2E32
Title : Structural basis for selection of glycosylated substrate by SCFFbs1 ubiquitin ligase
Authors : Mizushima, T.; Yoshida, Y.; Kumanomidou, T.; Hasegawa, Y.; Yamane, T.; Tanaka, K.
Deposited on : 2006-11-20
Resolution : 3.52 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1256 (3.74-3.30)
Clashscore	79885	1020 (3.68-3.36)
Ramachandran outliers	78287	1082 (3.70-3.34)
Sidechain outliers	78261	1082 (3.70-3.34)
RSRZ outliers	66119	1256 (3.74-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	297	
1	C	297	
2	B	166	
2	D	166	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5970 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F-box only protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	240	Total	C	N	O	S	0	0	0
			1948	1249	324	369	6			
1	C	240	Total	C	N	O	S	0	0	0
			1948	1249	324	369	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	LYS	ARG	SEE REMARK 999	UNP Q80UW2
C	151	LYS	ARG	SEE REMARK 999	UNP Q80UW2

- Molecule 2 is a protein called S-phase kinase-associated protein 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	130	Total	C	N	O	S	0	0	0
			1037	660	167	206	4			
2	D	130	Total	C	N	O	S	0	0	0
			1037	660	167	206	4			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	CLONING ARTIFACT	UNP P63208
B	-1	PRO	-	CLONING ARTIFACT	UNP P63208
B	0	HIS	-	CLONING ARTIFACT	UNP P63208
D	-2	GLY	-	CLONING ARTIFACT	UNP P63208
D	-1	PRO	-	CLONING ARTIFACT	UNP P63208
D	0	HIS	-	CLONING ARTIFACT	UNP P63208

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: F-box only protein 2

Chain A:



- Molecule 1: F-box only protein 2


Chain C:



- Molecule 2: S-phase kinase-associated protein 1A

Chain B:





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.94Å 109.82Å 153.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.53 – 3.52 56.53 – 3.52	Depositor EDS
% Data completeness (in resolution range)	89.7 (56.53-3.52) 89.7 (56.53-3.52)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.13 (at 3.49Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.223 , 0.299 0.219 , 0.286	Depositor DCC
R_{free} test set	651 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	68.3	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 12930 reflections	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	5970	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.06	2/2003 (0.1%)	1.00	2/2725 (0.1%)
1	C	1.01	1/2003 (0.0%)	0.97	1/2725 (0.0%)
2	B	1.13	5/1052 (0.5%)	1.11	3/1422 (0.2%)
2	D	1.10	3/1052 (0.3%)	1.09	2/1422 (0.1%)
All	All	1.06	11/6110 (0.2%)	1.02	8/8294 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
2	B	0	2
2	D	0	1
All	All	0	4

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	113	LYS	CD-CE	8.71	1.73	1.51
2	D	113	LYS	CD-CE	8.16	1.71	1.51
1	A	296	GLU	CG-CD	7.51	1.63	1.51
1	A	279	TYR	CE2-CZ	6.92	1.47	1.38
2	D	150	GLU	CB-CG	6.13	1.63	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	61	ARG	NE-CZ-NH1	8.41	124.50	120.30
1	A	129	ARG	NE-CZ-NH1	5.93	123.26	120.30
2	D	113	LYS	CB-CG-CD	5.85	126.81	111.60
2	B	113	LYS	CB-CG-CD	5.77	126.61	111.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	226	ARG	NE-CZ-NH2	-5.21	117.69	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	146	THR	Peptide
2	B	83	ASP	Peptide
1	C	165	GLN	Peptide
2	D	83	ASP	Peptide

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1948	0	1847	139	1
1	C	1948	0	1847	128	0
2	B	1037	0	1041	123	1
2	D	1037	0	1041	123	0
All	All	5970	0	5776	497	1

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 42.

The worst 5 of 497 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:D:148:GLU:O	2:D:149:GLU:HG3	1.41	1.19
2:B:64:HIS:CE1	2:D:128:LYS:HE2	1.81	1.16
2:B:64:HIS:HE1	2:D:128:LYS:HE2	1.05	1.15
2:B:148:GLU:O	2:B:149:GLU:HG3	1.57	1.03
2:B:64:HIS:HE1	2:D:128:LYS:CE	1.72	1.01

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:165:GLN:NE2	2:B:33:ASP:O[4.455]	2.09	0.11

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	236/297 (80%)	188 (80%)	32 (14%)	16 (7%)	2	27
1	C	236/297 (80%)	185 (78%)	36 (15%)	15 (6%)	2	29
2	B	124/166 (75%)	90 (73%)	21 (17%)	13 (10%)	1	14
2	D	124/166 (75%)	90 (73%)	20 (16%)	14 (11%)	1	13
All	All	720/926 (78%)	553 (77%)	109 (15%)	58 (8%)	1	21

5 of 58 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	101	LEU
1	A	136	ASP
1	A	143	VAL
1	A	230	GLU
1	A	260	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/250 (82%)	167 (81%)	39 (19%)	2	13
1	C	206/250 (82%)	168 (82%)	38 (18%)	2	14
2	B	119/152 (78%)	91 (76%)	28 (24%)	1	7
2	D	119/152 (78%)	85 (71%)	34 (29%)	0	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	650/804 (81%)	511 (79%)	139 (21%)	1 9

5 of 139 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	146	THR
1	C	113	HIS
2	D	127	ILE
2	B	150	GLU
1	C	69	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	60	GLN
1	C	97	GLN
2	D	60	GLN
2	B	64	HIS
1	A	116	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	240/297 (80%)	0.12	1 (0%) 90 71	31, 50, 66, 71	0
1	C	240/297 (80%)	0.09	0 100 100	31, 49, 67, 71	0
2	B	130/166 (78%)	0.15	1 (0%) 83 53	31, 52, 70, 82	0
2	D	130/166 (78%)	0.00	0 100 100	31, 52, 70, 82	0
All	All	740/926 (79%)	0.10	2 (0%) 91 76	31, 51, 67, 82	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	LYS	2.5
2	B	7	GLN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.