



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 05:54 AM GMT

PDB ID : 1E5X
Title : STRUCTURE OF THREONINE SYNTHASE FROM ARABIDOPSIS
THALIANA
Authors : Thomazeau, K.; Curien, G.; Dumas, R.; Biou, V.
Deposited on : 2000-08-04
Resolution : 2.25 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	486	
1	B	486	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7289 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THREONINE SYNTHASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	451	Total	C	N	O	S	Se	0	2	0
			3514	2236	592	664	9	13			
1	B	433	Total	C	N	O	S	Se	0	3	0
			3387	2158	569	638	9	13			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	309	MSO	MET	MODIFIED RESIDUE	UNP Q39144
A	77	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	146	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	167	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	180	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	221	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	245	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	322	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	388	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	391	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	399	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	453	MSE	MET	MODIFIED RESIDUE	UNP Q39144
A	472	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	309	MSO	MET	MODIFIED RESIDUE	UNP Q39144
B	77	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	146	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	167	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	180	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	221	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	245	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	322	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	388	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	391	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	399	MSE	MET	MODIFIED RESIDUE	UNP Q39144
B	453	MSE	MET	MODIFIED RESIDUE	UNP Q39144

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	472	MSE	MET	MODIFIED RESIDUE	UNP Q39144


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	203	Total 203	O 203	0	0
2	B	185	Total 185	O 185	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: THREONINE SYNTHASE



• Molecule 1: THREONINE SYNTHASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.76Å 62.14Å 76.59Å 109.48° 97.61° 112.74°	Depositor
Resolution (Å)	29.81 – 2.25 29.81 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.9 (29.81-2.25) 98.0 (29.81-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.38 (at 2.24Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.222 , 0.243 0.225 , 0.221	Depositor DCC
R_{free} test set	2071 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	33.3	Xtriage
Anisotropy	0.537	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 81903 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7289	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3571	0.62	1/4822 (0.0%)
1	B	0.37	0/3442	0.62	2/4646 (0.0%)
All	All	0.37	0/7013	0.62	3/9468 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	387	LEU	CA-CB-CG	7.06	131.54	115.30
1	B	387	LEU	CA-CB-CG	7.01	131.43	115.30
1	B	51	GLY	N-CA-C	5.25	126.22	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3514	0	3445	121	0
1	B	3387	0	3314	95	0
2	A	203	0	0	7	0
2	B	185	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7289	0	6759	209	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 15.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:56:TYR:OH	1:A:73:VAL:HG13	1.49	1.12
1:B:56:TYR:OH	1:B:73:VAL:HG13	1.53	1.09
1:A:62:VAL:HG12	1:A:64:ARG:HH12	1.41	0.85
1:B:294:GLY:HA3	1:B:326:GLN:HE22	1.44	0.81
1:A:391:MSE:HA	1:A:401:ILE:HD11	1.65	0.79

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	444/486 (91%)	410 (92%)	33 (7%)	1 (0%)	56	63
1	B	427/486 (88%)	396 (93%)	27 (6%)	4 (1%)	25	20
All	All	871/972 (90%)	806 (92%)	60 (7%)	5 (1%)	33	33

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	51	GLY
1	B	49	ALA
1	B	50	PRO
1	B	47	ASN
1	A	184	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	378/389 (97%)	365 (97%)	13 (3%)	49	57
1	B	364/389 (94%)	350 (96%)	14 (4%)	44	52
All	All	742/778 (95%)	715 (96%)	27 (4%)	45	54

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	387	LEU
1	B	50	PRO
1	B	376	ASN
1	B	18	HIS
1	A	64	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	376	ASN
1	B	153	HIS
1	B	299	ASN
1	A	326	GLN
1	B	271	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	MSO	A	309	1	8,8,9	8.09	5 (62%)	7,9,11	4.75	3 (42%)
1	MSO	B	309	1	8,8,9	8.29	5 (62%)	7,9,11	4.53	4 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MSO	A	309	1	-	1/4/7/9	0/0/0/0
1	MSO	B	309	1	-	1/4/7/9	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	309	MSO	O-C	17.47	1.23	1.11
1	A	309	MSO	O-C	17.05	1.23	1.11
1	B	309	MSO	SE-CE	-11.35	1.77	1.93
1	A	309	MSO	SE-CE	-10.18	1.79	1.93
1	A	309	MSO	SE-OE	-10.14	1.47	1.67

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	309	MSO	OE-SE-CE	11.15	112.97	101.81
1	B	309	MSO	OE-SE-CE	10.01	111.83	101.81
1	B	309	MSO	CE-SE-CG	4.73	103.93	97.61
1	A	309	MSO	CE-SE-CG	3.92	102.84	97.61
1	B	309	MSO	C-CA-N	-3.89	109.94	113.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	309	MSO	CE-SE-CG-CB
1	A	309	MSO	CE-SE-CG-CB

There are no ring outliers.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	451/486 (92%)	0.63	54 (11%) 5 6	25, 43, 79, 93	1 (0%)
1	B	433/486 (89%)	0.52	51 (11%) 5 6	26, 43, 75, 89	1 (0%)
All	All	884/972 (90%)	0.58	105 (11%) 5 6	25, 43, 76, 93	2 (0%)

The worst 5 of 105 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	TYR	8.2
1	B	241	PHE	7.6
1	B	24	ILE	6.9
1	B	363	VAL	6.6
1	A	27	GLU	6.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	MSO	B	309	9/10	0.15	-0.08	41,42,44,44	0
1	MSO	A	309	9/10	0.16	-0.86	41,41,43,44	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

There are no ligands in this entry.

6.5 Other polymers ⓘ

There are no such residues in this entry.