



# wwPDB X-ray Structure Validation Summary Report i

Feb 27, 2014 – 11:20 PM GMT

PDB ID : 4E5G  
Title : Crystal structure of avian influenza virus PAn bound to compound 2  
Authors : DuBois, R.M.; Slavish, P.J.; Webb, T.R.; White, S.W.  
Deposited on : 2012-03-14  
Resolution : 2.65 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2393 (2.68-2.60)
Clashscore	79885	2915 (2.68-2.60)
Ramachandran outliers	78287	2865 (2.68-2.60)
Sidechain outliers	78261	2865 (2.68-2.60)
RSRZ outliers	66119	2393 (2.68-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	187	
1	B	187	
1	C	187	
1	D	187	

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6111 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polymerase protein PA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	B	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	C	181	Total	C	N	O	S	0	0	0
			1489	937	257	284	11			
1	D	179	Total	C	N	O	S	0	0	0
			1475	929	255	280	11			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	50A	GLY	-	LINKER	UNP Q5EP34
A	50B	GLY	-	LINKER	UNP Q5EP34
A	50C	SER	-	LINKER	UNP Q5EP34
A	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
A	197	LEU	-	EXPRESSION TAG	UNP Q5EP34
A	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
A	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
A	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
A	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
A	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
B	50A	GLY	-	LINKER	UNP Q5EP34
B	50B	GLY	-	LINKER	UNP Q5EP34
B	50C	SER	-	LINKER	UNP Q5EP34
B	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
B	197	LEU	-	EXPRESSION TAG	UNP Q5EP34

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
B	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
B	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
B	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
B	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
C	50A	GLY	-	LINKER	UNP Q5EP34
C	50B	GLY	-	LINKER	UNP Q5EP34
C	50C	SER	-	LINKER	UNP Q5EP34
C	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196C	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	196D	GLU	-	EXPRESSION TAG	UNP Q5EP34
C	197	LEU	-	EXPRESSION TAG	UNP Q5EP34
C	198	ALA	-	EXPRESSION TAG	UNP Q5EP34
C	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
C	200	VAL	-	EXPRESSION TAG	UNP Q5EP34
C	201	PRO	-	EXPRESSION TAG	UNP Q5EP34
C	202	ARG	-	EXPRESSION TAG	UNP Q5EP34
D	50A	GLY	-	LINKER	UNP Q5EP34
D	50B	GLY	-	LINKER	UNP Q5EP34
D	50C	SER	-	LINKER	UNP Q5EP34
D	196A	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	196B	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	197	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	198	GLU	-	EXPRESSION TAG	UNP Q5EP34
D	199	LEU	-	EXPRESSION TAG	UNP Q5EP34
D	200	ALA	-	EXPRESSION TAG	UNP Q5EP34
D	201	LEU	-	EXPRESSION TAG	UNP Q5EP34
D	202	VAL	-	EXPRESSION TAG	UNP Q5EP34
D	203	PRO	-	EXPRESSION TAG	UNP Q5EP34
D	204	ARG	-	EXPRESSION TAG	UNP Q5EP34

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Mn	0	0
			2	2		
3	A	2	Total	Mn	0	0
			2	2		
3	D	2	Total	Mn	0	0
			2	2		
3	C	2	Total	Mn	0	0
			2	2		

- Molecule 4 is 2-4-DIOXO-4-PHENYLBUTANOICACID (three-letter code: XI7) (formula:  $C_{10}H_8O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			14	10	4		
4	A	1	Total	C	O	0	0
			14	10	4		
4	B	1	Total	C	O	0	0
			14	10	4		
4	B	1	Total	C	O	0	0
			14	10	4		
4	C	1	Total	C	O	0	0
			14	10	4		
4	C	1	Total	C	O	0	0
			14	10	4		
4	D	1	Total	C	O	0	0
			14	10	4		
4	D	1	Total	C	O	0	0
			14	10	4		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	O	0	0
			4	4		
5	B	1	Total	O	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	1	Total	O	0	0
			1	1		
5	D	3	Total	O	0	0
			3	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Polymerase protein PA

Chain A: 




- Molecule 1: Polymerase protein PA

Chain B: 




- Molecule 1: Polymerase protein PA

Chain C: 



- Molecule 1: Polymerase protein PA

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.89Å 133.66Å 126.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.84 – 2.65 44.84 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.84-2.65) 99.8 (44.84-2.65)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.34 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.243 , 0.285 0.245 , 0.286	Depositor DCC
$R_{free}$ test set	1599 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.9	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 25.8	EDS
Estimated twinning fraction	0.003 for -k,-h,-l	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 31674 reflections	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	6111	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 28.08 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.9629e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, SO4, XI7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/1519	0.58	0/2040
1	B	0.43	0/1519	0.59	0/2040
1	C	0.44	0/1519	0.56	0/2040
1	D	0.43	0/1505	0.57	0/2021
All	All	0.44	0/6062	0.58	0/8141

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1489	0	1450	24	0
1	B	1489	0	1450	30	0
1	C	1489	0	1450	33	0
1	D	1475	0	1439	32	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	28	0	14	1	0
4	B	28	0	14	2	0
4	C	28	0	14	2	0
4	D	28	0	14	2	0
5	A	4	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	3	0	0	0	0
All	All	6111	0	5845	110	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 9.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:D:73:LYS:CD	1:D:73:LYS:H	1.57	1.17
1:B:73:LYS:H	1:B:73:LYS:HE3	1.13	1.12
1:C:73:LYS:CE	1:C:73:LYS:H	1.64	1.09
1:A:73:LYS:HE3	1:A:73:LYS:H	0.90	1.06
1:A:73:LYS:HE3	1:A:73:LYS:N	1.73	1.03

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/187 (96%)	175 (98%)	4 (2%)	0	100	100
1	B	179/187 (96%)	176 (98%)	3 (2%)	0	100	100
1	C	179/187 (96%)	174 (97%)	5 (3%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	177/187 (95%)	174 (98%)	3 (2%)	0	100	100
All	All	714/748 (96%)	699 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/165 (97%)	157 (98%)	3 (2%)	69	90
1	B	160/165 (97%)	155 (97%)	5 (3%)	52	80
1	C	160/165 (97%)	156 (98%)	4 (2%)	60	85
1	D	159/165 (96%)	157 (99%)	2 (1%)	80	94
All	All	639/660 (97%)	625 (98%)	14 (2%)	64	88

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	104	LYS
1	B	124	ARG
1	C	153	GLU
1	B	96	ASN
1	C	96	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	96	ASN
1	D	92	ASN
1	C	92	ASN
1	B	92	ASN
1	C	41	HIS

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 8 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	SO4	A	301	-	4,4,4	0.17	0	6,6,6	0.15	0
2	SO4	A	302	-	4,4,4	0.17	0	6,6,6	0.11	0
4	XI7	A	305	3	14,14,14	2.04	3 (21%)	18,18,18	0.83	1 (5%)
4	XI7	A	306	-	14,14,14	1.96	3 (21%)	18,18,18	0.99	1 (5%)
2	SO4	B	301	-	4,4,4	0.23	0	6,6,6	0.12	0
2	SO4	B	302	-	4,4,4	0.25	0	6,6,6	0.08	0
4	XI7	B	305	3	14,14,14	2.02	3 (21%)	18,18,18	0.92	1 (5%)
4	XI7	B	306	-	14,14,14	1.95	3 (21%)	18,18,18	0.95	1 (5%)
2	SO4	C	301	-	4,4,4	0.20	0	6,6,6	0.10	0
2	SO4	C	302	-	4,4,4	0.18	0	6,6,6	0.08	0
4	XI7	C	305	3	14,14,14	2.02	3 (21%)	18,18,18	0.72	0
4	XI7	C	306	-	14,14,14	1.96	3 (21%)	18,18,18	1.03	1 (5%)
2	SO4	D	301	-	4,4,4	0.20	0	6,6,6	0.13	0
2	SO4	D	302	-	4,4,4	0.22	0	6,6,6	0.11	0
4	XI7	D	305	3	14,14,14	1.96	3 (21%)	18,18,18	0.98	2 (11%)
4	XI7	D	306	-	14,14,14	2.00	3 (21%)	18,18,18	1.37	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	301	-	-	0/0/0/0	0/0/0/0
2	SO4	A	302	-	-	0/0/0/0	0/0/0/0
4	XI7	A	305	3	-	0/12/12/12	0/1/1/1
4	XI7	A	306	-	-	0/12/12/12	0/1/1/1
2	SO4	B	301	-	-	0/0/0/0	0/0/0/0
2	SO4	B	302	-	-	0/0/0/0	0/0/0/0
4	XI7	B	305	3	-	0/12/12/12	0/1/1/1
4	XI7	B	306	-	-	0/12/12/12	0/1/1/1
2	SO4	C	301	-	-	0/0/0/0	0/0/0/0
2	SO4	C	302	-	-	0/0/0/0	0/0/0/0
4	XI7	C	305	3	-	0/12/12/12	0/1/1/1
4	XI7	C	306	-	-	0/12/12/12	0/1/1/1
2	SO4	D	301	-	-	0/0/0/0	0/0/0/0
2	SO4	D	302	-	-	0/0/0/0	0/0/0/0
4	XI7	D	305	3	-	0/12/12/12	0/1/1/1
4	XI7	D	306	-	-	0/12/12/12	0/1/1/1

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	305	XI7	C8-C9	-4.67	1.34	1.51
4	D	306	XI7	C8-C9	-4.64	1.34	1.51
4	B	305	XI7	C8-C9	-4.58	1.34	1.51
4	B	306	XI7	C8-C9	-4.58	1.34	1.51
4	D	306	XI7	O8-C9	4.55	1.33	1.23

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	306	XI7	C9-C8-C7	3.18	126.42	113.39
4	C	306	XI7	C9-C8-C7	3.03	125.79	113.39
4	A	306	XI7	C9-C8-C7	2.74	124.59	113.39
4	B	306	XI7	C9-C8-C7	2.42	123.32	113.39
4	B	305	XI7	C9-C8-C7	2.40	123.21	113.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/187 (96%)	0.20	3 (1%) 67 66	29, 46, 68, 84	0
1	B	181/187 (96%)	0.20	4 (2%) 59 56	31, 47, 73, 86	0
1	C	181/187 (96%)	0.18	2 (1%) 77 79	29, 46, 71, 76	0
1	D	179/187 (95%)	0.22	4 (2%) 59 56	32, 46, 68, 73	0
All	All	722/748 (96%)	0.20	13 (1%) 64 64	29, 46, 70, 86	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	124	ARG	3.7
1	D	141	GLU	2.9
1	D	125	ARG	2.9
1	D	124	ARG	2.8
1	D	7	GLN	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors



of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	302	5/5	0.27	-	99,99,99,100	0
3	MN	C	303	1/1	0.18	-	42,42,42,42	0
3	MN	A	304	1/1	0.17	-	37,37,37,37	0
2	SO4	C	301	5/5	0.17	-	87,88,88,88	0
3	MN	D	303	1/1	0.18	-	44,44,44,44	0
4	XI7	C	305	14/14	0.23	-	42,48,52,52	0
4	XI7	A	305	14/14	0.25	-	53,54,55,55	0
2	SO4	D	302	5/5	0.20	-	116,116,116,116	0
4	XI7	A	306	14/14	0.37	-	64,67,67,67	0
4	XI7	C	306	14/14	0.44	-	67,69,73,74	0
2	SO4	B	301	5/5	0.20	-	101,101,102,102	0
2	SO4	D	301	5/5	0.21	-	100,100,101,101	0
3	MN	D	304	1/1	0.17	-	56,56,56,56	0
2	SO4	C	302	5/5	0.22	-	90,90,91,91	0
4	XI7	B	305	14/14	0.21	-	50,54,57,57	0
4	XI7	D	306	14/14	0.42	-	62,65,66,66	0
3	MN	B	303	1/1	0.19	-	59,59,59,59	0
4	XI7	D	305	14/14	0.24	-	53,57,62,62	0
3	MN	C	304	1/1	0.20	-	53,53,53,53	0
2	SO4	A	301	5/5	0.12	-	82,83,83,83	0
3	MN	B	304	1/1	0.18	-	38,38,38,38	0
2	SO4	A	302	5/5	0.25	-	108,108,108,108	0
4	XI7	B	306	14/14	0.34	-	61,66,69,69	0
3	MN	A	303	1/1	0.24	-	57,57,57,57	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.