



wwPDB X-ray Structure Validation Summary Report

Feb 27, 2014 – 11:47 PM GMT

PDB ID : 2E6B
Title : Crystal structure of the stationary phase survival protein SurE from *Thermus thermophilus* HB8 in complex with magnesium and tungstate
Authors : Iwasaki, W.; Miki, K.
Deposited on : 2006-12-26
Resolution : 2.50 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	2784 (2.50-2.50)
Clashscore	79885	3562 (2.50-2.50)
Ramachandran outliers	78287	3480 (2.50-2.50)
Sidechain outliers	78261	3482 (2.50-2.50)
RSRZ outliers	66119	2785 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	244	
1	B	244	
1	C	244	
1	D	244	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	MG	A	301	-	X
2	MG	B	1301	-	X
2	MG	C	2301	-	X
2	MG	D	3301	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Geometry	Electron density
4	SO4	A	404	-	X
4	SO4	B	1402	-	X
4	SO4	B	1403	-	X
5	GOL	A	501	-	X
5	GOL	B	502	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7414 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 5'-nucleotidase surE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	237	Total	C	N	O	S	0	0	0
			1834	1189	321	321	3			
1	B	233	Total	C	N	O	S	0	0	0
			1810	1175	315	317	3			
1	C	232	Total	C	N	O	S	0	0	0
			1803	1169	316	315	3			
1	D	231	Total	C	N	O	S	0	0	0
			1794	1164	314	313	3			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is TUNGSTATE(VI)ION (three-letter code: WO4) (formula: O₄W).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	W	0	0
			5	4	1		
3	B	1	Total	O	W	0	0
			5	4	1		
3	C	1	Total	O	W	0	0
			5	4	1		
3	D	1	Total	O	W	0	0
			5	4	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	27	Total 27	O 27	0	0
6	B	27	Total 27	O 27	0	0
6	C	16	Total 16	O 16	0	0
6	D	17	Total 17	O 17	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 5'-nucleotidase surE

Chain A: 




• Molecule 1: 5'-nucleotidase surE

Chain B: 




• Molecule 1: 5'-nucleotidase surE


Chain C: 



• Molecule 1: 5'-nucleotidase surE

Chain D: 





4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	132.02Å 132.02Å 130.82Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.46 – 2.50 46.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.46-2.50) 99.9 (46.46-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.46 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.198 , 0.230 0.195 , 0.225	Depositor DCC
R_{free} test set	2293 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 31.0	EDS
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 46021 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7414	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SO4, WO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1893	0.63	0/2594
1	B	0.33	0/1867	0.63	0/2556
1	C	0.32	0/1861	0.60	0/2549
1	D	0.33	0/1852	0.62	0/2537
All	All	0.33	0/7473	0.62	0/10236

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1834	0	1815	66	0
1	B	1810	0	1795	48	0
1	C	1803	0	1782	64	0
1	D	1794	0	1774	52	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	20	0	0	2	0
4	B	15	0	0	1	0
4	C	5	0	0	0	0
4	D	10	0	0	1	0
5	A	6	0	8	4	0
5	B	6	0	8	0	0
6	A	27	0	0	0	0
6	B	27	0	0	2	0
6	C	16	0	0	0	0
6	D	17	0	0	0	0
All	All	7414	0	7182	196	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 14.

The worst 5 of 196 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:B:60:LEU:HD12	1:B:60:LEU:H	1.20	1.04
1:A:231:LEU:H	1:B:177:GLN:NE2	1.58	1.00
1:C:60:LEU:H	1:C:60:LEU:HD23	1.30	0.94
1:A:231:LEU:H	1:B:177:GLN:HE22	1.16	0.90
1:C:238:GLN:HA	1:C:238:GLN:HE21	1.36	0.89

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	233/244 (96%)	228 (98%)	4 (2%)	1 (0%)	43 66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	227/244 (93%)	218 (96%)	8 (4%)	1 (0%)	43	66
1	C	228/244 (93%)	216 (95%)	8 (4%)	4 (2%)	13	20
1	D	227/244 (93%)	223 (98%)	4 (2%)	0	100	100
All	All	915/976 (94%)	885 (97%)	24 (3%)	6 (1%)	30	50

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	132	ASN
1	C	192	PRO
1	C	48	ALA
1	B	132	ASN
1	C	133	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/194 (97%)	186 (98%)	3 (2%)	75	93
1	B	188/194 (97%)	183 (97%)	5 (3%)	57	83
1	C	186/194 (96%)	177 (95%)	9 (5%)	35	60
1	D	185/194 (95%)	182 (98%)	3 (2%)	75	93
All	All	748/776 (96%)	728 (97%)	20 (3%)	57	83

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	60	LEU
1	C	119	LEU
1	C	238	GLN
1	B	209	GLU
1	C	7	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	7	ASN
1	C	100	ASN
1	D	64	HIS
1	B	177	GLN
1	B	238	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	WO4	A	302	1,2	2,4,4	0.16	0	0,6,6	0.00	-
4	SO4	A	401	-	4,4,4	0.16	0	6,6,6	0.12	0
4	SO4	A	402	-	4,4,4	0.29	0	6,6,6	0.06	0
4	SO4	A	403	-	4,4,4	0.30	0	6,6,6	0.06	0
4	SO4	A	404	-	4,4,4	0.34	0	6,6,6	0.08	0
5	GOL	A	501	-	5,5,5	0.13	0	5,5,5	0.29	0
3	WO4	B	1302	1,2	2,4,4	0.23	0	0,6,6	0.00	-
4	SO4	B	1401	-	4,4,4	0.11	0	6,6,6	0.11	0
4	SO4	B	1402	-	4,4,4	0.31	0	6,6,6	0.07	0
4	SO4	B	1403	-	4,4,4	0.85	0	6,6,6	0.19	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	B	502	-	5,5,5	0.30	0	5,5,5	0.35	0
3	WO4	C	2302	1,2	2,4,4	0.22	0	0,6,6	0.00	-
4	SO4	C	2401	-	4,4,4	0.11	0	6,6,6	0.10	0
3	WO4	D	3302	1,2	2,4,4	0.20	0	0,6,6	0.00	-
4	SO4	D	3401	-	4,4,4	0.11	0	6,6,6	0.11	0
4	SO4	D	3402	-	4,4,4	0.31	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WO4	A	302	1,2	-	0/0/0/0	0/0/0/0
4	SO4	A	401	-	-	0/0/0/0	0/0/0/0
4	SO4	A	402	-	-	0/0/0/0	0/0/0/0
4	SO4	A	403	-	-	0/0/0/0	0/0/0/0
4	SO4	A	404	-	-	0/0/0/0	0/0/0/0
5	GOL	A	501	-	-	0/4/4/4	0/0/0/0
3	WO4	B	1302	1,2	-	0/0/0/0	0/0/0/0
4	SO4	B	1401	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1402	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1403	-	-	0/0/0/0	0/0/0/0
5	GOL	B	502	-	-	0/4/4/4	0/0/0/0
3	WO4	C	2302	1,2	-	0/0/0/0	0/0/0/0
4	SO4	C	2401	-	-	0/0/0/0	0/0/0/0
3	WO4	D	3302	1,2	-	0/0/0/0	0/0/0/0
4	SO4	D	3401	-	-	0/0/0/0	0/0/0/0
4	SO4	D	3402	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	237/244 (97%)	-0.17	4 (1%) 67 69	19, 33, 55, 67	0
1	B	233/244 (95%)	-0.21	3 (1%) 74 76	20, 33, 64, 78	0
1	C	232/244 (95%)	-0.12	5 (2%) 59 61	23, 38, 65, 74	0
1	D	231/244 (94%)	-0.22	4 (1%) 67 69	21, 34, 65, 75	0
All	All	933/976 (95%)	-0.18	16 (1%) 67 69	19, 35, 63, 78	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	193	MET	3.3
1	B	132	ASN	3.1
1	A	36	THR	3.0
1	C	192	PRO	2.9
1	B	192	PRO	2.7

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	A	301	1/1	0.22	14.75	38,38,38,38	0
4	SO4	B	1403	5/5	0.31	12.13	67,69,73,77	0
5	GOL	A	501	6/6	0.32	9.53	53,58,60,63	0
5	GOL	B	502	6/6	0.34	9.01	53,56,58,60	0
2	MG	C	2301	1/1	0.24	8.31	49,49,49,49	0
2	MG	B	1301	1/1	0.18	6.16	51,51,51,51	0
4	SO4	B	1402	5/5	0.25	4.26	84,84,85,85	0
2	MG	D	3301	1/1	0.18	3.24	43,43,43,43	0
4	SO4	A	404	5/5	0.25	2.91	78,80,83,83	0
4	SO4	B	1401	5/5	0.15	1.61	36,38,42,42	0
4	SO4	D	3401	5/5	0.14	0.96	40,42,43,46	0
4	SO4	D	3402	5/5	0.18	0.47	82,82,84,84	0
3	WO4	B	1302	5/5	0.12	0.05	41,41,48,55	0
4	SO4	A	401	5/5	0.13	-0.11	36,37,38,40	0
4	SO4	C	2401	5/5	0.13	-0.12	36,36,38,40	0
3	WO4	D	3302	5/5	0.13	-0.23	43,46,53,56	0
3	WO4	C	2302	5/5	0.12	-0.33	33,49,50,53	0
3	WO4	A	302	5/5	0.12	-0.55	39,41,48,49	0
4	SO4	A	402	5/5	0.12	-0.67	92,93,94,94	0
4	SO4	A	403	5/5	0.12	-0.77	69,70,73,73	0

6.5 Other polymers ⓘ

There are no such residues in this entry.