



wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 01:02 PM GMT

PDB ID : 1EA4
Title : TRANSCRIPTIONAL REPRESSOR COPG/22BP DSDNA COMPLEX
Authors : Gomis-Rueth, F.X.; Costa, M.; Sola, M.; Acebo, P.; Eritja, R.; Espinosa, M.;
Solar, G.D.; Coll, M.
Deposited on : 2000-11-05
Resolution : 2.95 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 21963
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	2029 (3.00-2.92)
Ramachandran outliers	78287	1955 (3.00-2.92)
Sidechain outliers	78261	1958 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	45	
1	B	45	
1	D	45	
1	E	45	
1	F	45	
1	G	45	
1	H	45	
1	J	45	
1	K	45	
1	L	45	
2	U	22	
2	W	22	
2	Y	22	
3	V	22	
3	X	22	
3	Z	22	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6013 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRANSCRIPTIONAL REPRESSOR COPG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	42	Total	C	N	O	S	0	0	0
			327	204	57	63	3			
1	B	41	Total	C	N	O	S	0	0	0
			322	202	55	61	4			
1	D	43	Total	C	N	O	S	0	0	0
			335	209	58	64	4			
1	E	44	Total	C	N	O	S	0	0	0
			344	214	59	67	4			
1	F	45	Total	C	N	O	S	0	0	0
			354	220	61	69	4			
1	G	42	Total	C	N	O	S	0	0	0
			326	204	56	62	4			
1	H	44	Total	C	N	O	S	0	0	0
			346	215	60	68	3			
1	J	44	Total	C	N	O	S	0	0	0
			344	214	59	67	4			
1	K	43	Total	C	N	O	S	0	0	0
			335	209	58	64	4			
1	L	44	Total	C	N	O	S	0	0	0
			344	214	59	67	4			

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*AP*AP*CP*CP*GP*TP*GP*CP*AP*CP*TP*CP*AP*AP*TP*GP*CP*AP*AP*TP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	U	20	Total	C	N	O	P	0	0	0
			388	184	74	111	19			
2	W	22	Total	C	N	O	P	0	0	0
			443	213	81	128	21			
2	Y	21	Total	C	N	O	P	0	21	0
			420	203	79	119	19			

- Molecule 3 is a DNA chain called DNA(5'-D(*AP*GP*AP*TP*TP*GP*CP*AP*TP*TP*

GP*AP*GP*TP*GP*CP*AP*CP*GP*GP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	V	21	Total 417	C 197	N 79	O 121	P 20	0	0	0
3	X	22	Total 437	C 207	N 81	O 128	P 21	0	0	0
3	Z	21	Total 430	C 207	N 81	O 123	P 19	0	21	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total 11	O 11	0	0
4	B	3	Total 3	O 3	0	0
4	D	8	Total 8	O 8	0	0
4	E	5	Total 5	O 5	0	0
4	F	6	Total 6	O 6	0	0
4	G	5	Total 5	O 5	0	0
4	H	4	Total 4	O 4	0	0
4	J	5	Total 5	O 5	0	0
4	K	6	Total 6	O 6	0	0
4	L	6	Total 6	O 6	0	0
4	U	11	Total 11	O 11	0	0
4	V	9	Total 9	O 9	0	0
4	W	2	Total 2	O 2	0	0
4	X	8	Total 8	O 8	0	0
4	Y	9	Total 9	O 9	0	0
4	Z	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain A: 



• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain B: 



• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain D: 



• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain E: 



• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain F: 



• Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain G: 



- Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain H: 



- Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain J: 



- Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain K: 




- Molecule 1: TRANSCRIPTIONAL REPRESSOR COPG

Chain L: 




- Molecule 2: DNA (5'-D(*TP*AP*AP*CP*CP*GP*TP*GP*CP*AP*CP*TP*CP*AP*AP*TP*GP*CP*AP*AP*TP*C)-3')

Chain U: 




- Molecule 2: DNA (5'-D(*TP*AP*AP*CP*CP*GP*TP*GP*CP*AP*CP*TP*CP*AP*AP*TP*GP*CP*AP*AP*TP*C)-3')

Chain W: 



- Molecule 2: DNA (5'-D(*TP*AP*AP*CP*CP*GP*TP*GP*CP*AP*CP*TP*CP*AP*AP*TP*GP*CP*AP*AP*TP*C)-3')

Chain Y: 




- Molecule 3: DNA(5'-D(*AP*GP*AP*TP*TP*GP*CP*AP*TP*TP*GP*AP*GP*TP*GP*CP*AP*CP*GP*GP*TP*T)-3')

Chain V: 


A201	G202	A203	T204	T205	G206	C207	A208	T209	T210	G211	A212	G213	T214	G215	C216	A217	C218	G219	G220	T221	DT
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	----

● Molecule 3: DNA(5'-D(*AP*GP*AP*TP*TP*GP*CP*AP*TP*TP*GP*AP*GP*TP*GP*CP*AP*CP*GP*GP*TP*T)-3')

Chain X: 

A201	G202	A203	T204	T205	G206	C207	A208	T209	T210	G211	A212	G213	T214	G215	C216	A217	C218	G219	G220	T221	T222
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

● Molecule 3: DNA(5'-D(*AP*GP*AP*TP*TP*GP*CP*AP*TP*TP*GP*AP*GP*TP*GP*CP*AP*CP*GP*GP*TP*T)-3')

Chain Z: 

A201	G202	A203	T204	T205	G206	C207	A208	T209	T210	G211	A212	G213	T214	G215	C216	A217	C218	G219	G220	T221	DT
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	213.40Å 76.04Å 50.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.95	Depositor
% Data completeness (in resolution range)	96.9 (40.00-2.95)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.230 , 0.307	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6013	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/327	0.71	0/432
1	B	0.63	0/322	0.86	1/425 (0.2%)
1	D	0.53	0/335	0.75	0/442
1	E	0.55	0/344	0.71	0/454
1	F	0.62	1/354 (0.3%)	0.82	1/465 (0.2%)
1	G	0.51	0/326	0.68	0/430
1	H	0.66	1/346 (0.3%)	0.80	1/455 (0.2%)
1	J	0.58	0/344	0.75	0/454
1	K	0.52	0/335	0.71	0/442
1	L	0.50	0/344	0.75	0/454
2	U	0.52	0/435	0.78	0/669
2	W	0.64	0/496	0.82	0/762
2	Y	0.46	0/470	0.80	0/720
3	V	0.57	0/468	0.80	0/723
3	X	0.64	0/490	0.91	0/757
3	Z	0.56	1/482 (0.2%)	0.86	0/742
All	All	0.57	3/6218 (0.0%)	0.79	3/8826 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Z	211[B]	DG	N1-C2	6.49	1.43	1.37
1	F	45	LYS	CE-NZ	-6.16	1.33	1.49
1	H	45	LYS	CE-NZ	-6.12	1.33	1.49

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	45	LYS	CD-CE-NZ	5.64	124.68	111.70
1	F	45	LYS	CD-CE-NZ	5.45	124.23	111.70
1	B	40	LYS	C-N-CA	5.06	134.35	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	327	0	355	30	0
1	B	322	0	356	27	0
1	D	335	0	367	40	0
1	E	344	0	373	39	0
1	F	354	0	386	29	0
1	G	326	0	359	24	0
1	H	346	0	374	41	0
1	J	344	0	373	35	0
1	K	335	0	367	39	0
1	L	344	0	373	34	0
2	U	388	0	213	36	0
2	W	443	0	249	38	0
2	Y	420	0	230	35	0
3	V	417	0	226	40	0
3	X	437	0	238	37	0
3	Z	430	0	224	39	0
4	A	11	0	0	1	0
4	B	3	0	0	0	0
4	D	8	0	0	0	0
4	E	5	0	0	1	0
4	F	6	0	0	1	0
4	G	5	0	0	0	0
4	H	4	0	0	0	0
4	J	5	0	0	0	0
4	K	6	0	0	0	0
4	L	6	0	0	0	0
4	U	11	0	0	1	0
4	V	9	0	0	0	0
4	W	2	0	0	0	0
4	X	8	0	0	0	0
4	Y	9	0	0	0	0
4	Z	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	6013	0	5063	478	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 44.

The worst 5 of 478 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:Y:206[A]:DG:N2	2:Y:220[A]:DA:H2	1.05	1.14
1:B:40:LYS:HA	1:B:40:LYS:HE2	1.33	1.08
1:D:1:MET:HB2	1:E:11:GLU:HB2	1.41	1.03
2:Y:202[A]:DA:H4'	2:Y:203[A]:DA:H5''	1.38	1.01
1:J:15:GLU:O	1:J:19:LYS:HD3	1.64	0.98

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	40/45 (89%)	36 (90%)	4 (10%)	0	100	100
1	B	39/45 (87%)	34 (87%)	4 (10%)	1 (3%)	8	37
1	D	41/45 (91%)	33 (80%)	8 (20%)	0	100	100
1	E	42/45 (93%)	37 (88%)	5 (12%)	0	100	100
1	F	43/45 (96%)	35 (81%)	7 (16%)	1 (2%)	10	41
1	G	40/45 (89%)	36 (90%)	3 (8%)	1 (2%)	9	38
1	H	42/45 (93%)	31 (74%)	11 (26%)	0	100	100
1	J	42/45 (93%)	36 (86%)	6 (14%)	0	100	100
1	K	41/45 (91%)	34 (83%)	6 (15%)	1 (2%)	9	40
1	L	42/45 (93%)	33 (79%)	6 (14%)	3 (7%)	2	7
All	All	412/450 (92%)	345 (84%)	60 (15%)	7 (2%)	14	52

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	44	GLU
1	K	3	LYS
1	L	2	LYS
1	L	3	LYS
1	B	40	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	37/40 (92%)	36 (97%)	1 (3%)	57	90
1	B	37/40 (92%)	32 (86%)	5 (14%)	6	22
1	D	38/40 (95%)	34 (90%)	4 (10%)	10	35
1	E	39/40 (98%)	36 (92%)	3 (8%)	18	54
1	F	40/40 (100%)	36 (90%)	4 (10%)	11	38
1	G	37/40 (92%)	31 (84%)	6 (16%)	3	15
1	H	39/40 (98%)	35 (90%)	4 (10%)	10	36
1	J	39/40 (98%)	34 (87%)	5 (13%)	6	25
1	K	38/40 (95%)	33 (87%)	5 (13%)	6	23
1	L	39/40 (98%)	35 (90%)	4 (10%)	10	36
All	All	383/400 (96%)	342 (89%)	41 (11%)	10	33

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	8	THR
1	G	38	ASN
1	L	3	LYS
1	G	12	SER
1	G	16	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	16	ASN
1	J	16	ASN
1	K	43	GLN
1	G	38	ASN
1	K	16	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.