



# wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 11:21 PM GMT

PDB ID : 1EAV  
Title : CRYSTAL STRUCTURES OF HUMAN GEPHYRIN AND PLANT CNX1 G  
DOMAINS - COMPARATIVE ANALYSIS AND FUNCTIONAL IMPLICA-  
TIONS  
Authors : Schwarz, G.; Schrader, N.; Mendel, R.R.; Hecht, H.J.  
Deposited on : 2001-07-17  
Resolution : 2.60 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	1718 (2.60-2.60)
Clashscore	79885	2154 (2.60-2.60)
Ramachandran outliers	78287	2113 (2.60-2.60)
Sidechain outliers	78261	2113 (2.60-2.60)
RSRZ outliers	66119	1718 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	162	
1	B	162	
1	C	162	
1	D	162	
1	E	162	
1	F	162	
1	G	162	
1	H	162	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9501 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	B	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	C	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	D	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	E	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	F	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			
1	G	160	Total	C	N	O	S	Se	0	0	0
			1183	745	200	231	1	6			
1	H	160	Total	C	N	O	S	Se	0	0	0
			1181	745	200	229	1	6			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
A	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
A	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
A	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
A	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
A	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
B	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
C	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
C	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
C	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
C	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
C	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
D	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
E	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
F	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
G	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	72	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	108	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	109	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	120	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	137	MSE	MET	MODIFIED RESIDUE	UNP Q39054
H	148	MSE	MET	MODIFIED RESIDUE	UNP Q39054

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	10	Total O 10 10	0	0
2	B	7	Total O 7 7	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	5	Total 5	O 5	0	0
2	D	3	Total 3	O 3	0	0
2	E	5	Total 5	O 5	0	0
2	F	8	Total 8	O 8	0	0
2	G	7	Total 7	O 7	0	0
2	H	6	Total 6	O 6	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain A: 




#### • Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain B: 




#### • Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain C: 




#### • Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain D: 




- Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain E:




- Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain F:




- Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain G:



- Molecule 1: MOLYBDOPTERIN BIOSYNTHESIS CNX1 PROTEIN

Chain H:



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	175.30Å 175.30Å 175.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.00 – 2.60 26.43 – 2.60	Depositor EDS
% Data completeness (in resolution range)	90.8 (27.00-2.60) 90.8 (26.43-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.35 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.0.36	Depositor
R, $R_{free}$	0.223 , 0.251 0.209 , 0.217	Depositor DCC
$R_{free}$ test set	2582 reflections (5.43%)	DCC
Wilson B-factor (Å <sup>2</sup> )	58.3	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 12.4	EDS
Estimated twinning fraction	0.457 for l,-k,h	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	17 of 50135 reflections (0.034%)	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9501	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.44	8/1192 (0.7%)	1.32	6/1608 (0.4%)
1	B	1.37	4/1192 (0.3%)	1.30	9/1608 (0.6%)
1	C	1.36	4/1192 (0.3%)	1.36	7/1608 (0.4%)
1	D	1.32	5/1192 (0.4%)	1.36	10/1608 (0.6%)
1	E	1.36	1/1192 (0.1%)	1.30	11/1608 (0.7%)
1	F	1.34	1/1192 (0.1%)	1.29	7/1608 (0.4%)
1	G	1.46	4/1194 (0.3%)	1.40	12/1609 (0.7%)
1	H	1.36	7/1192 (0.6%)	1.31	6/1608 (0.4%)
All	All	1.38	34/9538 (0.4%)	1.33	68/12865 (0.5%)

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	125	ALA	CA-CB	-7.19	1.37	1.52
1	D	71	GLU	CD-OE1	7.03	1.33	1.25
1	B	88	VAL	CA-CB	-6.99	1.40	1.54
1	A	122	SER	CB-OG	6.85	1.51	1.42
1	G	122	SER	CB-OG	6.85	1.51	1.42

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	35	ASP	CB-CG-OD2	12.88	129.90	118.30
1	A	35	ASP	CB-CG-OD2	11.81	128.93	118.30
1	B	35	ASP	CB-CG-OD2	11.21	128.39	118.30
1	C	35	ASP	CB-CG-OD2	10.18	127.46	118.30
1	G	24	ASP	CB-CG-OD2	9.50	126.85	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1181	0	1228	35	0
1	B	1181	0	1228	27	0
1	C	1181	0	1228	31	0
1	D	1181	0	1228	28	0
1	E	1181	0	1228	24	0
1	F	1181	0	1228	32	0
1	G	1183	0	1228	38	0
1	H	1181	0	1228	34	0
2	A	10	0	0	1	0
2	B	7	0	0	0	0
2	C	5	0	0	0	0
2	D	3	0	0	0	0
2	E	5	0	0	0	0
2	F	8	0	0	1	0
2	G	7	0	0	0	0
2	H	6	0	0	1	0
All	All	9501	0	9824	234	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

The worst 5 of 234 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:C:148:MSE:CE	1:C:148:MSE:SE	2.18	1.41
1:A:32:SER:HB2	1:G:35:ASP:OD1	1.65	0.97
1:E:125:ALA:H	1:E:136:ASN:HD22	1.23	0.86
1:C:125:ALA:H	1:C:136:ASN:HD22	1.24	0.85
1:D:125:ALA:H	1:D:136:ASN:HD22	1.23	0.84

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/162 (98%)	152 (96%)	6 (4%)	0	100	100
1	B	158/162 (98%)	152 (96%)	5 (3%)	1 (1%)	33	63
1	C	158/162 (98%)	149 (94%)	9 (6%)	0	100	100
1	D	158/162 (98%)	145 (92%)	11 (7%)	2 (1%)	18	35
1	E	158/162 (98%)	148 (94%)	9 (6%)	1 (1%)	33	63
1	F	158/162 (98%)	150 (95%)	7 (4%)	1 (1%)	33	63
1	G	158/162 (98%)	151 (96%)	6 (4%)	1 (1%)	33	63
1	H	158/162 (98%)	148 (94%)	9 (6%)	1 (1%)	33	63
All	All	1264/1296 (98%)	1195 (94%)	62 (5%)	7 (1%)	33	63

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	130	GLY
1	B	14	SER
1	D	14	SER
1	E	14	SER
1	F	14	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	131/127 (103%)	126 (96%)	5 (4%)	44	74
1	B	131/127 (103%)	125 (95%)	6 (5%)	37	66
1	C	131/127 (103%)	126 (96%)	5 (4%)	44	74

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	131/127 (103%)	126 (96%)	5 (4%)	44	74
1	E	131/127 (103%)	127 (97%)	4 (3%)	52	81
1	F	131/127 (103%)	125 (95%)	6 (5%)	37	66
1	G	131/127 (103%)	125 (95%)	6 (5%)	37	66
1	H	131/127 (103%)	127 (97%)	4 (3%)	52	81
All	All	1048/1016 (103%)	1007 (96%)	41 (4%)	43	74

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	36	SER
1	E	32	SER
1	H	23	PRO
1	D	124	SER
1	D	131	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	136	ASN
1	H	136	ASN
1	F	136	ASN
1	C	136	ASN
1	E	136	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	160/162 (98%)	-0.04	3 (1%) 64 61	33, 47, 77, 104	0
1	B	160/162 (98%)	-0.01	3 (1%) 64 61	34, 47, 78, 105	0
1	C	160/162 (98%)	-0.01	4 (2%) 54 52	35, 47, 77, 107	0
1	D	160/162 (98%)	-0.03	2 (1%) 74 75	35, 47, 78, 104	0
1	E	160/162 (98%)	0.03	4 (2%) 54 52	35, 48, 77, 102	0
1	F	160/162 (98%)	-0.04	2 (1%) 74 75	34, 47, 79, 106	0
1	G	160/162 (98%)	-0.02	2 (1%) 74 75	34, 45, 77, 104	0
1	H	160/162 (98%)	-0.03	2 (1%) 74 75	36, 49, 79, 104	0
All	All	1280/1296 (98%)	-0.02	22 (1%) 67 66	33, 47, 78, 107	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	3	GLY	12.4
1	D	3	GLY	9.1
1	G	3	GLY	8.6
1	F	4	PRO	8.5
1	B	3	GLY	7.3

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.