



# wwPDB X-ray Structure Validation Summary Report

Feb 28, 2014 – 10:54 PM GMT

PDB ID : 1EAY  
Title : CHEY-BINDING (P2) DOMAIN OF CHEA IN COMPLEX WITH CHEY FROM ESCHERICHIA COLI  
Authors : Mcevoy, M.M.; Hausrath, A.C.; Randolph, G.B.; Remington, S.J.; Dahlquist, F.W.  
Deposited on : 1998-04-23  
Resolution : 2.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	6188 (2.00-2.00)
Ramachandran outliers	78287	6102 (2.00-2.00)
Sidechain outliers	78261	6100 (2.00-2.00)
RSRZ outliers	66119	4890 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	128	
1	B	128	
2	C	74	
2	D	74	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3058 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHEY.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	128	Total	C	N	O	S	0	0	0
			953	612	154	181	6			
1	B	128	Total	C	N	O	S	0	0	0
			966	617	158	185	6			

- Molecule 2 is a protein called CHEA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	67	Total	C	N	O	S	0	0	0
			501	317	80	103	1			
2	D	69	Total	C	N	O	S	0	0	0
			514	325	82	106	1			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	54	Total	O	0	0
			54	54		
3	B	32	Total	O	0	0
			32	32		
3	C	27	Total	O	0	0
			27	27		
3	D	11	Total	O	0	0
			11	11		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: CHEY

Chain A:




#### • Molecule 1: CHEY

Chain B:




#### • Molecule 2: CHEA

Chain C:



#### • Molecule 2: CHEA

Chain D:



D219	
T225	
Y226	
GLU	
VAL	

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.50Å 64.20Å 158.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 21.21 – 1.95	Depositor EDS
% Data completeness (in resolution range)	78.0 (20.00-2.00) 74.3 (21.21-1.95)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	10.71 (at 1.95Å)	Xtriage
Refinement program	TNT V. 5-F	Depositor
R, $R_{free}$	0.217 , (Not available) 0.249 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	8.3	Xtriage
Anisotropy	1.127	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 170.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning	$\langle  L  \rangle = 0.30$ , $\langle L^2 \rangle = 0.13$	Xtriage
Outliers	0 of 30701 reflections	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	3058	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.37	8/965 (0.8%)	1.58	15/1302 (1.2%)
1	B	1.32	7/978 (0.7%)	1.57	11/1318 (0.8%)
2	C	1.41	5/505 (1.0%)	1.72	17/687 (2.5%)
2	D	1.41	5/518 (1.0%)	1.79	20/706 (2.8%)
All	All	1.37	25/2966 (0.8%)	1.64	63/4013 (1.6%)

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	27	GLU	CD-OE2	6.55	1.32	1.25
1	A	67	GLU	CD-OE1	6.47	1.32	1.25
2	C	178	GLU	CD-OE1	6.44	1.32	1.25
2	D	178	GLU	CD-OE1	6.43	1.32	1.25
2	D	217	GLU	CD-OE1	6.38	1.32	1.25

The worst 5 of 63 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	207	ASP	CB-CG-OD1	9.63	126.97	118.30
1	B	38	ASP	CB-CG-OD1	8.45	125.91	118.30
1	B	38	ASP	CB-CG-OD2	-8.08	111.03	118.30
1	A	18	ARG	NE-CZ-NH1	7.84	124.22	120.30
1	A	38	ASP	CB-CG-OD1	7.65	125.19	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit,

and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	953	0	957	96	0
1	B	966	0	977	109	0
2	C	501	0	505	44	0
2	D	514	0	518	36	0
3	A	54	0	0	6	0
3	B	32	0	0	4	0
3	C	27	0	0	0	0
3	D	11	0	0	0	0
All	All	3058	0	2957	277	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 47.

The worst 5 of 277 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:54:VAL:HG12	1:A:56:SER:HB2	1.48	0.94
1:B:122:LYS:HE3	2:D:215:VAL:HG13	1.53	0.90
2:C:175:LEU:HD11	2:C:216:ILE:HD13	1.53	0.89
2:C:166:ARG:NH2	2:C:220:GLN:HA	1.88	0.87
2:D:179:LEU:HB2	2:D:185:LEU:HD11	1.57	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/128 (98%)	119 (94%)	6 (5%)	1 (1%)	27	17
1	B	126/128 (98%)	112 (89%)	11 (9%)	3 (2%)	9	3
2	C	65/74 (88%)	61 (94%)	4 (6%)	0	100	100
2	D	67/74 (90%)	64 (96%)	3 (4%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	384/404 (95%)	356 (93%)	24 (6%)	4 (1%)	22	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	4	LYS
1	B	79	SER
1	A	79	SER
1	B	76	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/102 (93%)	84 (88%)	11 (12%)	8	4
1	B	99/102 (97%)	85 (86%)	14 (14%)	5	2
2	C	55/64 (86%)	48 (87%)	7 (13%)	6	3
2	D	57/64 (89%)	49 (86%)	8 (14%)	5	2
All	All	306/332 (92%)	266 (87%)	40 (13%)	6	3

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	47	GLN
1	B	111	PHE
2	D	199	LEU
1	B	91	LYS
1	B	126	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	B	23	ASN
1	B	32	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	44	ASN
1	B	121	ASN

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

### 5.7 Other polymers ⓘ

There are no such residues in this entry.

### 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	128/128 (100%)	-0.24	2 (1%) 68 69	17, 33, 65, 83	0
1	B	128/128 (100%)	-0.24	0 100 100	20, 32, 70, 85	0
2	C	67/74 (90%)	-0.28	2 (2%) 48 47	21, 36, 55, 75	0
2	D	69/74 (93%)	-0.18	1 (1%) 72 72	23, 37, 64, 78	0
All	All	392/404 (97%)	-0.24	5 (1%) 74 74	17, 35, 67, 85	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	ALA	2.7
2	C	169	ALA	2.7
1	A	3	ASP	2.2
2	D	169	ALA	2.2
2	C	225	THR	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers ⓘ

There are no such residues in this entry.