



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 03:43 AM GMT

PDB ID : 3EAX
Title : Crystal structure PTP1B complex with small molecule compound LZIP-6
Authors : Zhang, Z.-Y.; Liu, S.; Zhang, L.-F.; Yu, X.; Xue, T.; Gunawan, A.M.; Long, Y.-Q.
Deposited on : 2008-08-26
Resolution : 1.90 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	dev-1323
EDS	:	stable22639
Percentile statistics	:	21963
Refmac	:	5.8.0049
CCP4	:	6.3.0 (Settle)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	3684 (1.90-1.90)
Clashscore	79885	4465 (1.90-1.90)
Ramachandran outliers	78287	4413 (1.90-1.90)
Sidechain outliers	78261	4414 (1.90-1.90)
RSRZ outliers	66119	3686 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	321	

The following table lists non-polymeric compounds that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Geometry	Electron density
2	LZP	A	322	-	X

2 Entry composition i


There are 3 unique types of molecules in this entry. The entry contains 2434 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	282	2296	1458	397	425	16	24	0	0

- Molecule 2 is 4,4'-PIPERAZINE-1,4-DIYLBIS{1-[3-(BENZYLOXY)PHENYL]-4-OXOBUTANE-1,3-DIONE} (three-letter code: LZP) (formula: C₃₈H₃₄N₂O₈).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	48	38	2	8	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	90	Total	O	0	0
			90	90		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.61Å 88.61Å 104.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.90 36.01 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.90) 86.7 (36.01-1.60)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.244 0.217 , 0.246	Depositor DCC
R_{free} test set	1838 reflections (5.07%)	DCC
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 56.6	EDS
Estimated twinning fraction	0.023 for -h,-k,l	Xtriage
L-test for twinning	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 54313 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2434	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2348	0.56	0/3164

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2296	0	2272	51	0
2	A	48	0	34	4	0
3	A	90	0	0	1	0
All	All	2434	0	2306	51	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 11.

All (51) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:246:ILE:HD12	1:A:246:ILE:H	1.27	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:161:GLU:HG3	1:A:168:THR:HG22	1.54	0.90
1:A:219:ILE:HD12	1:A:262:GLN:NE2	2.03	0.73
1:A:3:MET:HE2	1:A:275:ILE:HG13	1.70	0.73
1:A:221:ARG:HE	2:A:322:LZP:H6	1.58	0.67
1:A:281:ILE:HG22	1:A:282:MET:HG3	1.76	0.67
1:A:220:GLY:HA3	2:A:322:LZP:H7A	1.79	0.64
1:A:34:VAL:HG11	1:A:56:ARG:NE	2.15	0.62
1:A:34:VAL:HG12	1:A:53:ASP:OD1	1.99	0.62
1:A:116:LYS:HE3	1:A:181:ASP:O	2.04	0.57
1:A:79:ARG:CZ	1:A:233:LEU:HD11	2.36	0.56
1:A:193:ASN:O	1:A:197:LYS:HG2	2.05	0.56
1:A:15:SER:O	1:A:19:ILE:HG12	2.08	0.54
1:A:72:ILE:HD12	1:A:83:LEU:HD13	1.90	0.53
1:A:162:ASN:OD1	1:A:164:THR:HB	2.09	0.52
1:A:234:LEU:O	1:A:238:ARG:HG2	2.09	0.52
1:A:34:VAL:HG11	1:A:56:ARG:CD	2.39	0.51
1:A:45:ARG:HB2	1:A:88:LEU:HD21	1.92	0.51
1:A:79:ARG:HG2	1:A:81:TYR:CZ	2.46	0.50
1:A:219:ILE:HG13	1:A:220:GLY:N	2.26	0.50
1:A:77:ALA:HB2	1:A:234:LEU:HD13	1.95	0.49
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.78	0.49
1:A:260:LEU:O	1:A:261:ILE:HB	2.13	0.48
1:A:100:TRP:CZ2	1:A:162:ASN:HB2	2.48	0.48
1:A:116:LYS:HE2	1:A:182:PHE:CZ	2.49	0.47
1:A:23:ILE:HD13	1:A:250:LEU:HD23	1.96	0.47
1:A:62:GLU:O	1:A:64:ASN:N	2.47	0.47
1:A:130:GLU:CD	1:A:130:GLU:H	2.19	0.46
1:A:246:ILE:CD1	1:A:246:ILE:H	2.05	0.46
1:A:172:LEU:HD23	1:A:201:SER:HB2	1.97	0.46
1:A:161:GLU:CG	1:A:168:THR:HG22	2.37	0.46
1:A:216:SER:HB2	2:A:322:LZP:H10	1.98	0.45
1:A:21:GLN:OE1	1:A:24:ARG:NH2	2.49	0.45
1:A:39:LYS:HE3	3:A:334:HOH:O	2.17	0.44
1:A:205:SER:HB3	1:A:207:GLU:OE1	2.18	0.44
1:A:161:GLU:HG3	1:A:168:THR:CG2	2.36	0.44
1:A:221:ARG:NE	2:A:322:LZP:H6	2.29	0.43
1:A:116:LYS:HE2	1:A:182:PHE:CE1	2.53	0.43
1:A:259:GLY:O	1:A:260:LEU:C	2.57	0.42
1:A:141:LYS:HB2	1:A:163:LEU:HD11	2.01	0.42
1:A:162:ASN:O	1:A:166:GLN:N	2.51	0.42
1:A:245:ASP:HB3	1:A:248:LYS:HB2	2.01	0.42
1:A:140:LEU:HD23	1:A:162:ASN:HA	2.00	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:46:TYR:CE2	1:A:217:ALA:HB2	2.56	0.41
1:A:165:THR:O	1:A:166:GLN:HB2	2.20	0.41
1:A:153:TYR:CE2	1:A:155:VAL:HG23	2.56	0.41
1:A:34:VAL:HG13	1:A:35:ALA:N	2.35	0.41
1:A:235:MET:HG3	1:A:244:VAL:HG21	2.02	0.41
1:A:45:ARG:NH2	1:A:88:LEU:HD22	2.36	0.40
1:A:2:GLU:HG3	1:A:4:GLU:H	1.86	0.40
1:A:73:LYS:HB2	1:A:73:LYS:NZ	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	280/321 (87%)	268 (96%)	10 (4%)	2 (1%)	30 15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	ASP
1	A	261	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	254/294 (86%)	240 (94%)	14 (6%)	30 16

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	11	ASP
1	A	22	ASP
1	A	26	GLU
1	A	59	LEU
1	A	105	ARG
1	A	130	GLU
1	A	215	CYS
1	A	234	LEU
1	A	236	ASP
1	A	238	ARG
1	A	248	LYS
1	A	265	ASP
1	A	266	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	LZP	A	322	-	52,52,52	4.09	38 (73%)	70,70,70	1.30	9 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LZP	A	322	-	-	0/40/52/52	0/5/5/5

All (38) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	322	LZP	C6-C5	8.55	1.58	1.38
2	A	322	LZP	C33-C34	8.52	1.58	1.38
2	A	322	LZP	C3-C2	6.90	1.57	1.37
2	A	322	LZP	C37-C36	6.88	1.57	1.37
2	A	322	LZP	C13-C12	6.05	1.48	1.39
2	A	322	LZP	C9-C8	5.97	1.51	1.38
2	A	322	LZP	C11-C12	5.97	1.49	1.39
2	A	322	LZP	C26-C27	5.89	1.48	1.39
2	A	322	LZP	C28-C27	5.84	1.49	1.39
2	A	322	LZP	C30-C31	5.71	1.50	1.38
2	A	322	LZP	C17-N1	5.63	1.44	1.34
2	A	322	LZP	C13-C8	5.60	1.49	1.38
2	A	322	LZP	C26-C31	5.48	1.49	1.38
2	A	322	LZP	C4-C5	4.73	1.49	1.38
2	A	322	LZP	C19-N2	4.70	1.55	1.47
2	A	322	LZP	C18-N1	4.60	1.55	1.47
2	A	322	LZP	C35-C34	4.55	1.49	1.38
2	A	322	LZP	C21-N1	4.24	1.54	1.47
2	A	322	LZP	C3-C4	3.92	1.48	1.39
2	A	322	LZP	C22-N2	3.81	1.41	1.34
2	A	322	LZP	C36-C35	3.73	1.47	1.39
2	A	322	LZP	C10-C9	3.60	1.47	1.39
2	A	322	LZP	C29-C30	3.50	1.47	1.39
2	A	322	LZP	C29-C28	3.39	1.46	1.39
2	A	322	LZP	C38-C33	3.35	1.46	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	322	LZP	C7-C5	3.34	1.58	1.50
2	A	322	LZP	C1-C6	3.33	1.46	1.39
2	A	322	LZP	C10-C11	3.12	1.46	1.39
2	A	322	LZP	O1-C8	3.11	1.44	1.37
2	A	322	LZP	C32-C34	2.93	1.57	1.50
2	A	322	LZP	C2-C1	2.92	1.46	1.37
2	A	322	LZP	C38-C37	2.90	1.46	1.37
2	A	322	LZP	C20-N2	2.82	1.52	1.47
2	A	322	LZP	O8-C31	2.74	1.44	1.37
2	A	322	LZP	C27-C25	2.70	1.54	1.49
2	A	322	LZP	C12-C14	2.37	1.53	1.49
2	A	322	LZP	C19-C18	2.15	1.60	1.51
2	A	322	LZP	C17-C16	2.06	1.56	1.53

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	322	LZP	C9-C8-C13	-3.71	115.23	120.56
2	A	322	LZP	C30-C31-C26	-3.61	115.37	120.56
2	A	322	LZP	C27-C26-C31	3.45	123.63	119.61
2	A	322	LZP	C12-C13-C8	3.29	123.45	119.61
2	A	322	LZP	C7-O1-C8	2.68	124.82	117.66
2	A	322	LZP	C16-C17-N1	2.38	121.68	119.11
2	A	322	LZP	O6-C23-C22	2.29	122.12	119.34
2	A	322	LZP	C32-O8-C31	2.28	123.75	117.66
2	A	322	LZP	O1-C7-C5	2.15	115.73	109.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/321 (87%)	0.04	9 (3%) 45 46	25, 39, 63, 79	8 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	62	GLU	5.3
1	A	63	ASP	4.7
1	A	152	TYR	4.6
1	A	7	PHE	2.9
1	A	213	VAL	2.6
1	A	2	GLU	2.5
1	A	283	GLY	2.1
1	A	47	ARG	2.0
1	A	14	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	LZP	A	322	48/48	0.24	3.29	45,54,58,59	0

6.5 Other polymers ⓘ

There are no such residues in this entry.