



wwPDB X-ray Structure Validation Summary Report i

Feb 28, 2014 – 05:57 AM GMT

PDB ID : 1EBO
Title : CRYSTAL STRUCTURE OF THE EBOLA VIRUS MEMBRANE-FUSION SUBUNIT, GP2, FROM THE ENVELOPE GLYCOPROTEIN ECTODOMAIN
Authors : Weissenhorn, W.; Carfi, A.; Lee, K.H.; Skehel, J.J.; Wiley, D.C.
Deposited on : 1998-11-03
Resolution : 3.00 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1594 (3.00-3.00)
Ramachandran outliers	78287	1537 (3.00-3.00)
Sidechain outliers	78261	1540 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	131	
1	B	131	
1	C	131	
1	D	131	
1	E	131	
1	F	131	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5331 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	112	Total	C	N	O	S	0	0	0
			901	571	159	169	2			
1	B	112	Total	C	N	O	S	0	0	0
			881	559	151	169	2			
1	C	111	Total	C	N	O	S	0	0	0
			893	567	158	166	2			
1	D	112	Total	C	N	O	S	0	0	0
			876	556	149	169	2			
1	E	111	Total	C	N	O	S	0	0	0
			881	560	153	166	2			
1	F	113	Total	C	N	O	S	0	0	0
			894	568	154	170	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLU	MISSING FROM SEQUENCE	UNP Q913A3
B	?	-	GLU	MISSING FROM SEQUENCE	gb 4389202
C	?	-	GLU	MISSING FROM SEQUENCE	GB 4389203
D	?	-	GLU	MISSING FROM SEQUENCE	GB 4389204
E	?	-	GLU	MISSING FROM SEQUENCE	GB 4389205
F	?	-	GLU	MISSING FROM SEQUENCE	GB 4389206

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	F	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2

Chain A: 




- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2

Chain B: 



- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2

Chain C: 



- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2


Chain D: 






- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2

Chain E:



- Molecule 1: EBOLA VIRUS ENVELOPE PROTEIN CHIMERA CONSISTING OF A FRAGMENT OF GCN4 ZIPPER CLONED N-TERMINAL TO A FRAGMENT OF GP2

Chain F:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	171.70Å 32.69Å 168.86Å 90.00° 119.23° 90.00°	Depositor
Resolution (Å)	20.00 – 3.00	Depositor
% Data completeness (in resolution range)	99.3 (20.00-3.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.239 , 0.256	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5331	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	2/915 (0.2%)	0.83	4/1237 (0.3%)
1	B	0.65	2/895 (0.2%)	0.83	1/1215 (0.1%)
1	C	0.42	0/907	0.70	0/1226
1	D	0.58	3/890 (0.3%)	0.79	3/1208 (0.2%)
1	E	0.73	2/895 (0.2%)	0.74	0/1211
1	F	0.42	0/908	0.79	1/1230 (0.1%)
All	All	0.57	9/5410 (0.2%)	0.78	9/7327 (0.1%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	108	ARG	CD-NE	-16.16	1.19	1.46
1	B	73	ARG	CD-NE	-12.20	1.25	1.46
1	A	131	ASP	C-N	-8.57	1.14	1.34
1	E	27	LYS	CB-CG	-7.80	1.31	1.52
1	D	55	GLU	CB-CG	-7.37	1.38	1.52

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	73	ARG	CD-NE-CZ	13.99	143.18	123.60
1	D	55	GLU	CA-CB-CG	10.20	135.84	113.40
1	A	52	GLY	N-CA-C	-6.93	95.78	113.10
1	A	27	LYS	CB-CA-C	6.58	123.56	110.40
1	D	51	ASP	CA-CB-CG	-5.93	100.35	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	901	0	894	75	0
1	B	881	0	851	64	0
1	C	893	0	891	84	0
1	D	876	0	842	63	0
1	E	881	0	869	62	0
1	F	894	0	875	58	0
2	A	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
3	B	1	0	0	1	0
3	D	1	0	0	1	0
All	All	5331	0	5222	307	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 29.

The worst 5 of 307 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:53:LEU:HD23	1:C:53:LEU:HD22	1.42	0.99
1:C:126:ILE:HA	1:C:130:VAL:CG1	1.98	0.93
1:F:32:ALA:O	1:F:53:LEU:HG	1.67	0.93
1:F:3:GLN:O	1:F:3:GLN:HG2	1.65	0.92
1:E:26:LYS:HG3	1:F:25:ILE:HD11	1.49	0.92

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	110/131 (84%)	103 (94%)	5 (4%)	2 (2%)	13	53
1	B	110/131 (84%)	99 (90%)	10 (9%)	1 (1%)	25	73
1	C	109/131 (83%)	102 (94%)	6 (6%)	1 (1%)	25	73
1	D	110/131 (84%)	103 (94%)	6 (6%)	1 (1%)	25	73
1	E	109/131 (83%)	102 (94%)	7 (6%)	0	100	100
1	F	111/131 (85%)	101 (91%)	9 (8%)	1 (1%)	25	73
All	All	659/786 (84%)	610 (93%)	43 (6%)	6 (1%)	25	73

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	32	ALA
1	D	32	ALA
1	A	29	ILE
1	F	31	GLU
1	B	29	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/117 (82%)	78 (81%)	18 (19%)	2	12
1	B	92/117 (79%)	74 (80%)	18 (20%)	2	11
1	C	95/117 (81%)	75 (79%)	20 (21%)	1	8
1	D	91/117 (78%)	74 (81%)	17 (19%)	2	12
1	E	93/117 (80%)	75 (81%)	18 (19%)	2	11
1	F	94/117 (80%)	83 (88%)	11 (12%)	8	31
All	All	561/702 (80%)	459 (82%)	102 (18%)	2	13

5 of 102 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	60	LEU
1	D	4	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	53	LEU
1	C	65	THR
1	C	84	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	94	GLN
1	D	20	ASN
1	F	62	ASN
1	D	17	HIS
1	D	62	ASN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.