



# Full wwPDB X-ray Structure Validation Report

Feb 27, 2014 – 06:55 AM GMT

PDB ID : 3ENC  
Title : Crystal structure of Pyrococcus furiosus PCC1 dimer  
Authors : Neculai, D.  
Deposited on : 2008-09-25  
Resolution : 2.63 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>

---


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.15 2013  
Xtriage (Phenix) : dev-1323  
EDS : stable22639  
Percentile statistics : 21963  
Refmac : 5.8.0049  
CCP4 : 6.3.0 (Settle)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)  
Validation Pipeline (wwPDB-VP) : stable22683

# 1 Overall quality at a glance

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	66092	2393 (2.68-2.60)
Clashscore	79885	2915 (2.68-2.60)
Ramachandran outliers	78287	2865 (2.68-2.60)
Sidechain outliers	78261	2865 (2.68-2.60)
RSRZ outliers	66119	2393 (2.68-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	87	
1	B	87	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1276 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called protein PCC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	79	Total	C	N	O	Se	0	0	0
			638	412	108	117	1			
1	B	79	Total	C	N	O	Se	0	0	0
			638	412	108	117	1			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	EXPRESSION TAG	UNP Q8TZI1
A	-3	ALA	-	EXPRESSION TAG	UNP Q8TZI1
A	-2	MSE	-	EXPRESSION TAG	UNP Q8TZI1
A	-1	ASP	-	EXPRESSION TAG	UNP Q8TZI1
A	0	PRO	-	EXPRESSION TAG	UNP Q8TZI1
A	12	MSE	ILE	ENGINEERED	UNP Q8TZI1
B	-4	GLY	-	EXPRESSION TAG	UNP Q8TZI1
B	-3	ALA	-	EXPRESSION TAG	UNP Q8TZI1
B	-2	MSE	-	EXPRESSION TAG	UNP Q8TZI1
B	-1	ASP	-	EXPRESSION TAG	UNP Q8TZI1
B	0	PRO	-	EXPRESSION TAG	UNP Q8TZI1
B	12	MSE	ILE	ENGINEERED	UNP Q8TZI1

i

- Molecule 1: protein PCC1

- Molecule 1: protein PCC1

GLY  
ALA  
MSE  
ASP  
PRO  
MSE  
K2  
L3  
K4  
R5  
M12  
P15  
S16  
E17  
D18  
V23  
Y24  
V27  
L28  
H31  
Y36  
R37  
E40  
I41  
D42  
F43  
K44  
L45  
K48  
K55  
D58  
S59  
L62  
T65  
V66  
N67  
S68  
Y69  
L70  
R71  
I72  
W73  
L80  
GLU

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.81Å 78.81Å 60.02Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	68.25 – 2.63 68.25 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.8 (68.25-2.63) 97.1 (68.25-2.50)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.62 (at 2.51Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.192 , 0.222 0.233 , 0.253	Depositor DCC
$R_{free}$ test set	507 reflections (7.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.7	Xtriage
Anisotropy	0.268	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 111.7	EDS
Estimated twinning fraction	0.503 for h,-h-k,-l 0.499 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.503 for h,-h-k,-l	Depositor
L-test for twinning	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Outliers	0 of 7234 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1276	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	124.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/647	0.62	0/869
1	B	0.43	0/647	0.56	0/869
All	All	0.45	0/1294	0.59	0/1738

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	638	0	0	12	0
1	B	638	0	0	11	0
All	All	1276	0	0	20	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 16.

All (20) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:43:PHE:CZ	1:A:50:ILE:CG2	2.74	0.70
1:B:66:VAL:O	1:B:70:LEU:N	2.36	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:60:SER:C	1:A:62:LEU:N	2.57	0.58
1:A:67:ASN:ND2	1:B:67:ASN:OD1	2.39	0.55
1:A:13:GLU:O	1:B:4:LYS:N	2.40	0.55
1:A:69:TYR:CD1	1:A:69:TYR:N	2.74	0.55
1:B:27:VAL:O	1:B:31:HIS:N	2.42	0.52
1:A:24:TYR:CG	1:A:45:LEU:CB	2.93	0.51
1:A:17:GLU:O	1:A:21:LYS:N	2.44	0.50
1:A:11:GLU:OE1	1:A:49:LYS:NZ	2.45	0.50
1:B:31:HIS:ND1	1:B:69:TYR:OH	2.46	0.49
1:B:16:SER:C	1:B:18:ASP:N	2.66	0.48
1:A:10:ILE:N	1:A:52:LEU:O	2.49	0.45
1:A:71:ARG:O	1:A:75:ALA:N	2.50	0.45
1:B:5:ARG:CB	1:B:55:LYS:NZ	2.81	0.43
1:B:23:VAL:O	1:B:24:TYR:C	2.55	0.42
1:A:63:ARG:NH1	1:B:71:ARG:NH1	2.69	0.40
1:B:28:LEU:CD2	1:B:43:PHE:CD2	3.04	0.40
1:B:58:ASP:O	1:B:59:SER:C	2.60	0.40
1:A:34:VAL:CG2	1:A:34:VAL:O	2.69	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	77/87 (88%)	67 (87%)	8 (10%)	2 (3%)	8	13
1	B	77/87 (88%)	69 (90%)	7 (9%)	1 (1%)	18	34
All	All	154/174 (88%)	136 (88%)	15 (10%)	3 (2%)	12	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	15	PRO
1	A	15	PRO

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	36	TYR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	69/72 (96%)	52 (75%)	17 (25%)	1	1
1	B	69/72 (96%)	57 (83%)	12 (17%)	3	4
All	All	138/144 (96%)	109 (79%)	29 (21%)	1	2

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	4	LYS
1	A	5	ARG
1	A	13	GLU
1	A	17	GLU
1	A	28	LEU
1	A	39	SER
1	A	45	LEU
1	A	48	LYS
1	A	55	LYS
1	A	59	SER
1	A	62	LEU
1	A	65	THR
1	A	68	SER
1	A	69	TYR
1	A	73	ILE
1	A	77	ILE
1	B	2	LYS
1	B	4	LYS
1	B	5	ARG
1	B	12	MSE
1	B	37	ARG
1	B	45	LEU
1	B	48	LYS

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	55	LYS
1	B	59	SER
1	B	62	LEU
1	B	68	SER
1	B	73	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA chains in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	79/87 (90%)	0.70	5 (6%) 19 17	74, 118, 172, 192	0
1	B	79/87 (90%)	0.72	7 (8%) 10 7	73, 117, 192, 225	0
All	All	158/174 (90%)	0.71	12 (7%) 14 11	73, 117, 183, 225	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	36	TYR	3.9
1	A	4	LYS	3.2
1	B	65	THR	2.9
1	A	23	VAL	2.7
1	A	48	LYS	2.7
1	B	12	MSE	2.6
1	B	37	ARG	2.3
1	B	41	ILE	2.2
1	B	40	GLU	2.2
1	A	36	TYR	2.2
1	A	39	SER	2.1
1	B	2	LYS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.