



Full wwPDB X-ray Structure Validation Report

Feb 28, 2014 – 09:17 PM GMT

PDB ID : 1EO2
Title : CRYSTAL STRUCTURE OF ACINETOBACTER SP. ADP1 PROTOCAT-
ECHUATE 3,4-DIOXYGENASE
Authors : Vetting, M.W.; D'Argenio, D.A.; Ornston, L.N.; Ohlendorf, D.H.
Deposited on : 2000-03-21
Resolution : 2.25 Å(reported)

This is a full wwPDB validation report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.15 2013
Xtriage (Phenix) : dev-1323
EDS : stable22639
Percentile statistics : 21963
Refmac : 5.8.0049
CCP4 : 6.3.0 (Settle)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : stable22683

1 Overall quality at a glance



The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	66092	1108 (2.28-2.24)
Clashscore	79885	1326 (2.28-2.24)
Ramachandran outliers	78287	1291 (2.28-2.24)
Sidechain outliers	78261	1291 (2.28-2.24)
RSRZ outliers	66119	1110 (2.28-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Mol	Chain	Length	Quality of chain
1	A	209	
2	B	241	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3712 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	0	0	0
			1602	1010	285	307			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	GLY	SER	SEE REMARK 999	UNP P20371

- Molecule 2 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	0	0
			1926	1224	345	352	5			

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Fe	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	66	Total	O	0	0
			66	66		
4	B	117	Total	O	0	0
			117	117		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN



• Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE BETA CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	145.50Å 145.50Å 145.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.25 25.72 – 2.27	Depositor EDS
% Data completeness (in resolution range)	86.8 (20.00-2.25) 86.8 (25.72-2.27)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.26Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.176 , 0.218 0.270 , 0.295	Depositor DCC
R_{free} test set	932 reflections (4.49%)	DCC
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 27.0	EDS
Estimated twinning fraction	0.035 for -l,-k,-h	Xtriage
L-test for twinning	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 23933 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	3712	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1638	0.63	0/2223
2	B	0.33	0/1981	0.69	2/2691 (0.1%)
All	All	0.32	0/3619	0.66	2/4914 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	451	ASN	N-CA-C	-6.40	93.71	111.00
2	B	452	ARG	N-CA-C	-6.02	94.74	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1548	43	0
2	B	1926	0	1857	49	0
3	B	1	0	0	0	0
4	A	66	0	0	1	0
4	B	117	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3712	0	3405	85	0

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 12.

All (85) close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
2:B:369:LYS:H	2:B:422:ASN:HD22	1.21	0.84
2:B:410:HIS:HD2	2:B:412:ASN:HB2	1.44	0.80
2:B:369:LYS:H	2:B:422:ASN:ND2	1.86	0.74
1:A:64:ARG:HD3	1:A:99:PHE:O	1.88	0.74
1:A:37:ASN:HB3	1:A:95:THR:HB	1.73	0.70
1:A:68:ILE:HD12	1:A:130:ILE:HD13	1.72	0.69
2:B:428:ARG:C	2:B:429:MET:HG3	2.15	0.66
1:A:28:ILE:HG22	1:A:30:VAL:HG23	1.79	0.65
2:B:453:ILE:HD12	2:B:454:ASN:H	1.64	0.63
1:A:133:ARG:HG2	2:B:326:THR:HG21	1.81	0.61
1:A:5:LEU:HD12	1:A:5:LEU:N	2.17	0.60
1:A:95:THR:HG22	1:A:106:PHE:CZ	2.37	0.59
2:B:403:ASN:HD22	2:B:403:ASN:C	2.05	0.59
2:B:376:GLU:OE1	2:B:519:ARG:NH2	2.36	0.59
2:B:309:ALA:HB3	2:B:387:GLN:OE1	2.03	0.59
1:A:21:LEU:HD12	1:A:67:LEU:HD22	1.85	0.58
2:B:453:ILE:HD12	2:B:454:ASN:N	2.19	0.57
1:A:68:ILE:HD12	1:A:130:ILE:CD1	2.34	0.57
1:A:67:LEU:HD21	1:A:94:ARG:HD2	1.84	0.57
2:B:315:ASP:HB3	2:B:500:GLN:NE2	2.21	0.56
2:B:428:ARG:O	2:B:429:MET:HG3	2.07	0.54
1:A:177:ARG:O	1:A:178:ASP:HB3	2.08	0.54
2:B:387:GLN:HG2	2:B:529:GLY:N	2.23	0.54
2:B:403:ASN:ND2	2:B:405:SER:H	2.07	0.53
2:B:486:ILE:HG23	2:B:492:LEU:CD2	2.39	0.53
1:A:69:GLU:OE2	1:A:94:ARG:HD3	2.08	0.52
1:A:9:PRO:HG2	2:B:504:LEU:HD21	1.91	0.52
1:A:39:LEU:HD11	1:A:93:GLY:HA3	1.90	0.52
2:B:315:ASP:HB3	2:B:500:GLN:HE21	1.75	0.52
2:B:432:ASP:OD2	2:B:436:TYR:HB3	2.10	0.51
1:A:131:PHE:O	1:A:132:ALA:HB2	2.10	0.51
1:A:16:TYR:HD1	1:A:19:ILE:HD12	1.76	0.51
1:A:130:ILE:HD11	1:A:187:ILE:HD13	1.93	0.50
1:A:40:VAL:HG22	1:A:109:ILE:HD13	1.93	0.50
2:B:400:TRP:HA	2:B:425:GLY:O	2.12	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:194:GLU:OE1	2:B:333:LYS:HG3	2.12	0.49
2:B:410:HIS:CD2	2:B:412:ASN:HB2	2.35	0.48
2:B:467:ALA:HB3	2:B:472:GLN:NE2	2.29	0.47
2:B:429:MET:HE2	2:B:439:PHE:HB3	1.95	0.47
1:A:19:ILE:HG23	1:A:26:ALA:CB	2.45	0.47
2:B:507:LEU:N	2:B:507:LEU:HD22	2.30	0.47
1:A:130:ILE:HD11	1:A:187:ILE:CD1	2.45	0.46
1:A:108:THR:OG1	1:A:109:ILE:N	2.49	0.46
2:B:448:PRO:HD3	2:B:456:TRP:CZ3	2.51	0.46
2:B:403:ASN:C	2:B:403:ASN:ND2	2.69	0.46
2:B:429:MET:HE1	2:B:439:PHE:CD1	2.51	0.45
2:B:490:PRO:HA	2:B:493:LYS:HZ2	1.81	0.45
1:A:175:GLU:OE1	1:A:177:ARG:NH2	2.50	0.45
2:B:403:ASN:ND2	2:B:407:ARG:H	2.15	0.45
1:A:118:LYS:HE3	4:A:851:HOH:O	2.17	0.44
2:B:510:LYS:HA	2:B:513:PHE:CE1	2.52	0.44
2:B:370:ASP:H	2:B:422:ASN:HD21	1.66	0.44
1:A:21:LEU:HD12	1:A:67:LEU:CD2	2.46	0.44
1:A:196:VAL:HG13	2:B:337:ILE:HD12	2.00	0.44
1:A:177:ARG:HD2	1:A:177:ARG:N	2.33	0.43
1:A:30:VAL:HG22	2:B:367:TYR:CD1	2.54	0.43
2:B:495:ILE:HA	2:B:496:PRO:HD3	1.87	0.43
1:A:55:VAL:O	1:A:62:PRO:HA	2.18	0.43
2:B:312:ASN:C	2:B:312:ASN:HD22	2.22	0.43
1:A:98:ASP:OD2	1:A:101:THR:HG23	2.19	0.43
1:A:128:LEU:HD23	1:A:129:ILE:N	2.33	0.43
2:B:468:ASP:H	2:B:472:GLN:HE22	1.66	0.43
1:A:11:GLN:HE22	2:B:492:LEU:HA	1.84	0.42
2:B:368:ALA:HB1	2:B:371:GLY:O	2.19	0.42
2:B:410:HIS:CD2	2:B:412:ASN:H	2.37	0.42
2:B:441:THR:OG1	2:B:442:ILE:N	2.53	0.42
2:B:363:LEU:N	2:B:363:LEU:HD12	2.35	0.42
1:A:38:ASN:ND2	1:A:40:VAL:H	2.17	0.42
1:A:137:ILE:HD11	2:B:317:PRO:O	2.20	0.41
1:A:143:VAL:HG22	1:A:185:PHE:CE2	2.54	0.41
1:A:129:ILE:HD13	1:A:131:PHE:CZ	2.55	0.41
1:A:155:ASP:HA	1:A:156:PRO:HD3	1.79	0.41
2:B:343:LEU:HA	2:B:346:VAL:HG22	2.02	0.41
2:B:510:LYS:HA	2:B:513:PHE:CD1	2.55	0.41
1:A:188:ARG:HH11	1:A:188:ARG:HG3	1.85	0.41
1:A:79:TYR:O	1:A:81:SER:N	2.51	0.41
2:B:469:GLY:H	2:B:472:GLN:NE2	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Distance(Å)	Clash(Å)
1:A:155:ASP:O	1:A:159:ASN:ND2	2.54	0.41
2:B:387:GLN:HG2	2:B:529:GLY:H	1.84	0.40
1:A:48:ARG:NH1	1:A:109:ILE:HD11	2.36	0.40
2:B:400:TRP:CE2	2:B:462:HIS:CB	3.04	0.40
2:B:489:CYS:HA	2:B:490:PRO:HD3	1.83	0.40
1:A:53:GLY:HA3	1:A:185:PHE:O	2.21	0.40
1:A:5:LEU:CD1	1:A:5:LEU:N	2.83	0.40
2:B:353:ALA:HA	2:B:356:PHE:HD2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/209 (96%)	191 (96%)	8 (4%)	1 (0%)	38	38
2	B	236/241 (98%)	221 (94%)	15 (6%)	0	100	100
All	All	436/450 (97%)	412 (94%)	23 (5%)	1 (0%)	56	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/175 (97%)	167 (99%)	2 (1%)	82	89

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	201/204 (98%)	195 (97%)	6 (3%)	53	62
All	All	370/379 (98%)	362 (98%)	8 (2%)	64	74

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	177	ARG
2	B	312	ASN
2	B	333	LYS
2	B	381	HIS
2	B	403	ASN
2	B	412	ASN
2	B	429	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	18	HIS
1	A	33	HIS
1	A	38	ASN
1	A	72	GLN
2	B	312	ASN
2	B	403	ASN
2	B	410	HIS
2	B	412	ASN
2	B	422	ASN
2	B	472	GLN
2	B	499	GLN

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/209 (96%)	1.32	32 (15%) 3 2	9, 24, 47, 70	0
2	B	238/241 (98%)	0.68	11 (4%) 31 35	9, 15, 37, 47	0
All	All	440/450 (97%)	0.97	43 (9%) 8 9	9, 18, 41, 70	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	PHE	6.4
1	A	178	ASP	5.6
2	B	303	ILE	5.3
1	A	46	GLY	4.0
1	A	87	GLY	3.2
1	A	177	ARG	3.2
1	A	26	ALA	3.2
1	A	173	LYS	3.1
1	A	102	GLY	3.1
2	B	370	ASP	3.0
2	B	538	ASP	2.9
1	A	179	GLY	2.9
1	A	88	LYS	2.9
1	A	43	ASN	2.9
1	A	44	THR	2.8
1	A	42	ASP	2.8
1	A	33	HIS	2.7
1	A	88(B)	VAL	2.7
2	B	355	LYS	2.4
1	A	100	GLY	2.4
1	A	118	LYS	2.4
2	B	470	TRP	2.4
1	A	128	LEU	2.3
1	A	104	TRP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	353	ALA	2.3
2	B	412	ASN	2.2
1	A	30	VAL	2.2
1	A	71	TRP	2.2
2	B	410	HIS	2.2
1	A	97	ALA	2.2
2	B	367	TYR	2.2
2	B	356	PHE	2.1
1	A	85	THR	2.1
1	A	183	TYR	2.1
1	A	61	LEU	2.1
1	A	101	THR	2.1
1	A	45	GLN	2.1
1	A	70	ILE	2.1
2	B	441	THR	2.1
1	A	149	ALA	2.1
1	A	82	GLN	2.1
1	A	67	LEU	2.0
1	A	119	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FE	B	600	1/1	0.10	-	14,14,14,14	0

6.5 Other polymers ⓘ

There are no such residues in this entry.