



wwPDB X-ray Structure Validation Summary Report

Feb 26, 2014 – 11:53 PM GMT

PDB ID : 1EVW
Title : L116A MUTANT OF THE HOMING ENDONUCLEASE I-PPOI COM-
PLEXED TO HOMING SITE DNA.
Authors : Galburt, E.A.; Jurica, M.S.; Chevalier, B.S.; Erho, D.; Stoddard, B.L.
Deposited on : 2000-04-20
Resolution : 3.10 Å(reported)

This is a wwPDB validation summary report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at <http://wwpdb.org/ValidationPDFNotes.html>


The following versions of software and data (see [references](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.15 2013
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	21963
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP)	:	stable22683

1 Overall quality at a glance

The reported resolution of this entry is 3.10 Å.













Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	79885	1078 (3.16-3.04)
Ramachandran outliers	78287	1044 (3.16-3.04)
Sidechain outliers	78261	1044 (3.16-3.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	E	12	
1	G	12	
2	M	8	
2	N	8	
3	F	12	
3	H	12	
4	O	8	
4	P	8	
5	A	163	
5	B	163	
5	C	163	
5	D	163	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6612 atoms, of which 0 are hydrogen and 0 are deuterium.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*GP*GP*CP*TP*AP*CP*CP*TP*TP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	12	Total	C	N	O	P	0	0	0
			241	117	42	71	11			
1	G	12	Total	C	N	O	P	0	0	0
			241	117	42	71	11			

- Molecule 2 is a DNA chain called DNA (5'-D(P*GP*AP*GP*AP*GP*TP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	M	8	Total	C	N	O	P	0	0	0
			169	79	35	47	8			
2	N	8	Total	C	N	O	P	0	0	0
			169	79	35	47	8			

- Molecule 3 is a DNA chain called DNA (5'-D(*TP*GP*AP*CP*TP*CP*TP*CP*TP*TP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	12	Total	C	N	O	P	0	0	0
			239	117	39	72	11			
3	H	12	Total	C	N	O	P	0	0	0
			239	117	39	72	11			

- Molecule 4 is a DNA chain called DNA (5'-D(P*GP*GP*TP*AP*GP*CP*CP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	O	8	Total	C	N	O	P	0	0	0
			167	78	33	48	8			
4	P	8	Total	C	N	O	P	0	0	0
			167	78	33	48	8			

- Molecule 5 is a protein called I-PPOI HOMING ENDONUCLEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	A	162	Total	C	N	O	S	0	0	0
			1242	783	232	219	8			
5	B	162	Total	C	N	O	S	0	0	0
			1242	783	232	219	8			
5	C	162	Total	C	N	O	S	0	0	0
			1242	783	232	219	8			
5	D	162	Total	C	N	O	S	0	0	0
			1242	783	232	219	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	116	ALA	LEU	engineered	UNP Q94702
B	116	ALA	LEU	engineered	UNP Q94702
C	116	ALA	LEU	engineered	UNP Q94702
D	116	ALA	LEU	engineered	UNP Q94702

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	2	Total	Zn	0	0
			2	2		
6	A	2	Total	Zn	0	0
			2	2		
6	D	2	Total	Zn	0	0
			2	2		
6	C	2	Total	Zn	0	0
			2	2		

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	P	1	Total	Mg	0	0
			1	1		
7	B	1	Total	Mg	0	0
			1	1		
7	A	1	Total	Mg	0	0
			1	1		
7	N	1	Total	Mg	0	0
			1	1		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: DNA (5'-D(*TP*GP*GP*CP*TP*AP*CP*CP*TP*TP*AP*A)-3')

Chain E: 



- Molecule 1: DNA (5'-D(*TP*GP*GP*CP*TP*AP*CP*CP*TP*TP*AP*A)-3')

Chain G: 




- Molecule 2: DNA (5'-D(P*GP*AP*GP*AP*GP*TP*CP*A)-3')

Chain M: 




- Molecule 2: DNA (5'-D(P*GP*AP*GP*AP*GP*TP*CP*A)-3')

Chain N: 




- Molecule 3: DNA (5'-D(*TP*GP*AP*CP*TP*CP*TP*CP*TP*TP*AP*A)-3')

Chain F: 




- Molecule 3: DNA (5'-D(*TP*GP*AP*CP*TP*CP*TP*CP*TP*TP*AP*A)-3')

Chain H: 



- Molecule 4: DNA (5'-D(P*GP*GP*TP*AP*GP*CP*CP*A)-3')

Chain O:




- Molecule 4: DNA (5'-D(P*GP*GP*TP*AP*GP*CP*CP*A)-3')

Chain P:




- Molecule 5: I-PPOI HOMING ENDONUCLEASE

Chain A:




- Molecule 5: I-PPOI HOMING ENDONUCLEASE

Chain B:




- Molecule 5: I-PPOI HOMING ENDONUCLEASE

Chain C:



- Molecule 5: I-PPOI HOMING ENDONUCLEASE

Chain D:



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	182.00Å 73.10Å 92.70Å 90.00° 95.40° 90.00°	Depositor
Resolution (Å)	50.00 – 3.10	Depositor
% Data completeness (in resolution range)	(Not available) (50.00-3.10)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.273 , 0.320	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6612	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	E	0.48	0/269	0.74	0/413
1	G	0.61	0/269	0.95	0/413
2	M	0.81	1/190 (0.5%)	0.74	0/290
2	N	0.98	1/190 (0.5%)	0.91	0/290
3	F	0.50	0/266	0.78	0/408
3	H	0.56	0/266	0.79	0/408
4	O	0.97	1/187 (0.5%)	0.99	0/285
4	P	0.95	2/187 (1.1%)	1.10	1/285 (0.4%)
5	A	0.56	0/1283	0.75	2/1756 (0.1%)
5	B	0.63	2/1283 (0.2%)	0.78	2/1756 (0.1%)
5	C	0.55	0/1283	0.68	0/1756
5	D	0.64	3/1283 (0.2%)	0.79	3/1756 (0.2%)
All	All	0.63	10/6956 (0.1%)	0.79	8/9816 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
2	M	0	1
3	H	0	1
4	O	0	3
4	P	0	4
5	C	0	1
All	All	0	11

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	N	14	DG	OP3-P	-7.69	1.51	1.61
4	O	13	DG	OP3-P	-7.42	1.52	1.61
2	M	14	DG	OP3-P	-7.30	1.52	1.61
4	P	13	DG	OP3-P	-7.26	1.52	1.61
5	D	71	VAL	CB-CG2	-6.26	1.39	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	74	ARG	NE-CZ-NH2	8.77	124.68	120.30
5	D	157	ARG	CG-CD-NE	-8.76	93.40	111.80
5	B	139	LEU	CA-CB-CG	7.25	131.98	115.30
5	B	21	GLN	CA-CB-CG	-7.05	97.88	113.40
4	P	13	DG	OP1-P-OP2	-6.78	109.42	119.60

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	7	DA	Sidechain
2	M	17	DA	Sidechain
4	O	14	DG	Sidechain
4	O	15	DT	Sidechain
4	O	19	DC	Sidechain

5.2 Close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogens added by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, and the number in parentheses is this value normalized per 1000 atoms of the molecule in the chain. The Symm-Clashes column gives symmetry related clashes, in the same way as for the Clashes column.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	241	0	138	20	0
1	G	241	0	138	25	0
2	M	169	0	90	19	0
2	N	169	0	90	14	0
3	F	239	0	139	6	0
3	H	239	0	139	11	0
4	O	167	0	90	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	P	167	0	90	21	0
5	A	1242	0	1185	57	0
5	B	1242	0	1186	62	2
5	C	1242	0	1185	74	2
5	D	1242	0	1185	71	0
6	A	2	0	0	0	0
6	B	2	0	0	0	0
6	C	2	0	0	0	0
6	D	2	0	0	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
7	N	1	0	0	0	0
7	P	1	0	0	0	0
All	All	6612	0	5655	306	2

Clashscore is defined as the number of clashes calculated for the entry per 1000 atoms (including hydrogens) of the entry. The overall clashscore for this entry is 25.

The worst 5 of 306 close contacts within the same asymmetric unit are listed below.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:C:9:LEU:HD23	5:D:157:ARG:HH21	1.11	1.11
1:E:13:DA:H2'	2:M:14:DG:C8	1.87	1.10
2:M:14:DG:H5''	5:B:96:ALA:O	1.51	1.08
5:D:157:ARG:NH1	5:D:157:ARG:HG2	1.68	1.04
2:M:14:DG:H5'	5:B:119:ASN:ND2	1.78	0.98

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Distance(Å)	Clash(Å)
5:B:28:HIS:ND1	5:C:21:GLN:NE2[1_545]	1.94	0.26
5:B:28:HIS:ND1	5:C:21:GLN:CD[1_545]	2.17	0.03

5.3 Torsion angles

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	A	160/163 (98%)	146 (91%)	12 (8%)	2 (1%)	18	60
5	B	160/163 (98%)	147 (92%)	10 (6%)	3 (2%)	12	51
5	C	160/163 (98%)	144 (90%)	14 (9%)	2 (1%)	18	60
5	D	160/163 (98%)	140 (88%)	18 (11%)	2 (1%)	18	60
All	All	640/652 (98%)	577 (90%)	54 (8%)	9 (1%)	16	58

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	A	77	SER
5	C	77	SER
5	C	159	SER
5	A	137	VAL
5	B	77	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution. The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
5	A	132/133 (99%)	114 (86%)	18 (14%)	5	21
5	B	132/133 (99%)	119 (90%)	13 (10%)	12	40
5	C	132/133 (99%)	117 (89%)	15 (11%)	8	31
5	D	132/133 (99%)	113 (86%)	19 (14%)	5	19
All	All	528/532 (99%)	463 (88%)	65 (12%)	7	26

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	B	144	LEU
5	C	61	ARG
5	D	129	ASN
5	B	155	GLN
5	C	26	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
5	C	63	GLN
5	C	70	GLN
5	D	63	GLN
5	B	119	ASN
5	C	101	HIS

5.3.3 RNA ⓘ

There are no RNA chains in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.